

1 Minimizing complex functions with Wirtinger derivatives

1.1 Definition

Define for $z = x + iy \in \mathbb{C}$ with $x, y \in \mathbb{R}$

$$\frac{\partial}{\partial z} \equiv \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) \quad \& \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) \quad (1)$$

for functions of one complex variable, and

$$\frac{\partial}{\partial z_j} = \frac{1}{2} \left(\frac{\partial}{\partial x_j} - i \frac{\partial}{\partial y_j} \right) \quad \& \quad \frac{\partial}{\partial \bar{z}_j} = \frac{1}{2} \left(\frac{\partial}{\partial x_j} + i \frac{\partial}{\partial y_j} \right) \quad (2)$$

for functions of multiple complex variables.

1.2 Agreement with the complex derivative

If f is complex differentiable at a point (i.e. the limit $f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$ exists), then $\frac{\partial f}{\partial z}$ agrees with the complex derivative $\frac{df}{dz}$.

Proof. For $f(z) = u(z) + iv(z)$ complex differentiable,

$$\begin{aligned} \frac{\partial f}{\partial z} &\stackrel{(1)}{=} \frac{1}{2} \left(\frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} - i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \right) \\ &\stackrel{(1)}{=} \frac{\partial u}{\partial z} + i \frac{\partial v}{\partial z} \stackrel{\text{Def.}}{=} \frac{df}{dz}. \end{aligned} \quad (3)$$

□

1.3 The $\bar{z}$ -derivative and Cauchy–Riemann

Furthermore,

$$\begin{aligned} 0 &\stackrel{!}{=} \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) = \frac{1}{2} \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} + i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) \\ &\iff \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \wedge \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}, \end{aligned} \quad (4)$$

so that $0 = \frac{\partial f}{\partial \bar{z}}$ is equivalent to the Cauchy–Riemann equations, i.e. the necessary & sufficient condition for f to be complex differentiable.

1.4 Minimizing a real-valued function

Minimize a real-valued function¹

$$f: \mathbb{C} \rightarrow \mathbb{R} \tag{5}$$

$$f(z) = f(x, y), \quad (z = x + iy) \tag{6}$$

via the condition

$$\nabla f(x, y) = \begin{pmatrix} \partial_x f \\ \partial_y f \end{pmatrix} \stackrel{!}{=} 0. \tag{7}$$

Then

$$\partial_x f \stackrel{!}{=} 0 \stackrel{!}{=} \partial_y f \quad \Longleftrightarrow \quad \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) \stackrel{!}{=} 0, \tag{8}$$

so one can minimize a real-valued function by imposing $\boxed{\frac{\partial f}{\partial \bar{z}} = 0}$.

1.5 References

- *Wirtinger derivatives*, English Wikipedia.

¹Minimizing a complex-valued f does not make sense because $\mathbb{C}$ has no order.