

Solutions

1) (a) The given permutation is:

52134

The inversions are:

(5, 2), (5, 1), (5, 3), (5, 2)

(2, 1)

Thus there are five (odd) inversion, and so this is an odd permutation.

(b) The given permutation is:

45213

The inversions are:

(4, 2), (4, 1), (4, 3)

(5, 2), (5, 1), (5, 3)

(2, 1)

Here, number of permutations are:

$$3 + 3 + 1$$

$$= 7$$

which is odd, thus given permutation is odd permutation.

(c) 42135

The inversions are:

(4, 2), (4, 1), (4, 3)

(2, 1)

Here, the number of conversions are:

$$3 + 1$$

$$= 4$$

which is even

Thus, the given permutation is an even permutation.

(d) 54321

The inversions are:

$(5,4), (5,3), (5,2), (5,1)$

$(4,3), (4,2), (4,1)$

$(3,2), (3,1)$

$(2,1)$

Here, the number of inversions are:

$$4 + 3 + 2 + 1$$

$$= 10$$

which is even

Thus, the given permutation is an even permutation.

2) (a) 436215 and 416235

The number of inversions in 436215 is given by

$$3 + 2 + 3 + 1 + 0 + 0$$

$$= 9$$

The number of inversions in 416235 is given by

$$3 + 0 + 3 + 0 + 0 + 0$$

$$= 6$$

Thus, the number of inversions differ by:

$$9 - 6 = 3$$

which is odd. Hence verified.

(b) 623415 and 523416

The number of inversions in 623415 is given by

$$5 + 0 + 0 + 1 + 0 + 0$$

$$= 6$$

The number of inversions in 523416 is given by

$$4 + 1 + 1 + 1 + 0 + 0$$

$$= 7$$

Thus, the number of inversions differ by:

$$7 - 6 = 1$$

which is odd. Hence verified.

(c) 321564 and 341562

The number of inversions in 321564 is given by

$$2+1+0+1+1+0 \\ = 5$$

The number of inversions in 341562 is given by

$$2+2+0+1+1+0 \\ = 6$$

Thus, the number of inversions differ by:

$$6-5=1$$

which is odd. Hence verified.

(d) 123564 and 423561

The number of inversions in 123564 is given by

$$0+0+0+1+1+0 \\ = 2$$

The number of inversions in 423561 is given by

$$3+1+1+1+1+0 \\ = 7$$

Thus, the number of inversions differ by:

$$7-2=5$$

which is odd. Hence verified.

$$3) \text{ (a) } D = \begin{vmatrix} 1 & 3 \\ 2 & 4 \end{vmatrix}$$

$$= (1)(4) - (3)(2)$$

$$= 4 - 6$$

$$= -2$$

$$\text{(b) } D = \begin{vmatrix} 1 & 2 & 1 \\ 1 & 3 & 2 \\ 1 & 0 & 1 \end{vmatrix}$$

$$\begin{aligned}
&= (1) \begin{vmatrix} 3 & 2 \\ 0 & 1 \end{vmatrix} - (2) \begin{vmatrix} 1 & 2 \\ 1 & 1 \end{vmatrix} + (1) \begin{vmatrix} 1 & 3 \\ 1 & 0 \end{vmatrix} \\
&= (1)((3)(1) - (2)(0)) - (2)((1)(1) - (2)(1)) + (1)((1)(0) - (3)(1)) \\
&= (1)(3 - 0) - (2)(1 - 2) + (1)(0 - 3) \\
&= (1)(3) - (2)(-1) + (1)(-3) \\
&= 3 + 2 - 3 \\
&= 2
\end{aligned}$$

$$(c) D = \begin{vmatrix} -4 & 2 & 0 & 0 \\ 2 & 3 & 1 & 0 \\ 3 & 1 & 0 & 2 \\ 1 & 3 & 0 & 3 \end{vmatrix}$$

$$\begin{aligned}
&= (-4) \begin{vmatrix} 3 & 1 & 0 \\ 1 & 0 & 2 \\ 3 & 0 & 3 \end{vmatrix} - (2) \begin{vmatrix} 2 & 1 & 0 \\ 3 & 0 & 2 \\ 1 & 0 & 3 \end{vmatrix} + (0) \begin{vmatrix} 2 & 3 & 0 \\ 3 & 1 & 2 \\ 1 & 3 & 3 \end{vmatrix} - (0) \begin{vmatrix} 2 & 3 & 1 \\ 3 & 1 & 0 \\ 1 & 3 & 0 \end{vmatrix} \\
&= (-4)(3(0 - 0) - 1(3 - 6) + 0(0 - 0)) - 2(2(0 - 0) - 1(9 - 2) + 0(0 - 0)) + 0 - 0 \\
&= (-4)(0 - 1(-3) + 0) - 2(0 - 1(7) + 0) + 0 - 0 \\
&= (-4)(3) - 2(-7) \\
&= -12 + 14 \\
&= 2
\end{aligned}$$

$$(d) D = \begin{vmatrix} 4 & 2 & 2 & 0 \\ 2 & 0 & 0 & 0 \\ 3 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{vmatrix}$$

$$\begin{aligned}
&= (4) \begin{vmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{vmatrix} - (2) \begin{vmatrix} 2 & 0 & 0 \\ 3 & 0 & 1 \\ 0 & 1 & 0 \end{vmatrix} + (2) \begin{vmatrix} 2 & 0 & 0 \\ 3 & 0 & 1 \\ 0 & 0 & 0 \end{vmatrix} - (0) \begin{vmatrix} 2 & 0 & 0 \\ 3 & 0 & 0 \\ 0 & 0 & 1 \end{vmatrix}
\end{aligned}$$

If all elements of a row/column are zero, then value of determinant is zero.

$$D = (4)(0) - (2) \begin{vmatrix} 2 & 0 & 0 \\ 3 & 0 & 1 \\ 0 & 1 & 0 \end{vmatrix} + (2)(0) - (0) \begin{vmatrix} 2 & 0 & 0 \\ 3 & 0 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= 0 - (2)(2(0-1) - 0 + 0) + (0) - (0)$$

$$= -(2)(2(-1))$$

$$= -4$$

$$4) A = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix}$$

Given that,

$$\det(A) = |A| = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = 3 \quad \dots \quad (I)$$

Consider

$$\det(B) = |B| = \begin{vmatrix} a_1 + 2b_1 - 3c_1 & a_2 + 2b_2 - 3c_2 & a_3 + 2b_3 - 3c_3 \\ c_1 & c_2 & c_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

Interchanging the rows R2 and R3, sign of the determinant gets changed.

$$= - \begin{vmatrix} a_1 + 2b_1 - 3c_1 & a_2 + 2b_2 - 3c_2 & a_3 + 2b_3 - 3c_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

Splitting the determinant along R1,

$$= - \left(\begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} + \begin{vmatrix} 2b_1 & 2b_2 & 2b_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} + \begin{vmatrix} -3c_1 & -3c_2 & -3c_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \right)$$

Taking 2 common from R1 of second determinant, and -3 common from 3rd determinant.

$$= - \left(\begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} + 2 \begin{vmatrix} b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} - 3 \begin{vmatrix} c_1 & c_2 & c_3 \end{vmatrix} \right)$$

Since R1 and R2 of 2nd determinant are identical, so it is equal to zero

R1 and R3 of 3rd determinant are identical, so it is also zero.

$$= - \left(\begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} + 2(0) - 3(0) \right)$$

$$= - \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

$$= -\det(A)$$

$$= -3$$

Now, consider

$$\det(C) = \begin{vmatrix} a_1 & 3a_2 & a_3 \\ b_1 & 3b_2 & b_3 \\ c_1 & 3c_2 & c_3 \end{vmatrix}$$

Taking 3 common from C2

$$= 3 \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

$$= 3\det(A)$$

$$= 3(3)$$

$$= 9$$

Now, consider

$$\det(D) = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

$$= |A^T|$$

Since, determinants of a matrix and its transpose, are same.

$$\det(D) = |A|$$

$$= 3$$

$$5) (a) \quad D = \begin{vmatrix} 2 & 0 & 1 & 4 \\ 3 & 2 & -4 & -2 \\ 2 & 3 & -1 & 0 \\ 11 & 8 & -4 & 6 \end{vmatrix}$$

$$R_2 - R_3$$

$$D = \begin{vmatrix} 2 & 0 & 1 & 4 \\ 1 & -1 & -3 & -2 \\ 2 & 3 & -1 & 0 \\ 11 & 8 & -4 & 6 \end{vmatrix}$$

$$R_2 \leftrightarrow R_1$$

$$D = - \begin{vmatrix} 1 & -1 & -3 & -2 \\ 2 & 0 & 1 & 4 \\ 2 & 3 & -1 & 0 \\ 11 & 8 & -4 & 6 \end{vmatrix}$$

$$R_2 - 2R_1$$

$$R_3 - 2R_1$$

$$R_4 - 11R_1$$

$$D = - \begin{vmatrix} 1 & -1 & -3 & -2 \\ 0 & 2 & 7 & 8 \\ 0 & 5 & 5 & 4 \\ 0 & 19 & 29 & 28 \end{vmatrix}$$

Expand along C1

$$D = -(1) \begin{vmatrix} 2 & 7 & 8 \\ 1 & -9 & -12 \\ 19 & 29 & 28 \end{vmatrix}$$

Taking 4 common C3

$$D = -(1)(4) \begin{vmatrix} 2 & 7 & 2 \\ 1 & -9 & -3 \\ 19 & 29 & 7 \end{vmatrix}$$

$$R_1 - 2R_2$$

$$R_3 - 19R_2$$

$$D = -4 \begin{vmatrix} 0 & 25 & 8 \\ 1 & -9 & -3 \\ 0 & 200 & 64 \end{vmatrix}$$

Taking 8 common from R3

$$D = -4(8) \begin{vmatrix} 0 & 25 & 8 \\ 1 & -9 & -3 \\ 0 & 25 & 8 \end{vmatrix}$$

Since R1 and R3 are identical, so this determinant is zero

Thus given determinant is zero.

$$(b) D = \begin{vmatrix} 4 & 2 & 3 & -4 \\ 3 & -2 & 1 & 5 \\ -2 & 0 & 1 & -3 \\ 8 & -2 & 6 & 4 \end{vmatrix}$$

Taking 2 common from R4

$$D = 2 \begin{vmatrix} 4 & 2 & 3 & -4 \\ 3 & -2 & 1 & 5 \\ -2 & 0 & 1 & -3 \\ 4 & -1 & 3 & 2 \end{vmatrix}$$

$$R_1 + 2R_4$$

$$R_2 - 2R_4$$

$$D = 2 \begin{vmatrix} 12 & 0 & 9 & 0 \\ -5 & 0 & -5 & 1 \\ -2 & 0 & 1 & -3 \\ 4 & -1 & 3 & 2 \end{vmatrix}$$

Expand along C2

$$D = 2(-1) \begin{vmatrix} 12 & 9 & 0 \\ -5 & -5 & 1 \\ -2 & 1 & -3 \end{vmatrix}$$

Taking 3 common from R1

$$D = 2(-1)(3) \begin{vmatrix} 4 & 3 & 0 \\ -5 & -5 & 1 \\ -2 & 1 & -3 \end{vmatrix}$$

$R_3 + 3R_1$

$$D = 2(-1)(3) \begin{vmatrix} 4 & 3 & 0 \\ -5 & -5 & 1 \\ 13 & 16 & 0 \end{vmatrix}$$

Expand along C3

$$D = 2(-1)(3)(-1) \begin{vmatrix} 4 & 3 \\ 13 & 16 \end{vmatrix}$$

$$D = 2(-1)(3)(-1)(64 - 39)$$

$$D = 6(25)$$

$$D = 150$$

$$(c) D = \begin{vmatrix} 2 & 0 & 0 & 0 \\ -5 & 3 & 0 & 0 \\ 3 & 2 & 4 & 0 \\ 4 & 2 & 1 & -5 \end{vmatrix}$$

The given determinant is in Triangular form, i.e. all the entries above the diagonal are zero

Thus, Determinant is given by multiplying the diagonal entries only

$$D = (2)(3)(4)(-5)$$

$$D = -120$$

6) Consider,

$$D_n = \begin{vmatrix} 1 & \alpha_1 & \alpha_1^2 & & \alpha_1^{n-2} & \alpha_1^{n-1} \\ 1 & \alpha_2 & \alpha_2^2 & \dots & \alpha_2^{n-2} & \alpha_2^{n-1} \\ 1 & \alpha_3 & \alpha_3^2 & & \alpha_3^{n-2} & \alpha_3^{n-1} \\ & \vdots & & \ddots & & \vdots \\ 1 & \alpha_{n-1} & \alpha_{n-1}^2 & \dots & \alpha_{n-1}^{n-2} & \alpha_{n-1}^{n-1} \\ 1 & \alpha_n & \alpha_n^2 & & \alpha_n^{n-2} & \alpha_n^{n-1} \end{vmatrix}$$

Subtract R1 from each of the other rows and leave Dn unchanged:

$$D_n = \begin{vmatrix} 1 & \alpha_1 & \alpha_1^2 & & \alpha_1^{n-2} & \alpha_1^{n-1} \\ 0 & \alpha_2 - \alpha_1 & \alpha_2^2 - \alpha_1^2 & \dots & \alpha_2^{n-2} - \alpha_1^{n-2} & \alpha_2^{n-1} - \alpha_1^{n-1} \\ 0 & \alpha_3 - \alpha_1 & \alpha_3^2 - \alpha_1^2 & & \alpha_3^{n-2} - \alpha_1^{n-2} & \alpha_3^{n-1} - \alpha_1^{n-1} \\ & \vdots & & \ddots & & \vdots \\ 0 & \alpha_{n-1} - \alpha_1 & \alpha_{n-1}^2 - \alpha_1^2 & \dots & \alpha_{n-1}^{n-2} - \alpha_1^{n-2} & \alpha_{n-1}^{n-1} - \alpha_1^{n-1} \\ 0 & \alpha_n - \alpha_1 & \alpha_n^2 - \alpha_1^2 & & \alpha_n^{n-2} - \alpha_1^{n-2} & \alpha_n^{n-1} - \alpha_1^{n-1} \end{vmatrix}$$

Subtract in order, α_1 times C-(n-1) from C-(n),

$$\begin{aligned} & C_n - \alpha_1 C_{n-1} \\ & C_{n-1} - \alpha_1 C_{n-2} \\ & \dots \\ & C_2 - \alpha_1 C_1 \end{aligned}$$

$$D_n = \begin{vmatrix} 1 & 0 & 0 & & 0 & 0 \\ 0 & \alpha_2 - \alpha_1 & (\alpha_2 - \alpha_1)\alpha_2 & \dots & (\alpha_2 - \alpha_1)\alpha_2^{n-3} & (\alpha_2 - \alpha_1)\alpha_2^{n-2} \\ 0 & \alpha_3 - \alpha_1 & (\alpha_3 - \alpha_1)\alpha_3 & & (\alpha_3 - \alpha_1)\alpha_3^{n-3} & (\alpha_3 - \alpha_1)\alpha_3^{n-2} \\ & \vdots & & \ddots & & \vdots \\ 0 & \alpha_{n-1} - \alpha_1 & (\alpha_{n-1} - \alpha_1)\alpha_{n-1} & \dots & (\alpha_{n-1} - \alpha_1)\alpha_{n-1}^{n-3} & (\alpha_{n-1} - \alpha_1)\alpha_{n-1}^{n-2} \\ 0 & \alpha_n - \alpha_1 & (\alpha_n - \alpha_1)\alpha_n & & (\alpha_n - \alpha_1)\alpha_n^{n-3} & (\alpha_n - \alpha_1)\alpha_n^{n-2} \end{vmatrix}$$

From (k)th row, extracting $(\alpha_k - \alpha_1)$ as factors, we get,

$$D_n = \prod_{k=2}^n (\alpha_k - \alpha_1) \begin{vmatrix} 1 & 0 & 0 & & 0 & 0 \\ 0 & 1 & \alpha_2 & \dots & \alpha_2^{n-3} & \alpha_2^{n-2} \\ 0 & 1 & \alpha_3 & & \alpha_3^{n-3} & \alpha_3^{n-2} \\ & \vdots & & \ddots & & \vdots \\ 0 & 1 & \alpha_{n-1} & \dots & \alpha_{n-1}^{n-3} & \alpha_{n-1}^{n-2} \\ 0 & 1 & \alpha_n & & \alpha_n^{n-3} & \alpha_n^{n-2} \end{vmatrix}$$

Expand along C1

$$D_n = \prod_{k=2}^n (\alpha_k - \alpha_1) \begin{vmatrix} 1 & \alpha_2 & & \alpha_2^{n-3} & \alpha_2^{n-2} \\ 1 & \alpha_3 & \dots & \alpha_3^{n-3} & \alpha_3^{n-2} \\ & \vdots & \ddots & & \vdots \\ 1 & \alpha_{n-1} & & \alpha_{n-1}^{n-3} & \alpha_{n-1}^{n-2} \\ 1 & \alpha_n & & \alpha_n^{n-3} & \alpha_n^{n-2} \end{vmatrix}$$

It can be easily seen,

$$D_n = \prod_{k=2}^n (\alpha_k - \alpha_1) D_{n-1}$$

$$\Rightarrow D_n = \prod_{k=2}^n (\alpha_k - \alpha_1) \prod_{m=2}^n (\alpha_m - \alpha_2) D_{n-2}$$

and so on,...

$$\text{Thus, } \Rightarrow D_n = \prod_{1 \leq i < j \leq n} (\alpha_j - \alpha_i)$$

Hence proved.