

University College Dublin An Coláiste Ollscoile, Baile Átha Cliath

SEMESTER I EXAMINATION - 2009

CHEM 10030 Chemistry For Engineers

Professor Powell

Professor Wayne

Professor Waghorne

Dr. Sullivan *

Dr. Quinn

Time Allowed: 2 hours

Instructions for Candidates

All questions carry equal marks; the approximate assignment of marks to parts of a question is indicated (as a percentage) in parentheses.

Use a separate answer book provided for each question.

No loose rough work sheets are to be used. The rough work for each question should be included in the answer book for that question.

Instructions for Invigilators

The use of electronic calculators is permitted
A Periodic Table of the Elements is attached to these sheets
Graph paper should be provided

Page 1 of 7

Answer both questions 1 and 2

Question 1

Answer any *four* of the following (a) - (f)

- (a) Answer both sections (i) and (ii)
 - (i) Draw a schematic diagram of the cathode-ray tube apparatus used by Thompson and explain how his experiments furthered the development of atomic theory; (15)
 - (ii) Draw a schematic diagram of the experiment that confirmed the existence of the nucleus and label the individual components. (10)

- (b) Answer **both** sections (i) and (ii)
 - (i) Explain the principle of operation of a mass spectrometer;
 - (ii) An element exists as one of four possible isotopes. 83.7% of all atoms of the element have a relative atomic mass of 51.94, 4.3% have a mass of 49.95, 9.5% have a mass of 52.94 and the remainder has a relative atomic mass of 53.94. Identify the element and determine its average relative atomic mass.

 (10)

Continued.....

(c)	Answer bo	and sections (1) and (11)	
	(i)	Draw and label (using the graph paper provided) a potential energy diagram showing PE as a function of inter atomic distance for two Br atoms given that the bond strength of a Br–Br bond is 193 kJ mol ⁻¹ and the inter nuclear distance is 229 pm.	(15)
	(ii)	Explain the magnitude of the PE when the two Br nuclei are (a) 1 m, (b) 229 pm and (c) 25 pm apart.	(10)
(d)	Answer al	l sections	
	you show	is structures and VSEPR models of the following molecules. Ensure that electron counts, discriminate between lone pairs and bonding pairs of and state the final shapes of the molecules.	
		BF_3	(5)
		${ m SF_6} \ { m C_2H_4}$	(5) (5)
		H_2O	(5)
		PCl_5	(5)
(e)	Answer bo	oth sections (i) and (ii)	
	(i)	Explain why heat is given out when a hydrocarbon reacts with O_2 to form CO_2 and H_2O .	(15)
	(ii)	Determine the mass of CO_2 and the number of moles of H_2O formed when $20 kg$ of octane (C_8H_{18}) is fully burned.	(10)
	S	ection (f) is on the following page Continued	

- (f) Answer both sections (i) and (ii)
 - (i) Explain the appearance of the hydrogen emission spectrum using the Bohr model of the atom. (15)
 - (ii) Determine the wavelength of the photon emitted when an electron in a H atom falls from the n=5 to the n=2 level. In what region of the electromagnetic spectrum would light of this wavelength be found?

Planck's constant,
$$h = 6.626 \times 10^{-34} \text{ J s.}$$

Rydberg constant, $R = 2.18 \times 10^{-18} \text{ J.}$
Speed of light, $c = 3 \times 10^8 \text{ m s}^{-1}$. (10)

Question 2 is on the next page......

Question 2

Answer all parts (a)-(c)

- (a) Answer all parts (i), (ii) and (iii)
 - (i) State Hess's Law;
 - (ii) Define the standard state of an element;
 - (iii) Given the following information:

$$\begin{array}{lll} {\rm TiO_{2}(s)} \, + \, 4 \, {\rm HCl(g)} \, \to \, {\rm TiCl_{4}(l)} \, + \, 2 \, {\rm H_{2}O(l)} \, \, \Delta_{\rm r} H^{\ominus} = -22 \, {\rm kJ \; mol^{-1}} \\ {\rm Ti(s)} \, + \, 2 \, {\rm Cl_{2}(g)} \, \to \, {\rm TiCl_{4}(l)} \, & \Delta_{\rm r} H^{\ominus} = \, -763 \, {\rm kJ \; mol^{-1}} \\ {\rm H_{2}O(l)} \, \to \, {\rm H_{2}(g)} \, + \, {}^{1}\!\!\!/_{2} \, {\rm O_{2}(g)} \, & \Delta_{\rm r} H^{\ominus} = \, +286 \, {\rm kJ \; mol^{-1}} \\ {}^{1}\!\!\!/_{2} \, {\rm H_{2}(g)} \, + \, {}^{1}\!\!\!/_{2} \, {\rm Cl_{2}(g)} \, \to \, {\rm HCl(g)} \, & \Delta_{\rm r} H^{\ominus} = \, -92 \, {\rm kJ \; mol^{-1}} \end{array}$$

Calculate the standard enthalpy of formation of TiO₂;

(b) For the exothermic reaction:

$$2 \text{ HI}(g) \Rightarrow I_2(g) + H_2(g)$$

Write an expression for the equilibrium constant in terms of concentrations.

When 0.2 moles of pure HI is added to a 2 L flask at 726 K it was found that 0.156 moles of HI remained when equilibrium was reached.

What are the equilibrium concentrations of I_2 and H_2 ?

Calculate the equilibrium constant.

What effect would increasing the reaction temperature have on the amount of I_2 at equilibrium? Explain your reasoning. (33.3)

Continued.....

- (c) Answer all parts (i), (ii) and (iii).
 - (i) Explain briefly the meaning of the following terms in chemical kinetics:

 Rate equation Activation energy Reaction half-life
 - (ii) The radioactive isotope 123 I breaks down in a first-order process with a half-life of 4.7×10^4 s at 25 °C. Calculate the rate constant for the decay.
 - (iii) Using a reaction-energy diagram briefly explain how a catalyst increases the rate of reaction.

(33.3)

oOo

Periodic Table of the Elements

1 1 2 2 2 2 2 2 2 2																		
1	18 VIII	2 Helium He 4.0026	10 Neon Ne	20.18	Agou	39.948	36 Krypton	ż	83.80	54 Xenon	Xe	131.29	Radon	R	222			
1	47 IN		9 Fluorine F	18.9984	ဦး ၁ ဦ	35.453	35 Bromine	ģ	79.904	53 Iodine		126.904	Astatine	¥	210			
1 1 1 1 1 1 1 1 1 1	16 12		oxygen Oxygen	1		32.066	Selenium	တိ				127.60	Polonium	Ъ	209			
1	5 >		7 Nitrogen	14.0067					74.922	51 Antimony	dS S	121.75	Bismuth	洒	208.98			
1 1 1 1 1 1 1 1 1 1	4 >	,	و Carbon			28.086	32 Germanium	ලී				118.71	Lead	Pp	207.19			
1 Group (old notation) Group (old notat	≡ 13		5 Boron B	10.811	Auminium Auminium				69.723	49 Indium	_	114.82	Thallium	F	204.38			
1					12	c c	Zinc	Zn	65.39	48 Cadmium	ਲ	112.411	Mercury	Нg	200.59			
1					7	9	Copper	ರ	63.546	47 Silver	Ag	107.868	e plos	Au	196.967			
1 1 1 1 1 1 1 1 1 1					10	90	Z8 Nickel	Z	58.69	46 Palladium	Pd	106.42	Platinum	置	195.09			
1 1 1 1 1 1 1 1 1 1	ation) ition)	nber SS			O		2/ Cobalt	රි	58.933	45 Rhodium	格	102.905	Iridium	<u>_</u>	192.22	109 Meifnerium	Mt	268
1	o (new nota p (old nota	Atomic nun Name Symb			∞	90	8 <u>I</u> o	Fe	55.847	44 Ruthenium	Ru	101.07	Osmium	SO	190.2	108 Hassium	£	269
2 Beryllium Beryllium 12 Bae 9,0122 124.305 20 Calcium Ca Scandium 74 87.62 88.905 56 88.905 57 Barlum 137.34 138.91 88 Radium Adrinium Rut Ra AC 226 227 24.305 38 39 Srontium 7118.91 88 88 89 Radium Adrinium Rut Ra AC 226 227	Group	Name Mass				30	25 Manganese	M⊓	54.938	43 Technetium	T _C	88 14	Rhenium	Re	186.207	107 Bohrium	В	264
2 Beryllium Beryllium 12 Bae 9,0122 124.305 20 Calcium Ca Scandium 74 87.62 88.905 56 88.905 57 Barlum 137.34 138.91 88 Radium Adrinium Rut Ra AC 226 227 24.305 38 39 Srontium 7118.91 88 88 89 Radium Adrinium Rut Ra AC 226 227					φ	à	24 Chromium	ර්	51.996	42 Molybdenum	ω	95.94	Tungsten	≥	183.85	106 Seaboroium	Sq	266
2 Beryllium Beryllium 12 Bae 9,0122 124.305 20 Calcium Ca Scandium 74 87.62 88.905 56 88.905 57 Barlum 137.34 138.91 88 Radium Adrinium Rut Ra AC 226 227 24.305 38 39 Srontium 7118.91 88 88 89 Radium Adrinium Rut Ra AC 226 227					S	90	23 Vanadium	>	50.942	41 Niobium	<u>Q</u>	92.906	Tantalum	Та	180.948	105 Dubnjum	Dp	262
2 Beryllium Beryllium Beryllium Beryllium 12 Beryllium 30,0122 20 21 24,305 20 24,305 20 24,305 26 38 38 39 Strontium Sr Yttium Sr Y 87,62 88,906 56 57 Berlum Lanthan Ba La 137,34 138,9 88 89 Radium Actinium Actinium Radium Actinium					4	L				Ŋ		-			.	104 Rutherfordium	Ŗ	261
1 2 1 1 1 1 1 1 1 1						5	Scandium	လွ	44.956	39 Yîtrium	>	88.90	Lanthanum	Ľ	138.91	89 Actinitum	Ac	227
1 Hydrogen Hydrogen Linory 1 1 3 Lithium Linory 1 1 8 Sodium Na 22.989 1 1 9 Potassium R	2 =		4 Beryllium Be	9.0122	Mg	24.305												226
	- -	1 Hydrogen H 1.0079	3 Lithium	6.941	Na	22.989	19 Potassium	ᅩ	39.0983	37 Rubidium	Rb	85.468	Cesium	క	132.905	87 Francium	占	223

	58	59	09	61	_	63	ı	65	99	- 67	89	69	20	7.1
	Cerinm	Praesodymium	Neodymium	Promethium	-	Europium	ගී	Terbium	Dysprosium	Holmium	Erbinm	Thulium	Ytterbium	Lutetium
Lanthanides	S P	ፊ	Š	Pa	Sm	Ш	р О	P	٥	운	山	٤	Ϋ́	Ξ
	140.12	140.907	144.24	144.913	_	151.96		158.925	162.50	164.93	167.26	168.934	173.04	174.97
	06	91	92	93		95		- 62	86	66	100	101	102	103
	Thorium	Proactinium	Uranium	Neptunium	-	Americium		Berkelium	Californium	Einsteinium	Ferminm	Mendelevium	Nobelium	Lawrencium
Actinides	£	Ра	>	2 Z		Am		鮝	ざ	ß	표	Md	ŝ	۲
	232.038	231.036	238.03	237.048		243		247	242.058	254	257.095	258.10	259.101	260.105