# Algebra 1B

Fergus Munro

March 31, 2026

## Contents

**1 Vector Spaces and Linear Maps** **2**
1.1 Subspaces . . . 2
1.2 Linear Maps . . . 2
1.3 Subspaces from Linear Maps . . . 3

**2 Bases, Coordinates and Matrices** **4**
2.1 On Lists . . . 4
2.2 Spanning and Independent lists . . . 4
2.3 Spanning and Independence in $\mathbb{F}^n$ . . . 5
2.4 Bases and Isomorphisms . . . 6
2.5 Bases and Coordinates . . . 7
2.6 Bases, Linear Maps and Matrices . . . 7
2.7 Change of Basis . . . 8
2.8 Computations in $\mathbb{F}^n$ . . . 10

**3 Dimension and Rank** **10**
3.1 Dimension . . . 10
3.2 Rank . . . 13
3.3 The Rank-Nullity Theorem . . . 15
3.4 Applications to Sums and Products . . . 17

**4 Determinants and Inverses** **18**
4.1 Determinants: the Sum Formula . . . 18
4.2 Properties of Determinants . . . 18
4.3 Characterisation of det . . . 19
4.4 Minors, Adjugates and Inverses . . . 20

**5 Eigenvalues and Diagonalisation** **21**
5.1 Fundamental Theorem of Algebra . . . 21
5.2 Eigenvalues and Characteristic Polynomials of Square Matrices . 21
5.3 Diagonalisation . . . 23
5.4 Applications of Diagonalisation . . . 25
5.5 Multiplicity of Eigenvalues and Diagonalisation . . . 25

**6 Inner Product Spaces** **27**
6.1 Inner Products . . . 27
6.2 Cauchy-Schwarz Inequality . . . 28
6.3 Orthogonality . . . 28

# 1 Vector Spaces and Linear Maps

**Definition.** A **vector space** $V$ over a field $\mathbb{F}$ is a set $V$ with two operations, $+$ and $\times$.

All algebraic manipulation in a vector space ultimately comes from the Vector Space Axioms and Field Axioms.

## 1.1 Subspaces

**Definition.** Let $V$ be a vector space over $\mathbb{F}$. A subset $U \subseteq V$ is a (linear) **subspace** if

- $0 \in U$
- $U$ is closed under linear combinations, meaning

$$\forall u, w \in U,\ \lambda, \mu \in \mathbb{F} \quad \lambda u + \mu w \in U.$$

In this case, we write $U \leq V$.

**Remark.** Because addition and multiplication is defined, $U$ is a vector space in its own right.

Any $V$ has two special subspaces, $V$ itself and the trivial subspace $\{0\}$. A subspace $U$ with $U \neq V$ is said to be 'proper'.

## 1.2 Linear Maps

**Definition.** Let $V, W$ be vector spaces over $\mathbb{F}$. A map $\phi : V \to W$ is a **linear map** if

$$\phi(\lambda v + \mu w) = \lambda \phi(v) + \mu \phi(w) \quad \lambda, \mu \in \mathbb{F},\ v, w \in V.$$

If $V = W$, then $\phi$ is a linear operator on $V$.

**Lemma.**

- The identity map $\mathrm{id}_V : V \to V$, defined $\mathrm{id}_V(v) = v$, is a linear operator over $V$.
- If $\phi : V \to W$ is a linear map then

$$\phi(0) = 0, \quad \phi(-v) = -\phi(v).$$

- If $\phi : V \to W$, $\psi : W \to U$ are linear maps then $\psi \circ \phi : V \to U$ is also a linear map
- If $\phi : V \to W$ is a linear bijecion then $\phi^{-1} : W \to V$ is also linear.

**Proof.**

- $\mathrm{id}_V(\lambda v + \mu w) = \lambda v + \mu w = \lambda \mathrm{id}_V(v) + \mu \mathrm{id}_V(w)$, hence $\mathrm{id}_V$ is a linear operator.
- $\phi(0) = \phi(v + (-v)) = \phi(v) - \phi(v) = 0$
- $\psi \circ \phi(\lambda v + \mu w) = \psi(\lambda \phi(v) + \mu \phi(w)) = \lambda \psi \circ \phi(v) + \mu \psi \circ \phi(w)$ hence $\psi \circ \phi$ is linear
- Let $v, w \in W$. Hence $\phi(\phi^{-1}(w_i)) = w_i$ for arbitrary $w_i$.

$$\begin{aligned} \lambda v + \mu w &= \lambda \phi(\phi^{-1}(v)) + \mu \phi(\phi^{-1}(w)) \\ &= \phi\big(\lambda \phi^{-1}(v) + \mu \phi^{-1}(w)\big). \end{aligned}$$

  Then, by applying $\phi^{-1}$ to both sides

$$\begin{aligned} \phi^{-1}(\lambda v + \mu w) &= (\phi^{-1} \circ \phi)(\lambda \phi^{-1}(v) + \mu \phi^{-1}(w)) \\ &= \lambda \phi^{-1}(v) + \mu \phi^{-1}(w). \end{aligned}$$

□

**Definition.** A linear bijection $\phi : V \to W$ is called a (linear) **isomorphism**.
We say $V$ is **isomorphic to** $W$ if an isomorphism $\phi : V \to W$ exists, written $V \cong W$.

## 1.3 Subspaces from Linear Maps

**Definition.** Let $\phi : V \to W$ be linear.
The **kernel** of $\phi$ is $\ker \phi \subseteq V$ given

$$\ker \phi = \{v \in V \mid \phi(v) = 0\}.$$

The **image** of $\phi$ is $\operatorname{Im} \phi \subseteq W$ given by

$$\operatorname{Im} \phi = \{\phi(v) \mid v \in V\}$$

**Proposition.** Let $\phi : V \to W$ be linear. Then

- $\ker \phi \leq V$
- $\operatorname{Im} \phi \leq W$.

**Proof.**

- Let $v, w \in \ker \phi$. Then

$$\phi(\lambda v + \mu w) = \lambda \phi(v) + \mu \phi(w) = 0$$

hence $\lambda v + \mu w \in \ker \phi$, so $\ker \phi \leq V$.

- Let $\phi(v), \phi(w) \in \operatorname{Im} \phi$. Then

$$\lambda\phi(v) + \mu\phi(w) = \phi(\lambda v + \mu w) \in \operatorname{Im} V$$

Hence, $\operatorname{Im} V \leq V$.

□

**Proposition.** A linear map $\phi : V \to W$ is injective if and only if $\ker \phi = \{0\}$

**Proof.**
($\Rightarrow$) For any linear map, $\phi(0) = 0$ and 0 is always in the kernel. Hence if $\phi(v) = 0 = \phi(0)$, by injectivity $v = 0$, so $\ker \phi = \{0\}$.
($\Leftarrow$) Suppose $\phi(v) = \phi(w)$, therefore $\phi(v-w) = 0$ by linearity and since $\ker \phi = \{0\}$, $v - w = 0$, meaning $\phi$ is injective. □

**Corollary.** A linear map $\phi : V \to W$ is an isomorphism if and only if $\ker \phi = \{0\}$ and $\operatorname{Im} \phi = W$.

# 2 Bases, Coordinates and Matrices

## 2.1 On Lists

**Definition.** A **list** of vectors in a vector space $V$ is a finite sequence $v_1, ..., v_n$ of elements of $V$. A **sequence** is a map $\{1, ..., n\} \to V$ defined $i \to v_i$

There are some differences between lists and (sub)sets:

- lists are ordered
- lists allow the same element to appear more than once

## 2.2 Spanning and Independent lists

**Definition.** Let $\alpha : v_1, ..., v_n \in V$ be a list of vectors in a vector space $V$ over $\mathbb{F}$.
The **span** of $\alpha$ is given by

$$span\ \alpha = span\{v_1, ..., v_n\} := \left\{ \sum_{i=1}^{n} \lambda_i v_i | \lambda_1, ..., \lambda_n \in \mathbb{F} \right\} \leq V.$$

More generally, for $S \subseteq V$, the span of $S$ is

$$spanS := \left\{ \sum_{i=1}^{n} \lambda_i s_i | s_1, ..., s_n \in S, \lambda_1, ..., \lambda_n \in \mathbb{F}, n \in \mathbb{N} \right\} \leq V.$$

*span* $S$ is the smallest subspace of $V$ containing $S$ :

$$span\ S = \bigcap_{u \leq V \ | \ S \subseteq u} u.$$

**Definition.** Let $\alpha : v_1, ..., v_n \in V$ be a list of vectors in a vector space $V$ over $\mathbb{F}$.

- $\alpha$ is **(linearly) independent** if, whenever $\lambda_1, ..., \lambda_n \in \mathbb{F}$ have

$$\sum_{i=1}^{n} \lambda_i v_i = 0 \text{ then } \lambda_1 = ... = \lambda_n = 0.$$

  and **(linearly) dependent** otherwise.

- $\alpha$ is **spanning** or **spans** if

$$span\ \alpha = V.$$

- $\alpha$ is a **basis** of $V$ if it is both independent and spans.

**Remark.** Independence of a list $\alpha : v_1, ..., v_n \in V$ says that no element $v_i \in \alpha$ can be expressed as another element multiplied by a scalar $\lambda$, or furthermore

$$\sum_{i=1}^{n} \lambda_i v_i \neq \sum_{i=1}^{n} \mu_i v_i.$$

unless $\lambda_i = \mu_i$ for all $i$, meaning there is a one-to-one mapping from the list $\lambda_1, ..., \lambda_n$ to the span of $\alpha$.

**Lemma.**

- A single vector $v$ is independent if $v \neq 0$
- A two element list $v_1, v_2$ is dependent $\Leftrightarrow v_1 = \lambda v_2$ for some $\lambda \in \mathbb{F}$

## 2.3 Spanning and Independence in $\mathbb{F}^n$

**Proposition.** Let $\alpha : v_1, ..., v_n \in \mathbb{F}^M$ and define $A \in M_{M \times N}(\mathbb{F})$ by $col_i(A) = v_i$. Let $\hat{A}$ be an REF matrix obtained from $A$ by gaussian elimination. Then

- The list $\alpha$ is independent if and only if every column of $\hat{A}$ has a pivot.
- $\alpha$ is spanning if $\hat{A}$ has no zero rows.

**Proof.**

- $\alpha$ is independent $\Leftrightarrow \sum_{i=1}^{n} \lambda_i v_i \neq 0$ for nonzero vector $\lambda \Leftrightarrow N_A = \{0\}$ $\Leftrightarrow \hat{A}$ has a pivot in every collumn
- $\alpha$ spanning $\Leftrightarrow \forall v, v = \sum_{i=1}^{n} \lambda_i v_i$ for some vector $\lambda \Leftrightarrow Ax = b$ always has a solution $\Leftrightarrow A$ has no zero rows

$\square$

## 2.4 Bases and Isomorphisms

**Proposition.** Let $\alpha : v_1, ..., v_n$ be a list in $V$ and define $\phi_\alpha : \mathbb{F}^n \to V$

$$\phi_\alpha(\mathbf{x}) \to \sum_{i=1}^{n} v_i x_i.$$

Then

- $\alpha$ is independent if and only if $\phi_\alpha$ is injective
- $\alpha$ spans if and only if $\phi_\alpha$ is surjective.

Hence $\alpha$ is a basis if and only if $\phi_\alpha$ is an isomorphism.

**Proof.**

- $\alpha$ independent $\Leftrightarrow \phi_\alpha(v) = 0 \Rightarrow v = 0 \Leftrightarrow \phi_\alpha(v - w) = 0 \Rightarrow v - w = 0$ $\Leftrightarrow \phi_\alpha(v) = \phi_\alpha(w) \Rightarrow v = w \Leftrightarrow \phi_\alpha$ injective.
- $\alpha$ spans $\Leftrightarrow$ every $v$ can be written in terms of $\alpha \Leftrightarrow \forall w \exists v \phi_\alpha(v) = w$ $\Leftrightarrow \phi_\alpha$ surjective.

$\square$

**Corollary.** A list $\alpha : v_1, ..., v_n \in V$ is a basis if and only if every $v \in V$ can be written as a linear combination

$$v = \sum_{i=1}^{n} x_i v_i$$

for unique $x_1, ..., x_n \in \mathbb{F}$.

**Proposition.** Let $\phi : \mathbb{F}^n \to V$ be an isomorphism. Then $\phi = \phi_\alpha$ where $\alpha : \phi(e_1), ..., \phi(e_n) \in V$ is a basis.

**Proof.**

$$\phi(v) = \phi\left(\sum_{i=1}^{n} \lambda_i e_i\right) = \sum_{i=1}^{n} \lambda_i \phi(e_i) = \phi_\alpha(v)$$

$\square$

## 2.5 Bases and Coordinates

A basis allows us to represent abstract vectors by lists of scalars (coordinates).

**Definition.** Let $\alpha : v_1, ..., v_n \in V$ be a basis of $V$. For $v \in V$, we call the vector $(x_1, ..., x_n) \in \mathbb{F}^n$ defined by

$$v = \sum_{i=1}^{n} x_i v_i.$$

This is the **coordinate vector** of $v$ with respect to $\alpha$.

**Remark.**

- $v = \phi_\alpha(x_1, ..., x_n)$ so

$$(x_1, ..., x_n) = \phi_\alpha^{-1}(v).$$

- Reordering $\alpha$ gives a different basis with different coordinate vectors.

## 2.6 Bases, Linear Maps and Matrices

**Definition.** Let $V, W$ be vector spaces over $\mathbb{F}$, $\alpha : v_1, ..., v_n \in V$ be a basis of $V$, $\beta : w_1, ..., v_m \in W$ be a basis of W, and $\phi : V \to W$ is a linear map.

The **matrix** of $\phi$ with respect to $\alpha, \beta$ is $A \in M_{m,n}(\mathbb{F})$ given by

$$\phi(v_j) = \sum_{i=1}^{m} A_{ij} w_i.$$

This means the $j^{th}$ column of $A$ is $\phi(v_j)$ written in terms of $\beta$.

When $V = W$, we can take $\alpha = \beta$

**Definition.** Let $V$ be a vector space over $\mathbb{F}$ with basis $\alpha = v_1, ..., v_n$ and $\phi : V \to V$ is a linear map on $V$. The **matrix** of $\phi$ with respect to $\alpha$ is $A \in M_n(\mathbb{F})$ given by

$$\phi(v_j) = \sum_{i=1}^{n} A_{ij} v_i.$$

**Proposition.** Let $\alpha : v_1, ..., v_n$ be a basis of $V$, $\beta : w_1, ..., v_m$ be a basis of $W$, and $\phi : V \to W$ is a linear map with matrix $A$ with respect to $\alpha, \beta$.

- For $v \in V$ with coordinate vector $x$ with respect to $\alpha$, $\phi(v)$ has coordinates y with respect to $\beta$ where

$$\underline{y} = A\underline{x}.$$

- We have
$$\phi = \phi_\beta \circ \phi_A \circ \phi_\alpha^{-1}$$
or, equivalently
$$\phi_A = \phi_\beta^{-1} \circ \phi \circ \phi_\alpha.$$

**Proof.**

- To prove the first part:
$$\begin{aligned}\phi(v) &= \sum_{j=1}^{n} x_j \phi(v_j) \\ &= \sum_{j=1}^{n} x_i (\sum_{i=1}^{m} A_{ij} w_i) \\ &= \sum_{i=1}^{m} \sum_{j=1}^{n} A_{ij} x_j w_i \\ &= \sum_{i=1}^{m} (A\underline{x})_i w_i.\end{aligned}$$
hence $\phi(v)$ has coordinates $A\underline{x}$ with respect to $\beta$, so $y = A\underline{x}$.
- Let $w = \phi(v)$. Since $y = \phi_\beta^{-1}(w) = A\phi_\alpha^{-1}(v)$ we have
$$\phi_\beta^{-1} \circ \phi = \phi_A \circ \phi_\alpha^{-1}$$
from which we can obtain the necessary equations.

□

**Proposition.** Let $V, W, U$ be vector spaces over $\mathbb{F}$ with bases $\alpha, \beta, \gamma$ respectively. Let $\phi : V \to W$ and $\psi : W \to U$ be linear maps with matrices $A$ with respect to $\alpha, \beta$ and $B$ with respect to $\beta, \gamma$.

Then $\psi \circ \phi$ has matrix $BA$ with respect to $\alpha, \gamma$.

**Proof.**

$$\begin{aligned}\psi \circ \phi &= (\phi_\gamma \circ \phi_B \circ \phi_\beta^{-1}) \circ (\phi_\beta \circ \phi_A \circ \phi_\alpha^{-1}) \\ &= \phi_\gamma \circ \phi_B \circ (\phi_\beta^{-1} \circ \phi_\beta) \circ \phi_A \circ \phi_\alpha^{-1} \\ &= \phi_\gamma \circ \phi_B \circ \phi_A \circ \phi_\alpha^{-1} \\ &= \phi_\gamma \circ \phi_{BA} \circ \phi_\alpha^{-1}.\end{aligned}$$

hence $\psi \circ \phi$ has matrix $BA$ with respect to $\alpha, \gamma$.

□

## 2.7 Change of Basis

What happens if we want to compose two linear maps that use different bases?

**Definition.** Let $\alpha : v_1, ..., v_n$ and $\alpha' : v'_1, ..., v'_n$ be two bases of $V$. Then the change of basis matrix with respect to $\alpha, \alpha'$ is $P \in M_n(\mathbb{F})$ such that

$$v'_j = \sum_{i=1}^{n} P_{ij} v_i.$$

This means the $j^{th}$ column of $P$ is $v'_j$ written in terms of $\alpha$.

**Lemma.** $\phi_P = \phi_\alpha^{-1} \circ \phi_{\alpha'}$ and so is an isomorphism, hence $P$ is invertible.

**Proof.** $\phi_P$ is the identity map with respect to $\alpha', \alpha$, hence

$$\begin{aligned} \phi_P &= \phi_\alpha^{-1} \circ \mathrm{id}_V \circ \phi_{\alpha'} \\ &= \phi_\alpha^{-1} \circ \phi_{\alpha'}. \end{aligned}$$

and as it is the composition of isomorphism, it is itself and isomorphism and hence invertible. □

**Corollary.** The change of basis matrix from $\alpha'$ to $\alpha$ is $P^{-1}$.

**Proposition.** Let $P$ be a change of basis matrix from $\alpha$ to $\alpha'$. Let $v \in V$ have coordinate vector $x$ with respect to $\alpha$ and $x'$ with respect to $\alpha'$. Then

$$\underline{x} = P\underline{x}'.$$

**Proof.** $P$ is the matrix of $\mathrm{id}_V$ with respect to $\alpha'$ and $\alpha$, so $\mathrm{id}_V = \phi_\alpha \circ \phi_P \circ \phi_{\alpha'}^{-1}$ or $\phi_P = \phi_\alpha^{-1} \circ \phi_{\alpha'}$. Since

$$\phi_\alpha(\underline{x}) = \phi_{\alpha'}(\underline{x}')$$

this means

$$\begin{aligned} \underline{x} &= \phi_\alpha^{-1} \circ \phi_{\alpha'}(\underline{x}') \\ &= P\underline{x}'. \end{aligned}$$

□

**Proposition.** Let $\alpha, \alpha'$ be bases of $V$ with change of basis matrix $P$ from $\alpha$ to $\alpha'$. Let $\beta, \beta'$ be bases of $W$ with change of basis matrix $Q$ from $\beta$ to $\beta'$.

Let $\phi : V \to W$ is a linear map with matrix $A$ with respect to $\alpha, \beta$ and $A'$ with respect to $\alpha', \beta'$. Then

$$A' = Q^{-1}AP.$$

**Proof.** We can say $\phi = \mathrm{id}_W \circ \phi \circ \mathrm{id}_V$, with matrices $Q^{-1}, A$ and $P$ respectively. Therefore $A' = Q^{-1}AP$. □

**Corollary.** If $\alpha, \alpha'$ are bases of $V$ with change of basis matrix $P$ from $\alpha$ to $\alpha'$, then, if $\phi : V \to W$ is a linear operator on $V$ with matrix $A$ with respect to $\alpha$ and $A'$ with respect to $\alpha'$, then

$$A' = P^{-1}AP.$$

These results prompt the following definition.

**Definition.**

- Matrices $A, A' \in M_{M,N}(\mathbb{F})$ are **equivalent** if there exists invertible matrices $Q \in M_M(\mathbb{F}), P \in M_N(\mathbb{F})$ such that

$$A' = Q^{-1}AP.$$

- $A, A' \in M_N(\mathbb{F})$ are **similar** if there exists invertible $P \in M_N(\mathbb{F})$ such that

$$A' = P^{-1}AP.$$

Both equivalence and similarity are equivalence relations.

### 2.8 Computations in $\mathbb{F}^n$

**Example.** When $\alpha$ is the standard mases of $\mathbb{F}^n$, and $\alpha' : v_1', ..., v_n'$ another basis, then $P$ is easy to compute:

$$col_j(P) = v_j'.$$

The reason we care about all this: there may be a cunning choice of basis which gives a simpler matrix for a given linear map.

## 3 Dimension and Rank

### 3.1 Dimension

**Lemma.** (Fundamental Lemma)
Let $A \in M_{M,N}(\mathbb{F})$. If $n > m$ then there is a nonzero $x \in \mathbb{F}^n$ such that $A\underline{x} = 0$.

**Theorem.**

- If $v_1, ..., v_n \in V$ are an independent list of vector, and $w_1, ..., w_m$ are a spanning list, then $n \leq m$.
- If $v_1, ..., v_n$ and $w_1, ..., w_m$ are bases, then $n = m$.

**Proof.**

- We can write each $v_j = \sum_{i=1}^m A_{ij}w_i$ for some matrix $A \in M_{m,n}(\mathbb{F})$.

Hence suppose

$$\sum_{j=1}^{n}\sum_{i=1}^{m} A_{ij}w_i = 0 = A\mathbf{w}$$

. If $n > 0$, there will always be a nonzero solution, contradicting the independence of $v_1, ..., v_n$. Therefore $n \leq m$.

- Since both lists are a basis so both surjective and injective, $n \leq m$ and $m \leq n$. Therefore $n = m$.

□

**Definition.** Let $V$ be a vector space.

- $V$ is **finite dimensional** if it has a finite length basis and **infinite dimensional** otherwise
- If $V$ is finite dimensional, the **dimension** of $V$, denoted $\dim V$ is the length of any basis of $V$.

**Example.**

- $\dim \mathbb{F}^n = n$
- $\dim \mathbb{F}^0 = \dim\{0\} = 0$
- $\dim \mathbb{F}[x] = \infty$

Dimension is preserved by isomorphism.

**Proposition.** Let $V$ be a vector space over $\mathbb{F}$. Then $\dim V = n$ if and only if $\mathbb{F}^n \cong V$.

In this case, if $W \cong V$ then W is finite dimensional also with $\dim W = n$.

**Proof.** Any isomorphism $\phi : \mathbb{F}^n \to V$ creates a basis $\alpha$ of $V$ by applying $\phi$ to $e_i, ..., e_n$, and is uniquely determined by this basis. The second statement can be proved by composing the two isomorphisms, meaning $W \cong \mathbb{F}^n$ and so $\dim W = n$. □

**Proposition.**

- If $v_1, ..., v_n$ span $V$ and $v_n \in span\{v_1, ..., v_{n-1}\}$ then $v_1, ..., v_{n-1}$ span $V$.
- If $v_1, ..., v_{n-1}$ is an independent list and $v_n \notin span\{v_1, ..., v_{n-1}\}$ then $v_1, ..., v_n$ is also independent.

**Proof.**

$$
\begin{aligned}
V &= span\{v_1, ..., v_n\} \\
&= \{\sum_{i=1}^{n} \lambda_i v_i | \lambda_1, ..., \lambda_n \in \mathbb{F}, v_1, ..., v_n \in V\} \\
&= \{\sum_{i=1}^{n-1} \lambda_i v_i + v_n | \lambda_1, ..., \lambda_n \in \mathbb{F}, v_1, ..., v_n \in V\} \\
&= \{\sum_{i=1}^{n-1} \lambda_i v_i + \sum_{i=1}^{n-1} \mu_i v_i | \lambda_1, ..., \lambda_n \in \mathbb{F}, \mu_1, ..., \mu_n \in \mathbb{F}, v_1, ..., v_n \in V\} \\
&= \{\sum_{i=1}^{n-1} (\lambda_i + \mu_i) v_i | \lambda_1, ..., \lambda_n \in \mathbb{F}, \mu_1, ..., \mu_n \in \mathbb{F}, v_1, ..., v_n \in V\} \\
&= span\{v_1, ..., v_{n-1}\}.
\end{aligned}
$$

For the second statement, suppose $\sum_{i=1}^{n-1} \lambda_i v_i + v_n = 0$. Then if not independent, $v_n$ can be written as some linear combination of $v_1, ..., v_{n-1}$, so it would be in the span of those vectors, which is a contradiction. □

**Definition.** Let $\alpha$ be a list of vectors in $V$.

- a **sublist** of $\alpha$ is a list obtained by deleting some elements of $\alpha$
- a list $\beta$ **extends** $\alpha$ if $\beta$ is obtained by appending vectors to $\alpha$.

**Theorem.** Let $\beta$ be a finite spanning list of vectors in a vector space $V$ extending a (possibly empty) independent list $\alpha$. Then there is a basis $\gamma$ of $V$ which

- extends $\alpha$
- is a sublist of $\beta$.

**Proof.** Let $\gamma$ be the longest independent sublist of $\beta$. If all elements of $\beta$ can be written in terms of $\gamma$, it is a basis. If one cannot, then appending this vector to $\gamma$ gives an independent sublist of $\beta$, contradicting $\gamma$'s maximal length. □

**Corollary.**

- Any finite spanning list contains a basis
- If $V$ is finite dimensional, any independent list can be extended to a basis.

**Corollary.** Let $\dim V = n$.

- Any spanning list of length $n$ is a basis.
- Any independent list of length $n$ is a basis.

**Proof.** Both lists contain a length $n$ sublist that is a basis. Since they are themselves length $n$, that sublist is just the entire list. $\square$

**Theorem.** Let $V$ be finite dimensional and $W \leq V$. Then $W$ is finite dimensional also and $\dim W \leq \dim V$.

Futheremore, $\dim W = \dim V \Leftrightarrow W = V$.

**Proof.** Any independent list in $W$ has length $n \leq \dim V$. Consider an independent list $\alpha$ of maximum length in $W$. If it does not span $W$, a longer independent list exists so there is a contradiction. Therefore, this list is a basis of $W$ with $\dim W \leq \dim V$. If $\dim W = \dim V$, $\alpha$ is a basis of $V$ with $W = span\ \alpha = V$. $\square$

**Remark.** If two vector spaces have the same dimension, they are only equal if one is a subspace of the other. Equal dimension alone is not enough to show they are equal.

**Proposition.** Let $\alpha : v_1, ..., v_n \in \mathbb{F}^m$ and define $A \in M_{m,n}(\mathbb{F})$ by $col_j(A) = \bar{v}_j$.

Let $\hat{A}$ be a REF matrix obtained from $A$ by gaussian elimination with $r$ pivots in cols $j_1 < j_2 < ... < j_r$. Then $v_{j_1}, ..., v_{j_r}$ is a basis for $span\ \alpha \leq \mathbb{F}^M$.

**Proof.** Consider the matrix $B \in M_{m,r}(\mathbb{F})$ with columns from the list $\beta : v_{j_1}, ..., v_{j_r}$. Then since EROs act on each column separately, $\hat{B}$ has a pivot in every column, thus $\beta$ is independent.

To show $span\ \alpha = span\ \beta$, this is the same as proving that for every $v_j$, there exists $x^j$ such that

$$B\mathbf{x^j} = \mathbf{v_j}.$$

Suppose this is unsolvable. This is trivially solvable for $v_j$ in $\beta$, so suppose $v_j$ is not. Then the REF of the augmented matrix $(B|v_j)$ has a row where all entries in $B$-columns are zero and the $v_j$ entry is nonzero. This means this column contains a pivot, so $v_j$ would be in $\beta$, a contradiction. $\square$

## 3.2 Rank

**Definition.** The **rank** of a linear map $\phi : V \to W$ is

$$rank\ \phi := \dim \operatorname{Im} \phi.$$

**Lemma.** Given $\phi : V \to W$ and isomorphisms $f : V' \to V$ and $g : W' \to$

$W$, set $\phi' = g^{-1} \circ \phi \circ f : V' \to W'$. Then

$$rank\ \phi = rank\ \phi'.$$

**Proof.** Since $f$ is surjective, $\mathrm{Im}(\phi \circ f) = \mathrm{Im}\,\phi$. Then restricting $g$ to domain $\mathrm{Im}\,\phi$ and codomain $\mathrm{Im}(g \circ \phi)$ gives an isomorphism, hence the two images are isomorphic and so $\phi$ and $\phi'$ have the same rank. $\square$

**Definition.** Let $A \in M_{M,N}(\mathbb{F})$.

- The **column space** of $A$ is the span of the columns of $A$.

$$colspace\ A := \left\{ \sum_{j=1}^{n} x_j col_j(A) | x_1, ..., x_n \in \mathbb{F} \right\}.$$

  The **column rank** of $A$ is $colrank\ A := dim\ colspace\ A$.

- The **row space** of $A$ is the span of the rows of $A$

$$rowspace\ A := \left\{ \sum_{j=1}^{m} x_j row_j(A) | x_1, ..., x_n \in \mathbb{F} \right\}.$$

  The **row rank** of $A$ is $row\ rank\ A := dim\ rowspace\ A$.

- $rank\ A$ is the number of pivots in the REF of $A$.

**Theorem.** Let $A \in M_{m,n}(\mathbb{F})$. Then $rank\ A$ is equal to:

- $rank\ \phi_A$
- $colrank\ A$
- $rowrank\ A$
- The smallest number $r$ such that

$$A = BC$$

  with $B \in M_{M,r}, C \in M_{r,N}$.

**Proof.** Firstly, $colspace\ A = \mathrm{Im}\,A$ so $colrank\ A = rank\ \phi_A$. If each column of $A$ represents a list of vectors $\alpha$, the columns with pivots in $\hat{A}$ form a basis of $span\ \alpha$, so $rank\ A = colrank\ A$. It is possible to factorize $A = BC$ with $B \in M_{m,r}$ and $C \in M_{r,n}$ with $r = colrank\ A$, by setting the columns of $B$ to be a basis of the column space. Now suppose $r$ is not minimal, so there exists $s < r$. Then

$$col_k(A) = \sum_{j=1}^{s} c_{jk} col_j(B)$$

thus there exists a basis of length $s$ for *colspace* $A$, which is a contradiction. Therefore $r$ is the smallest value to factorize $A$. A similar argument can be used with the row rank. $\square$

**Corollary.** For $A \in M_{M,N}$, $rank\ A = rank\ A^T$.

**Proof.** $rank\ A = rowspace\ A = colspace A^T = rank\ A^T$. $\square$

**Remark.** To find the rank of a matrix $A$, and the 'smallest' matrices $B$ and $C$ such that $A = BC$, reduce $A$ into RREF. Then the number of pivots is $r = rank\ A$. The columns of $B$ are the columns of $A$ where there is a pivot in $\hat{A}$, and the columns of $C$ are the first $r$ rows of $\hat{A}$.

**Proposition.**

- Let $\phi : V \to W$ be a linear map with matrix $A$ with respect to some bases $\alpha, \beta$. Then
$$rank\ A = \operatorname{rank} \phi.$$
- Let $A, B \in M_{M,N}(\mathbb{F})$ be equivalent. Then
$$\operatorname{rank} A = \operatorname{rank} B.$$

**Proof.**

- $\phi = \phi_\beta \circ \phi_A \circ \phi_\alpha^{-1}$. Then because $\phi_\beta$ and $\phi_\alpha$ are isomorphisms
$$\operatorname{rank} \phi = \operatorname{rank} \phi_A = \operatorname{rank} A.$$
- Let $A = QBP^{-1}$ for invertible $P, Q$. Then
$$\phi_A = \phi_Q \circ \phi_B \circ \phi_P^{-1}$$
thus
$$\operatorname{rank} A = \operatorname{rank} \phi_A = \operatorname{rank} \phi_B = \operatorname{rank} B.$$

$\square$

**Remark.** The above statement has a converse: If $\operatorname{rank} A = r$ then $A$ is equivalent to $\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}$ Additionaly, matrices of the same size and rank are equivalent under an equivalence relation.

## 3.3 The Rank-Nullity Theorem

**Definition.** The **nullity** of a linear map $\phi$ is
$$null\ \phi = \dim \ker \phi.$$

**Theorem.** (Rank-Nullity Theorem)
If $V$ is finite dimensional and $\phi : V \to W$ is linear, then

$$rank\ \phi + null\ \phi = dim\ V.$$

**Proof.** Let $\dim V = n$, $\dim \ker \phi = k$. Then we must prove $\dim \operatorname{Im} \phi = n - k$. Consider a basis of $\ker \phi v_1, ..., v_k$ extended by $v_{k+1}, ..., v_n$ to form a basis of $V$, and we want to prove that $\phi(v_{k+1}), ..., \phi(v_n)$ is a basis of $\operatorname{Im} \phi$.

- (Independence). Suppose

$$\begin{aligned} \sum_{k+1=1}^{n} \lambda_i \phi(v_i) &= 0 \\ \phi\left(\sum_{i=k+1}^{n} \lambda_i v_i\right) &= 0 \\ \phi(w) &= 0. \end{aligned}$$

  so $w \in \ker \phi$. However this means that it could be written in terms of $v_1, .., v_k$, but this is a contradiction as the two sublists are independent due to being a basis of $V$.

- (Spanning). Let $v \in V$. Then

$$\begin{aligned} \sum_{i=1}^{n} \lambda_i \phi(v_i) &= v \\ \sum_{i=1}^{k} \lambda_i \phi(v_i) + \sum_{k+1=1}^{n} \lambda_i \phi(v_i) &= v \\ \sum_{k+1=1}^{n} \lambda_i \phi(v_i) &= v \\ &. \end{aligned}$$

  so $v_{k+1}, ..., v_k$ is a basis of $\operatorname{Im} \phi$. Hence

$$\operatorname{rank} \phi + \operatorname{null} \phi = \dim V.$$

□

**Remark.** The dimension of $W$ need not be finite.

**Theorem.** Let $\phi : V \to V$ be a linear operator on finite dimensional $V$. Then the following are equivalent:

- $\phi$ is an isomorphism
- $\phi$ is surjective

- $\phi$ is injective
- $\ker \phi = \{0\}$

**Proof.** $\phi$ is surjective $\Leftrightarrow \operatorname{Im}\phi = V \Leftrightarrow \operatorname{rank}\phi = \dim V \Leftrightarrow \operatorname{null}\phi = 0 \Leftrightarrow \ker\phi = \{0\} \Leftrightarrow \phi$ is injective. $\square$

**Remark.** This result fails if $V$ is infinite dimensional.

## 3.4 Applications to Sums and Products

**Definition.** Let $V, W$ be vector spaces over $\mathbb{F}$. Then the **product** of $V$ and $W$ is $V \times W$ with addition and scalar multiplication defined componentwise

$$(v, w) + (v', w') = (v + v', w + w')$$
$$\lambda(v, w) = (\lambda v, \lambda w).$$

$V \times W$ is a vector space over $\mathbb{F}$ with this addition and scalar multiplication.

**Proposition.** If $V$, $W$ are finite dimensional, so is $V \times W$ and

$$\dim(V \times W) = \dim V + \dim W.$$

**Proof.** Consider a linear projection map $\pi_V : (V \times W) \to V$, $\pi_(v, w) \mapsto v$. Then $\operatorname{rank}\pi_V = \dim V$ and $\operatorname{null}\pi_V = \dim W$. Thus

$$\dim(V \times W) = \dim V + \dim W.$$

$\square$

**Definition.** Let $U, W \leq V$. Then the **sum** of $U$ and $W$ is

$$U + W := \{u + w | u \in U, w \in W\}.$$

**Remark.** $U + W$ is itself a subspace of $V$.

**Proposition.** (Dimensional Formula)
If $U, W \leq V$ are finite dimensional. Then

$$\dim(U + W) + \dim U \cap W = \dim U + \dim W.$$

**Proof.** Consider $\phi : (U \times W) \to U + W$, $\phi(v, w) \mapsto u + w$. This clearly surjects so $\operatorname{rank}\phi = \dim(U + W)$, hence we just need to prove $\operatorname{null}\phi =$

$\dim U \cap W$.

$$\phi(u, w) = u + w = 0.$$
$$w = -u.$$

so $w \in u$ and $u \in W$. Thus $\ker \phi \cong U \cap W$ so they have the same dimension. $\square$

# 4 Determinants and Inverses

## 4.1 Determinants: the Sum Formula

**Definition.** Let $A = (a_{ij}) \in M_N(\mathbb{F})$. The determinant of $A$, denoted $\det A$ or $|A|$ is

$$\det A = \sum_{\sigma \in S_n} sign(\sigma) a_{\sigma(1)1} ... a_{\sigma(n)n}.$$

This is a signed product of one entry from each collummn, in a different row each time, all added together.

**Remark.** The are $n!$ summands so this formula is not easy to use as $n$ gets larger. Fortunately, there are other ways to compute $\det A$.

**Proposition.** If $A = \begin{pmatrix} A' & * \\ 0 & A'' \end{pmatrix}$ with $A' \in M_N(\mathbb{F}), A'' \in M_M(\mathbb{F})$ is block diagonal (or upper triangular) then $\det A = \det A' \det A''$.

**Proposition.** $\det A^T = \det A$.

## 4.2 Properties of Determinants

We view $\det A$ as a function of the columns of $A$:
write $A = (\underline{c}_1, ..., \underline{c}_n)$ $\quad \underline{c}_j = col_j(A)$ and view $\det : (\underline{c}_1, ..., \underline{c}_n) \mapsto \det A$

**Proposition.** $\det A = \det(\underline{c}_1, ..., \underline{c}_n)$ is

- **multilinear** (linear in each slot separately): if $c_j = \lambda \underline{v} + \mu \underline{w}$ where $v, w$ are column vectors, $\lambda, \mu \in \mathbb{F}$, then $\forall j, 1 \leq j \leq n$

$$\det(\underline{c}_1, ..., \underline{c}_j, ..., \underline{c}_n) = \lambda \det(\underline{c}_1, ..., \underline{v}, ..., \underline{c}_n) + \mu \det(\underline{c}_1, ..., \underline{w}, ..., \underline{c}_n)$$

- **alternating**: if $\underline{c}_i = \underline{c}_j$ for some $i \neq j$, then

$$\det(\underline{c}_1, ..., \underline{c}_n) = 0.$$

**Corollary.** Considering $A \in M_N(\mathbb{F})$ as a list of row vectors $\underline{r}_1, ..., \underline{r}_n \in M_N(\mathbb{F})$, and the function $\det A = \det(\underline{r}_1, ..., \underline{r}_n)$ is

- **multilinear** (linear in each slot separately): if $r_j = \lambda \underline{v} + \mu \underline{w}$ where $v, w$ are column vectors, $\lambda, \mu \in \mathbb{F}$, then $\forall j, 1 \leq j \leq n$

$$\det(\underline{r}_1, ..., \underline{r}_j, ..., \underline{r}_n) = \lambda \det(\underline{r}_1, ..., \underline{v}, ..., \underline{r}_n) + \mu \det(\underline{r}_1, ..., \underline{r}, ..., \underline{r}_n)$$

- **alternating**: if $\underline{r}_i = \underline{r}_j$ for some $i \neq j$, then

$$\det(\underline{r}_1, ..., \underline{r}_n) = 0.$$

**Proposition.** If $\hat{A}$ is obtained from $A$ by

- $\hat{\underline{c}}_i = \lambda \underline{c}_i$ then $\det \hat{A} = \lambda \det A$
- $\hat{\underline{c}}_i = \underline{c}_i + \lambda \underline{c}_j$ then $\det \hat{A} = \det A$
- $\underline{c}_i \leftrightarrow \underline{c}_j, (i \neq j)$ (swap columns $i$ and $j$) then $\det \hat{A} = -\det A$

**Remark.** Since $\det A^T = \det A$, this works just as well for EROs as it does for ECOs. This gives us a method to compute the determinant:

Use guassian elmination (also using ECOs) to turn $A$ into an upper or lower triangular $\hat{A}$, keeping track of the changes to the determinant.

**Corollary.** If $\hat{A}$ is obtained from $A$ by a mix of EROs and ECOs. Then

$$\det \hat{A} = 0 \Leftrightarrow \det A = 0.$$

**Corollary.** If $\hat{A}$ is obtained from $A$ by permuting the columns of $A$ (or the rows) by $\sigma \in S_n$ then

$$\det \hat{A} = \operatorname{sign} \sigma \det A.$$

## 4.3 Characterisation of det

Fix $n > 0$, and let $I := I_n \in M_n(\mathbb{F})$, the identity matrix.

**Theorem.** Let $D : M_N(\mathbb{F}) \to \mathbb{F}$ be a function that is multilinear and alternating in columns. Then

$$D(A) = D(I) \det A.$$

**Corollary.** If $D : M_N(\mathbb{F}) \to \mathbb{F}$ is multilinear and alternating and $D(I) = 1$, then

$$D = \det.$$

**Theorem.** (Product formula)
If $A, B \in M_n(\mathbb{F})$, then

$$\det(AB) = \det A \det B.$$

**Corollary.**

- If $P \in M_m(\mathbb{F})$ is invertible, then $\det(P_{-1}) = det(P)^{-1}$.
- If $A$ and $B$ are similar, then $\det A = \det B$.

**Remark.** If $\phi : V \to V$ is a linear operator and $A, B$ are the matrices of $\phi$ with respect to two bases, then $A$ and $B$ are similar, so $\det A = \det B$.
Hence the following is well defined.

**Definition.** Let $\phi : V \to V$ be a linear map, and $V$ be finite dimensional. The **determinant** of $\phi$, $\det \phi$, is

$$\det \phi := \det A$$

where $A$ is the matrix of $\phi$ with respect to some basis.

## 4.4 Minors, Adjugates and Inverses

We introduce a method for computing the determinant, and the inverse of a matrix.

**Definition.** For $A = (a_{ij}) \in M_n(\mathbb{F})$ the $(i, j)$-minor is

$$\mu_{ij}(A) := \det M_{ij}(A)$$

where $M_{ij}(A)$ is the matrix obtained by deleting the $i^{th}$ row and $j^{th}$ column of $A$.

**Proposition.** (column expansion and row expansion of determinant)

- For any fixed column $j$

$$\det(A) = \sum_{i=1}^{n} (-1)^{i+j} a_{ij} \mu_{ij}(A).$$

- For any fixed row $i$

$$\det(A) = \sum_{j=1}^{n} (-1)^{i+j} a_{ij} \mu_{ij}(A).$$

**Definition.** The **adjugate** of a matrix $A \in M_N(\mathbb{F})$ is

$$adj(A) = (b_{ij}) \in M_N(\mathbb{F})$$

where

$$b_{ij} = (-1)^{i+j}\mu_{ji}(A)$$

We say $(-1)^{i+j}\mu_{ij}$ is the $(i,j)$-cofactor and so $adj(A)$ is the transpose of the cofactor matrix.

**Theorem.**

$$adj(A)A = \det(A)I.$$

**Corollary.** If $\det A \neq 0$ and then $A$ is invertible and

$$A^{-1} = \frac{1}{\det A} ajd(A).$$

**Remark.** By the product formula, if $A$ is invertible then $\det A \neq 0$. We also have the converse:

$$A \text{ is invertible } \Leftrightarrow \det A \neq 0.$$

# 5 Eigenvalues and Diagonalisation

## 5.1 Fundamental Theorem of Algebra

**Theorem.** (Fundamental Theorem of Algebra)
Let $f(x) \in \mathbb{C}[x]$ be a non-constant polynomial of degree $d > 0$. The following equivalence statements hold:

- $f(x)$ has at least one complex root
- $f(x)$ has precisely $d$ roots counted with multiplicity
- $f(x)$ factorizes as a product of linear terms

## 5.2 Eigenvalues and Characteristic Polynomials of Square Matrices

**Definition.** Let $A \in M_n(\mathbb{F})$. A scalar $\lambda \in \mathbb{F}$ is an **eigenvalue** of $A$ if there exists a non-zero vector $v$ such that

$$A\mathbf{v} = \lambda\mathbf{v}.$$

In this case the vector $v$ is an **eigenvector** of $A$ with **eigenvalue** $\lambda$ or a $\lambda$**-eigenvector**.

**Remark.** If $v$ is an eigenvector with eigenvector $\lambda$ then every nonzero scalar multiple $av$ of $v$ is also a $\lambda$-eigenvector.

There is a method for computing eigenvectors!

**Definition.** The characteristic polynomial of a matrix $A \in M_n(\mathbb{F})$ is

$$\Delta_A(t) = \det(A - tI) \in \mathbb{F}[t].$$

Where $I \in M_n(\mathbb{F})$ is the identity matrix $I = \begin{pmatrix} 1 & \dots & 0 \\ \vdots & \ddots & 0 \\ 0 & \dots & 1 \end{pmatrix}$

**Proposition.** The eigenvalues of $A$ are the roots of $\Delta_A(t)$ in $\mathbb{F}$.

**Proof.** Let $\lambda \in \mathbb{F}$. Then the following are equivalent:

- there exists non-zero $v \in \mathbb{F}^n$ such that $Av = \lambda v$.
- there exists non-zero $v \in N_{A-\lambda I}$ i.e. $(A - \lambda I)v = 0$
- $\ker \phi_{A-\lambda I} \neq 0$
- $\phi_{A-\lambda I}$ is not bijective
- $A - \lambda I$ is not invertible
- $\det(A - \lambda I) = 0$
- $\lambda$ is a solution of $\Delta_A(t) = 0$

$\square$

**Lemma.** If $A \in M_n(\mathbb{F})$ then $\Delta_A(t)$ is a degree $n$ polynomial, with leading coefficient $(-1)^n$. Therefore $\Delta_A(t)$ has t most $n$ roots. If $\mathbb{F} = \mathbb{C}$ then $\Delta_A(t)$ has precisely $n$ roots counted with multiplicity.

**Proof.**

$$\begin{vmatrix} a_{11} - t & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} - t & & \vdots \\ \vdots & & \ddots & \vdots \\ a_{n1} & \dots & \dots & a_{nn} - t \end{vmatrix}.$$

has degree $n$ with the term $t^n$ arising from the summand $\sigma = Id$ permutation.

The summand is $(a_{11} - t)(a_{22} - t)...(a_{nn} - t)$ so the leading term is $(-t)^n$, and leading coefficient $(-1)^n$.

Hence this polynomail has maximum $n$ roots, with exactly $n$ over $\mathbb{C}$.

$\square$

**Lemma.** Similar matrices have the same characteristic polynomial, hence the same eigenvalues.

**Proof.** Suppose $A = PBP^{-1}$. Then

$$\begin{aligned}\Delta_A(t) &= \det(A - tI) \\ &= \det(PBP^{-1} - tI) \\ &= \det(P(B - tI)P^{-1}) \\ &= \det(B - tI) = \Delta_B(t)..\end{aligned}$$

□

**Definition.** Let $V$ be a vector space over $\mathbb{F}$, and $\phi : V \to V$ be a linear map.

- $\lambda \in \mathbb{F}$ is an **eigenvalue** of $\phi$ if there exists nonzero $v \in V$ such that

$$\phi(v) = \lambda v.$$

  Then $v$ is a $\lambda-$**eigenvector** of $\phi$.

- The $\lambda$-eigenspace of $\phi$ is

$$\begin{aligned}E_\phi(\lambda) &= \{v \in V \mid \phi(v) = \lambda v\} \\ &= \ker(\phi - \lambda \mathrm{id}_V).\end{aligned}$$

**Lemma.** Let $A \in M_n(\mathbb{F})$, $V$ be a vector space over $\mathbb{F}$, $\phi : V \to V$ be a linear map. If $A$ is the matrix of $\phi$ in some basis $\alpha$ then $A, \phi$ have the same eigenvalues and eigenvectors.

**Definition.** The **characteristic polynomial** of $\phi : V \to V$ is

$$\Delta_\phi(t) = \det(\phi - t\mathrm{id}_V).$$

Equivalently, if $A$ represents $\phi$ in some basis $\Delta_\phi(t) := \Delta_A(t)$.

**Remark.** The characteristic polynomial $\phi$ is well defined because similar matrices have equal determinants.

## 5.3 Diagonalisation

Let $\phi : V \to V$ be a linear map and $A \in M_n(\mathbb{F})$, the matrix of $\phi$ with respect to some basis $v_1, ..., v_n$. Then the following are equivalent.

- $v_i$ is a $\lambda_i-$eigenvector for $i = 1, ..., n$.

- $A$ is diagonal and $A = \begin{pmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{pmatrix}$

**Definition.**

- A linear operator $\phi : V \to V$ is **diagonalisable** if there is a basis consisting of eigenvectors of $\phi$. In this case we call it an **eigenbasis**.
- $A \in M_n(\mathbb{F})$ is **diagonalisable** (over $\mathbb{F}$) if there is an invertible $P \in M_n(\mathbb{F})$ such that
$$P^{-1}AP \quad \text{is diagonal.}$$
We say that $A$ is **diagonalised** by $P$.

**Remark.** $A \in M_n(\mathbb{F})$ is diagonalisable if and only if $\phi_A : \mathbb{F}^n \to \mathbb{F}^n$ is diagonalisable.

**Proposition.** If $\phi : V \to V$ has a matrix $A$ with respect to a basis $\beta$. Then the following are equivalent:

- $\phi$ is diagonalisable
- $\phi_A$ is diagonalisable
- $A$ is diagonalisable

**Proof.** $\phi_A = \phi_\beta^{-1} \circ \phi \circ \phi_\beta$. If $v \in V$ is an eigenvector $\phi$ with eigenvalue $\lambda$, then

$$\begin{aligned} \phi_A(\phi_\beta^{-1}(v)) &= \phi_\beta^{-1} \circ \phi \circ \phi_\beta(\phi_\beta^{-1}(v)) \\ &= \phi_\beta^{-1}(\phi(v)) \\ &= \phi_\beta^{-1}(\lambda v) \\ &= \lambda \phi_\beta^{-1}(v). \end{aligned}$$

So $\phi_\beta^{-1}(v)$ is a $\lambda-$eigenvector of $\phi_A$. Since $\phi_\beta^{-1}$ is an isomorphism any basis of $V$ will be mapped to a corresponding basis in $\mathbb{F}^n$, so if $\phi$ is diagonalisable so is $\phi_A$. Since $\phi = \phi_\beta \circ \phi_A \circ \phi_\beta^{-1}$, the same argument will work for in reverse.

If $A$ is diagonalisable, then there exists $P$ such that

$$\begin{aligned} P^{-1}AP = D &\Leftrightarrow AP = DP \\ &\Leftrightarrow Acol_j(P) = \lambda_j col_j(P) \quad \forall j \\ &\Leftrightarrow \text{the columns of } P \text{ are an eigenbasis of } A. \end{aligned}$$

Since the columns of $P$ form a basis, $\phi_A$ is diagonalisable if and only if $A$ diagonalisable. $\square$

**Remark.** Many matrices are not diagonalisable over $\mathbb{R}$, but are diagonalisable over $\mathbb{C}$. This happens when the characteristic polynomial has no real roots.

## 5.4 Applications of Diagonalisation

Diagonalisation can be extremely helpful for efficient calculation of powers of matrices.

**Proposition.** Suppose $A$ is diagonalisable, with $P^{-1}AP = D$ being diagonal. Then $A = PDP^{-1}$, so for $k > 0$

$$\begin{aligned} A^k &= (PDP^{-1})(PDP^{-1})...(PDP^{-1}) \\ &= PD^kP^{-1}. \end{aligned}$$

with $D^k = \begin{pmatrix} \lambda_1^k & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n^k \end{pmatrix}$. This allows us to calculate $A^k$ explicitly with only 2 matrix mutliplications.

## 5.5 Multiplicity of Eigenvalues and Diagonalisation

We want a necessary and sufficient condition for a matrix or operator to be diagonalisable.

If $\dim V = n$ and $\phi : V \to V$ is linear, then an eigenbasis needs $n$ linearly Independent eigenvectors.

**Theorem.** If $v_1, ..., v_n$ are eigenvectors $\phi$ with distinct eigenvalues $\lambda_1, ..., \lambda_n$ then $v_1, ..., v_n$ are independent.

**Proof.** By contradiction, suppose WLOG that $a_1v_1 + ... + a_kv_k = 0$ is the smallest dependent list of eigenvectors. Then

$$\begin{aligned} 0 = \lambda_1 \sum_{i=1}^{k} a_iv_i &= \phi(\sum_{i=1}^{n} a_iv_i) = \sum_{i=1}^{n} a_i\phi(v_i) \\ \sum_{i=1}^{k} a_i\lambda_kv_i &= \sum_{i=1}^{n} a_i\lambda_iv_i. \end{aligned}$$

Then, by subtracting the two sums, the $v_1$ cancels giving a smaller dependent list, a contradiction. $\square$

**Remark.** This can be generalized: if we conbine independent list from each eigenspace we get an independent list.

**Corollary.** Let $n = \dim V$ and $\phi : V \to V$ be a linear operator. If

$$\Delta_\phi(t) = (t - \lambda_1)...(t - \lambda_n)$$

with $\lambda_i$ distinct for all $i$, then $\phi$ is diagonalisable.

**Proof.** Because there are $n$ distinct eigenvalues, there are $n$ distinct eigenvectors and since they are independent, they form an eigenbasis. □

**Definition.** Let $\lambda$ be an eigenvector of $\phi : V \to V$. Then

- the **algebraic multiplicity** of $\lambda$ denoted $am(\lambda)$ is the largest $k$ such that $(\lambda - t)^k$ divides $\Delta_\phi(t)$.
- the **geometric multiplicity** of $\lambda$ is $gm(\lambda) := \dim E_\phi(\lambda)$.

**Remark.**

- $\dim E_\phi(\lambda)$ is the maximal number of independent $\lambda-$eigenvectors.
- Suppose $\lambda$ is a root of $\Delta_\phi(t)$. Then $(t - \lambda)$ divides $\Delta_\phi(t)$ so

  $$am(\lambda) \geq 1.$$

  If $E_\phi(\lambda) \neq 0$ then

  $$gm(\lambda) \geq 1.$$

**Proposition.** For any $\lambda \in \mathbb{F}$ and any linear map $\phi$:

$$am(\lambda) \geq gm(\lambda).$$

**Proof.** Let $v_1, ..., v_k$ be a basis of $E_\phi(\lambda)$. Extend this list to be a basis of $V$. Then the matrix of $\phi$ from the standard basis to this basis is diagonal in the top corner, and so if $A$ is the matrix in the bottom corner, then

$$\Delta_\phi(t) = (\lambda - t)^t \det(A - It).$$

Hence $(\lambda - t)$ divides $\Delta_\phi(t)$ at least $k$ times, so $am(\lambda) \geq gm(\lambda)$. □

**Theorem.** Let $\phi : V \to V$ be a linear operator. Then $\phi$ is diagonalisable if and only if $\Delta_\phi(t)$ is a product of linear factors and $am(\lambda) = gm(\lambda)$ for all $\lambda$.

**Proof.**
($\Rightarrow$) $\phi$ has a diagonal matrix wrt to its eigenbasis, and so in this diagonal matrix, $\Delta_\phi(t) = (\lambda_1 - t)...(\lambda_n - t)$. Then if $\lambda$ is not an eigenvector, $am(\lambda) = 0 = gm(\lambda)$. If $\lambda$ is an eigenvector, $gm(\lambda)$ is the number of $\lambda-$eigenvectors, and so the number of terms equal to $(\lambda - t)$ in $\Delta_\phi(t)$,

which is $am(\lambda)$.

($\Leftarrow$) Suppose $\Delta_\phi(t) = (\lambda_1 - t)^{k_1}...(\lambda_m - t)^{k_m}$. Then since there are $n$ total terms in $\Delta_\phi(t)$

$$\sum_{i=1}^{m} k_i = n.$$

Each eigenvalue $\lambda_i$ has $k_i$ independent $\lambda_i-$eigenvectors. If these $n$ eigenvectors are independent, then there is a basis of eigenvectors of $\phi$ so $\phi$ is diagonalisable.

Suppose these vectors are dependent. Then

$$\begin{aligned} 0 &= \sum_{i=1}^{m} \sum_{j=1}^{k_i} a_i v_{i_j} \\ &= \sum_{i=1}^{m} b_i v_i. \end{aligned}$$

for some $\lambda_i-$eigenvector $v_i$. However, since these eigenvectors have distinct eigenvalues, they are linearly independent, a contradiction. $\square$

# 6 Inner Product Spaces

## 6.1 Inner Products

Recall that the dot (or scalar) product on $\mathbb{R}^n$ is the map

$$\begin{aligned} \mathbb{R}^n \times \mathbb{R}^n &\to \mathbb{R} \\ x \cdot y &= \sum_{i=1}^{n} x_i y_i.. \end{aligned}$$

Then we can define the length of $x$ to be $|x| = x \cdot x$, and the angle between two vectors with $x \cdot y = |x||y| \cos\theta$.

We also have the dot product on $\mathbb{C}^n$, defined as

$$x \cdot y = \sum_{i=1}^{n} \bar{x}_i y_i.$$

Note that $x\dot{x} = \sum_{i=1}^{n} \bar{x}_i x_i = \sum_{i=1}^{n} |x_i|^2$.

**Definition.** Let $V$ be a vector space over $\mathbb{F}$ which we assume to be either $\mathbb{R}$ or $\mathbb{C}$. An **inner product** on $V$ is a map $V \times V \to \mathbb{F} : (v, w) \mapsto \langle v, w \rangle$ which is

- **congugate symmetric:** $\langle w, v \rangle = \overline{\langle v, w \rangle}$
- **linear in the second argument:**

$$\langle v, \lambda u + \mu w \rangle = \lambda \langle v, u \rangle + \mu \langle v, w \rangle$$

- **positive definite:** For all $v \in V$, $\langle v, v \rangle \geq 0$ with equality if and only if $v = 0$

A vector space with a fixed inner product is called an **inner product space**.

An inner product space comes with an inner product already defined, denoted $\langle \cdot, \cdot \rangle$.

**Remark.** $\langle v, v \rangle = \overline{\langle v, v \rangle}$ so the inner product of a vector on itself is always real.

- Suppose $\mathbb{F} = \mathbb{R}$. Then congugate symmetry is just symmetry, and so $\langle v, w \rangle = \langle w, v \rangle$. Furthermore this means the inner product is lienar in both arguments.
- Now suppose $\mathbb{F} = \mathbb{C}$. Then

$$\langle \lambda u + \mu v, w \rangle = \bar{\lambda} \langle u, w \rangle + \bar{\mu} \langle v, w \rangle.$$

**Definition.** Let $v$ be an inner product space over $\mathbb{R}$.

- The **norm** of a vector $v$ is $\|v\| = \sqrt{\langle v, v \rangle} \geq 0$.
  If $\|v\| = 1$ then we say $v$ is **unit-length**, and $\frac{v}{\|v\|}$ is the **normalisation** of $v$.
- Let $v, w \in V$. If $\langle v, w \rangle = 0$, then $v$ is **orthogonal** to $w$, written $v \perp w$.

## 6.2 Cauchy-Schwarz Inequality

**Theorem.** Let $V$ be an inner product space. For $v, w \in V$,

$$|\langle v, w \rangle| \leq \|v\| \, \|w\|$$

with equality if and only if $v$ and $w$ are dependent.

**Corollary.** (Triangle Inequality) For $v, w \in V$ we have

$$\|v + w\| \leq \|v\| + \|w\|$$

with equality if and only if $v$ and $w$ are dependent.

## 6.3 Orthogonality

**Definition.** A list $u_1, ..., u_k$ in an inner product space $V$ is **orthonormal** if for $1 \leq i, j \leq k$

$$\langle u_i, u_j \rangle = \delta_{ij} = \begin{cases} 1 & i = j \\ 0 & 1 \neq j \end{cases}.$$

**Remark.** A list is orthonormal if every $u_i$ has unit length and distinct $u_i, u_j$ are orthogonal.

**Lemma.** For an orthonormal basis $u_1, ..., u_n$ in an inner product space $V$, for a vector $v \in V$ we have

$$v = \sum_{i=1}^{n} \langle u_i, v \rangle u_i.$$

**Lemma.** Any orthonormal list is linearly independent.

**Proposition.** If $u_1, ..., u_n$ is an orthonormal basis for $V$ and

$$v = \sum_{i=1}^{n} x_i u_i$$
$$w = \sum_{i=1}^{n} y_i u_i$$

then

$$\langle v, w \rangle = \sum_{i=1}^{n} \bar{x}_i y_i = x \cdot y$$

**Proof.** Expand using sesquilinearity.

$$\begin{aligned} \langle v, w \rangle &= \langle \sum_{i=1}^{n} x_i u_i, \sum_{j=1}^{n} y_j u_j \rangle \\ &= \sum_{i,j=1}^{n} \bar{x}_i y_i \langle u_i, u_j \rangle \\ &= \sum_{i=1}^{n} \bar{x}_i y_i. \end{aligned}$$

□

**Theorem.** (Gram-Schmitt Orthogonalisation) Let $v_1, ..., v_n$ be an independent list in an inner product space $V$. Then there exists an orthonormal list $u_1, ..., u_n$ such that for $1 \leq k \leq u$ we have

$$span\ \{v_1, ..., v_k\} = span\ \{u_1, ..., u_k\}.$$

We construct $u_1, ..., u_k$ inductively.

For $k = 1$ we set $w_1 = v_1$, and $u_1 = \frac{w_1}{\|w_1\|}$

For $k > 1$, suppose we have defined $u_1, ..., u_k$ already and set

$$\begin{aligned} w_k &= v_k - \sum_{j=1}^{k-1} \langle_j, v_k \rangle u_j \\ &= v_k - \sum_{j=1}^{n} \frac{\langle w_j, v_k \rangle}{\|w_j\|^2}. \end{aligned}$$

and $u_k = \frac{w_k}{\|w_k\|}$.