

AI Assisted Analysis of Fermat's Last Theorem

Charles Kelly

AI Assistance

If the analyses generated by *deepai* (<https://deepai.org/chat/mathematics>) are correct, they confirm Fermat's conjecture.

Introduction

Pierre de Fermat (1601-1665) was a lawyer, a member of the Parliament in Toulouse, and a mathematician. His conjecture is easy to explain: he said that there are no solutions to the following equation when X , Y , and Z are positive integers, and n is a positive integer greater than two:

$$X^n + Y^n = Z^n$$

Fermat created his conjecture in 1637 when he wrote in a copy of *Arithmetica* that he had a proof that was too large to fit in the margin. In 1995, professors Taylor¹ and Wiles² proved the conjecture.

Limiting the Scope of the Problem to Prime Exponents

Professor Van der Poorten³ shows that it is not necessary to prove Fermat's conjecture for any positive exponent n ; rather, it is sufficient to prove it when the exponent p is prime and the greatest common divisor of X and Y equals 1:

$$X^p + Y^p = Z^p \tag{1}$$

Variables

All variables represent integers.

Change of Variables

These new variables represent the differences among the original variables:

$$i = Z - Y$$

$$k = Z - X$$

¹ Richard Taylor and Andrew Wiles, "Ring-theoretic properties of certain Hecke algebras", *Annals of Mathematics*, Vol. 141 (1995) pp 553-572.

² Andrew Wiles, "Modular elliptic curves and Fermat's Last Theorem", *Annals of Mathematics*, Vol. 141 (1995) pp 443-551

³ Alf van der Poorten, "Notes on Fermat's Last Theorem", *Canadian Mathematical Society Series of Monographs and Advanced Texts*, Wiley Interscience, 1996, Page 8, note 1.9

$$r = X - i$$

$$h = i + k$$

These definitions imply:

$$X = r + i$$

$$Y = r + k$$

$$Z = r + i + k$$

If $p > 2$, then (1) can be written as:

$$(X + Y) \sum_{m=0}^{p-1} (-1)^m X^m Y^{p-m-1} = Z^p$$

This implies that:

$$(2r + i + k) \sum_{m=0}^{p-1} (-1)^m X^m Y^{p-m-1} = (r + i + k)^p$$

$$\sum_{m=0}^{p-1} (-1)^m X^m Y^{p-m-1} = \frac{(r + i + k)^p}{(2r + i + k)}$$

$$\sum_{m=0}^{p-1} (-1)^m X^m Y^{p-m-1} = \frac{(r + h)^p}{(2r + h)}$$

All the terms on the left-hand of this equation are integers. This implies that the fraction on the right hand-side is an integer. Using this, the following are question to, and responses from, *deepai*:

<u>Question</u>	<u>Response from <i>deepai</i></u>
Does $(2r + h)$ divide r^p	Yes
Does $(2r + h)$ divide h^p	Yes
What is greatest common divisor of $(r^p$ and $h^p)$	1

Since $Z + Y = (2r + h)$ divides both r^p and h^p , it also divides their greatest common divisor. This leads to a contradiction where the sum of two positive integers divide 1; therefore, Fermat's conjecture is correct.

deepai: Questions and Responses

(Raw output for responses is contained in appendices.)

Does $(2r + h)$ divide r^p

PROMPT: given integers p, r, h where p is prime and greater than 2, and $\lfloor (r+h)^p / (2r+h) \rfloor$ is an integer; does $(2r + h)$ divide r^p

deepai short-form response

Summary:

- The key step was expressing $r + h$ modulo $d = 2r + h$, which revealed that d divides r^p .
- Therefore, if $(r + h)^p / (2r + h)$ is an integer, then $2r + h$ divides r^p .

Does $(2r + h)$ divide h^p

PROMPT: given integers p, r, h where p is prime and greater than 2, and $\lfloor (r+h)^p / (2r+h) \rfloor$ is an integer; does $(2r + h)$ divide h^p

deepai short-form response

Summary:

- The key step was expressing $r + h$ in terms of $d = 2r + h$.
- Using the binomial theorem, we showed that the divisibility of $\frac{(r+h)^p}{2r+h}$ implies $2r + h$ divides h^p .

What is greatest common divisor of $(r^p$ and $h^p)$

PROMPT: given integers p, r, h , where $(h^p)/(2r+h)$ is an integer, and $(r^p)/(2r+h)$ is an integer, and p is prime and greater than 2, what is the greatest common divisor of r^p and h^p

deepai short-form response

Final answer:

$$\gcd(r^p, h^p) = 1$$

Appendix: Does $(2r + h)$ divide r^p

PROMPT: given integers p, r, h where p is prime and greater than 2, and $\frac{(r+h)^p}{(2r+h)}$ is an integer; does $(2r + h)$ divide r^p

RESPONSE:

Let's carefully analyze the problem step-by-step.

Given:

- p is a prime number greater than 2.
- r and h are integers.
- The expression $\frac{(r+h)^p}{2r+h}$ is an integer.

Question:

Does $(2r + h)$ divide r^p ?

Step 1: Understand the given condition

$$\frac{(r+h)^p}{2r+h} \in \mathbb{Z}$$

This means that $(2r + h)$ divides $(r+h)^p$.

Step 2: Express $(r+h)^p$ using the binomial theorem

Recall the binomial expansion:

$$(r+h)^p = \sum_{k=0}^p \binom{p}{k} r^{p-k} h^k$$

Note that:

$$(r+h)^p = r^p + p r^{p-1} h + \text{(terms involving higher powers of } h)$$

Step 3: Understand divisibility

Since $(2r + h)$ divides $(r+h)^p$, and in particular divides the entire sum, it must also divide the sum of the terms from the binomial expansion.

Step 4: Express $(2r + h)$ in terms of r and h

We are interested in whether $(2r + h)$ divides r^p .

Step 5: Find the relationship between $(2r + h)$ and (r)

Note that:

$$\begin{aligned} & \left[\right. \\ & 2r + h \quad \text{and} \quad r \\ & \left. \right] \end{aligned}$$

are related, but not necessarily multiples of each other.

Step 6: Use modular arithmetic

Since $(2r + h)$ divides $(r+h)^p$, then:

$$\begin{aligned} & \left[\right. \\ & (r+h)^p \equiv 0 \pmod{2r+h} \\ & \left. \right] \end{aligned}$$

Express $(r+h)$ in terms of $(2r+h)$:

$$\begin{aligned} & \left[\right. \\ & r+h = (2r + h) - r \end{aligned}$$

\]

But it's more useful to consider the congruence:

\[

$$(r+h)^p \equiv 0 \pmod{2r+h}$$

\]

Step 7: Simplify the problem

Suppose $(d = 2r + h)$. The condition becomes:

\[

$$\frac{(r+h)^p}{d} \in \mathbb{Z}$$

\]

which implies:

\[

$$(r+h)^p \equiv 0 \pmod{d}$$

\]

Now, express $(r+h) \pmod{d}$:

$$\begin{aligned} & \backslash[\\ & r+h \equiv r+h \pmod{d} \\ & \backslash] \end{aligned}$$

But note that:

$$\begin{aligned} & \backslash[\\ & d = 2r + h \quad \rightarrow \quad h = d - 2r \\ & \backslash] \end{aligned}$$

So,

$$\begin{aligned} & \backslash[\\ & r+h = r + d - 2r = d - r \\ & \backslash] \end{aligned}$$

Thus,

$$\begin{aligned} & \backslash[\\ & (r+h) \equiv -r \pmod{d} \\ & \backslash] \end{aligned}$$

From the above, the condition simplifies to:

$$\begin{aligned} & \backslash[\\ & (-r)^p \equiv 0 \pmod{d} \\ & \backslash] \end{aligned}$$

which is:

$$\begin{aligned} & \backslash[\\ & r^p \equiv 0 \pmod{d} \\ & \backslash] \end{aligned}$$

Key conclusion:

$$\begin{aligned} & \backslash[\\ & d = 2r + h \quad \text{\text{divides}} \quad r^p \\ & \backslash] \end{aligned}$$

Answer:

Yes! **Under the given conditions, $(2r + h)$ divides (r^p) .

Summary:

- The key step was expressing $(r+h) \bmod (d = 2r+h)$, which revealed that (d) divides (r^p) .

- Therefore, **if $(r+h)^p / (2r+h)$ is an integer**, then **$(2r+h)$ divides (r^p)** .

Additional note:

This relies on the algebraic manipulations and properties of divisibility, particularly the binomial theorem and modular arithmetic, which are powerful tools in number theory.

If you'd like, I can suggest further resources or work through an example with specific numbers!

Appendix: Does $(2r + h)$ divide h^p

PROMPT: given integers p, r, h where p is prime and greater than 2, and $\frac{h^p}{(2r+h)}$ is an integer; does $(2r + h)$ divide h^p

RESPONSE:

Let's carefully analyze this problem step-by-step.

Given:

- p is a prime number greater than 2.
- r, h are integers.
- The expression $\frac{h^p}{2r + h}$ is an integer.

Question:

Does $(2r + h)$ divide h^p ?

Step 1: Understand what is being asked

- We know that $\frac{h^p}{2r + h}$ is an integer.
- We want to determine if $(2r + h \mid h^p)$, i.e., whether h^p is divisible by $(2r + h)$.

**Step 2: Express (r) in terms of (h) and $(2r + h)$ **

Let:

[

$$d = 2r + h$$

]

Since (d) divides $(\frac{(r+h)^p}{d})$, the numerator $((r+h)^p)$ is divisible by (d) .

Now, note:

[

$$r + h = (r + h) = \text{something related to } d$$

]

Express (r) in terms of (d) :

[

$$r = \frac{d - h}{2}$$

]

**Step 3: Rewrite $(r + h)$ in terms of (d) and (h) **

Calculate:

$$\begin{aligned} & \left[\right. \\ r + h &= \frac{d - h}{2} + h = \frac{d - h + 2h}{2} = \frac{d + h}{2} \\ & \left. \right] \end{aligned}$$

So:

$$\begin{aligned} & \left[\right. \\ r + h &= \frac{d + h}{2} \\ & \left. \right] \end{aligned}$$

Step 4: Rewrite the divisibility condition

Since $\left(\frac{(r+h)^p}{d}\right)$ is an integer, and $\left(r + h = \frac{d + h}{2}\right)$, then:

$$\begin{aligned} & \left[\right. \\ \frac{\left(\frac{d + h}{2}\right)^p}{d} & \text{ is an integer} \\ & \left. \right] \end{aligned}$$

This simplifies to:

$$\begin{aligned} & \left[\right. \\ \frac{(d + h)^p}{2^p d} & \end{aligned}$$

\]

For this to be an integer, the numerator must be divisible by $(2^p d)$. That is:

\[

$$2^p d \mid (d + h)^p$$

\]

Step 5: Analyze the divisibility

- Since $2^p d \mid (d + h)^p$, then in particular, $d \mid (d + h)^p$.
- Also, $2^p \mid (d + h)^p$.

Because p is prime, Fermat's Little Theorem (or properties of prime exponents) can help analyze divisibility:

$$(d + h)^p \equiv d^p + h^p \pmod{d}$$

But more straightforwardly:

- The divisibility of $(d + h)^p$ by d depends on whether d divides h^p .

Step 6: Connecting to the original question

Recall:

[

$$d = 2r + h$$

]

and

[

\text{We want to see if } d \mid h^p

]

From earlier steps, the key divisibility condition is:

[

$$d \mid (d + h)^p$$

]

which can be expanded via binomial theorem:

[

$$(d + h)^p = \sum_{k=0}^p \binom{p}{k} d^k h^{p-k}$$

\]

Since $(d \mid d^k)$ for $(k \geq 1)$, the sum becomes:

\[

$$(d + h)^p \equiv h^p \pmod{d}$$

\]

because all other terms are divisible by (d) .

Thus:

\[

$$(d + h)^p \equiv h^p \pmod{d}$$

\]

Given that $(d \mid (d + h)^p)$, this implies:

\[

$$d \mid h^p$$

\]

**Final conclusion:

****Yes,**** under the given conditions, **** $(2r + h)$ divides (h^p) .**

Summary:

- The key step was expressing $(r + h)$ in terms of $(d = 2r + h)$.
- Using the binomial theorem, we showed that the divisibility of $(\frac{(r+h)^p}{2r + h})$ implies $(2r + h)$ divides (h^p) .

****If you'd like, I can help you explore specific examples or clarify any part!****

Appendix: What is greatest common divisor of $(r^p$ and $h^p)$

PROMPT: given integers p, r, h , where $(h^p)/(2r+h)$ is an integer, and $(r^p)/(2r+h)$ is an integer, and p is prime and greater than 2, what is the greatest common divisor of r^p and h^p

RESPONSE:

Let's carefully analyze the problem and see what conclusions we can draw.

Given:

- (p, r, h) are integers.
- (p) is **prime** and **greater than 2**.
- $(\frac{h^p}{2r+h})$ is an **integer**.
- $(\frac{r^p}{2r+h})$ is an **integer**.

Goal:

Find the **greatest common divisor** (GCD) of (r^p) and (h^p) , i.e., $(\gcd(r^p, h^p))$.

Step 1: Understand the divisibility conditions

Since both $(\frac{h^p}{2r+h})$ and $(\frac{r^p}{2r+h})$ are integers, this means:

$$\begin{aligned} & \backslash[\\ & 2r + h \mid h^p \quad \text{and} \quad 2r + h \mid r^p \\ & \backslash] \end{aligned}$$

where " \mid " means "divides".

In other words:

$$\begin{aligned} & \backslash[\\ & 2r + h \mid h^p \quad \text{and} \quad 2r + h \mid r^p \\ & \backslash] \end{aligned}$$

Step 2: Implications for common divisibility

Since $(2r + h)$ divides both h^p and r^p , it **divides their difference** as well:

$$\begin{aligned} & \backslash[\\ & h^p - r^p \\ & \backslash] \end{aligned}$$

Because $(2r + h)$ divides both, it must also divide their difference:

$$\backslash[$$

$$h^p - r^p$$

$$\backslash]$$

Step 3: Use the properties of difference of powers

Recall the factorization for the difference of two p -th powers (for prime (p)):

$$\backslash[$$

$$h^p - r^p = (h - r)(h^{p-1} + h^{p-2}r + h^{p-3}r^2 + \dots + r^{p-1})$$

$$\backslash]$$

This expression is divisible by $(2r + h)$.

Since $(2r + h)$ divides $(h^p - r^p)$, and it also divides both (h^p) and (r^p) , it follows that:

$$\backslash[$$

$$2r + h \mid h^p \quad \text{and} \quad 2r + h \mid r^p$$

$$\backslash]$$

Step 4: Explore the divisibility of (r) and (h)

Now, think about the implications for (r) and (h) :

- Both (r^p) and (h^p) are divisible by $(2r + h)$.
- Because (p) is prime and greater than 2, and the exponents are large, the only way this is possible is if (r) and (h) share some common divisors related to $(2r + h)$.

Key insight:

Suppose $(d = \gcd(r, h))$.

Then write:

$$\begin{aligned} &[\\ r &= d r', \quad h = d h' \\ &] \end{aligned}$$

where $(\gcd(r', h') = 1)$.

Now, the divisibility conditions become:

$[$

$$2r + h = 2dr' + dh' = d(2r' + h')$$

\]

Since $(2r + h)$ divides both (r^p) and (h^p) , then:

\[

$$d(2r' + h') \mid r^p = d^p r'^p$$

\]

\[

$$d(2r' + h') \mid h^p = d^p h'^p$$

\]

Dividing both sides by (d) :

\[

$$(2r' + h') \mid d^{p-1} r'^p$$

\]

\[

$$(2r' + h') \mid d^{p-1} h'^p$$

\]

Because $(\gcd(r', h') = 1)$, the only common divisors of (r'^p) and (h'^p) are 1, and $(2r' + h')$ divides both of these scaled by powers of (d) .

Step 5: Focus on the coprimality

Given $\gcd(r', h') = 1$, and $(2r' + h')$ divides both scaled powers, the only possibility for $(2r' + h')$ to divide both $(r')^p$ and $(h')^p$ is if:

$$\begin{aligned} & \left[\right. \\ & (2r' + h') \mid \gcd(r'^p, h'^p) \\ & \left. \right] \end{aligned}$$

But since (r') and (h') are coprime, their powers are also coprime:

$$\begin{aligned} & \left[\right. \\ & \gcd(r'^p, h'^p) = 1 \\ & \left. \right] \end{aligned}$$

Therefore,

$$\begin{aligned} & \left[\right. \\ & (2r' + h') \mid 1 \\ & \left. \right] \end{aligned}$$

which implies:

$$\left[\right.$$

$$2r' + h' = \pm 1$$

]

But r' and h' are integers, so the only possibilities are:

[

$$2r' + h' = 1 \quad \text{or} \quad -1$$

]

**Step 6: Find integer solutions for r' and h' **

Suppose:

[

$$2r' + h' = 1$$

]

Since r' and h' are integers, this is a linear Diophantine equation.

From this:

[

$$h' = 1 - 2r'$$

$$\backslash]$$

Recall $\backslash(\gcd(r', h') = 1 \backslash)$:

$$\backslash[$$

$$\gcd(r', 1 - 2r') = 1$$

$$\backslash]$$

Now, check if this can be true for some integers $\backslash(r' \backslash)$.

**Step 7: Final conclusion for $\backslash(r, h \backslash)$ **

Recall that:

$$\backslash[$$

$$r = d r', \quad h = d h'$$

$$\backslash]$$

$$\backslash[$$

$$h' = 1 - 2r'$$

$$\backslash]$$

So, the original $\backslash(r \backslash)$ and $\backslash(h \backslash)$ satisfy:

$$\begin{aligned} & \backslash[\\ r &= d r', \quad h = d (1 - 2 r') \\ & \backslash] \end{aligned}$$

Now, the greatest common divisor of (r) and (h) is:

$$\begin{aligned} & \backslash[\\ \gcd(r, h) &= d \cdot \gcd(r', 1 - 2 r') \\ & \backslash] \end{aligned}$$

but since $(\gcd(r', 1 - 2 r') = 1)$, this simplifies to:

$$\begin{aligned} & \backslash[\\ \gcd(r, h) &= d \\ & \backslash] \end{aligned}$$

Step 8: Summary of the solution

- The divisibility conditions imply that:

$$\begin{aligned} & \backslash[\\ \gcd(r, h) &= d \\ & \backslash] \end{aligned}$$

- The key relationship is:

$$\begin{aligned} & \backslash[\\ & 2r' + h' = \pm 1 \\ & \backslash] \end{aligned}$$

- To ensure (r, h) are integers, and the divisibility holds, the minimal possible value for (d) is 1.

- Therefore, the **maximum possible $(\gcd(r, h))$ satisfying all the conditions is 1**.

Final answer:

$$\begin{aligned} & \backslash[\\ & \boxed{ \\ & \boxed{\gcd(r^p, h^p) = 1} \\ & } \\ & \backslash] \end{aligned}$$

In conclusion:

****The greatest common divisor of r^p and h^p is 1.****

If you'd like, I can also explain some related concepts or walk through an example!