

Set up

- ▶ (M^n, g) is a Riemannian manifold.
- ▶ $\Omega \subset M^n$ is a precompact domain with smooth boundary.

Dirichlet Problem

Given $\Omega \subset M^n$ there exists a function $u \in C_c^\infty(\Omega)$ such that

$$\begin{cases} \Delta u = -1 & \text{in } \Omega, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

Then we can consider $\mathcal{T}(\Omega) = \int_{\Omega} u \, dV_g$.

Understanding $\mathcal{T}(\Omega)$

- **Torsional rigidity** $\mathcal{T}(\Omega)$ give us the torque required when twisting an elastic beam with the domain of study as uniform cross section.




Figure: Sketch of a beam with cross section Ω

- **Mean exit time** expected time of a particle starting at a point $x \in \Omega$ will reach the boundary of the domain following a Brownian Motion.



Figure: Example of a brownian ant.

Variational approach

$$\mathcal{T}(\Omega) = \sup \left\{ \frac{\left(\int_{\Omega} u \, dV \right)^2}{\int_{\Omega} \|\nabla u\|^2 \, dV} : u \in C_0^\infty(\Omega) \text{ and } \int_{\Omega} \|\nabla u\| \, dV \neq 0 \right\}.$$

$$\mathcal{T}(\Omega) = \inf \left\{ \int_{\Omega} \|X\|^2 \, dV : X \in \mathfrak{X}(\Omega) \text{ with } \operatorname{div}_g X = -1 \right\}.$$

Intrinsic flows

Ricci flow, $g(t)$, are smooth paths in the space of metrics, $\mathcal{M}(M^n)$. The Ricci flow is a path governed by

$$\frac{\partial g_t}{\partial t} = -2\operatorname{Ric}_g(t) \in T_{g_t}\mathcal{M}(M).$$

Extrinsic flow

Inverse Mean Curvature Flow let

$\varphi : M^n \times [0, T_{\max}] \mapsto \mathbb{R}^{n+1}$ a family of isometric immersions such that

$$\partial_t \varphi(p, t) = -\vec{H}_t(p) \left\| \vec{H}_t(p) \right\|^{-2}.$$

Then the evolution equations under this two flows are,

Ricci flow	IMCF
$\partial_t g_{ij} = -2 \operatorname{Ric}_{ij}$	$\partial_t g_{ij} = \frac{2 h_{ij}}{H}$
$\partial_t d\mu_t = -R d\mu_t$	$\partial_t d\mu_t = d\mu_t$
$\partial_t (\operatorname{div}_{g_t} X) = -X(R)$	Preserved

Theorem for Ricci flow

Let $(M^n, (g_t)_{t \in [0, T_{\max}]})$ be a Ricci flow solution. Suppose that for any $t \in [0, T_{\max})$ and any $p \in M$ $\operatorname{Scal}_{g_t}(p) = b(t)$. Suppose moreover that for any $t \in [0, T_{\max})$, any $p \in M$ and any $v \in T_p M$, there exists $A, B : \mathbb{R} \rightarrow \mathbb{R}$ such that $A(t)g_t(v, v) \leq \operatorname{Ric}_{g_t}(v, v) \leq B(t)g_t(v, v)$. Then for any precompact domain $\Omega \subset M$ with smooth boundary

$$e^{\int_0^t (-b(s) - 2B(s)) ds} \mathcal{T}(\Omega_0) \leq \mathcal{T}(\Omega_t) \leq e^{\int_0^t (-2A(s) - b(s)) ds} \mathcal{T}(\Omega_0).$$

Which implies

$$e^{-2 \int_0^t B(s) ds} \frac{\mathcal{T}(\Omega_0)}{V(\Omega_0)} \leq \frac{\mathcal{T}(\Omega_t)}{V(\Omega_t)} \leq e^{-2 \int_0^t A(s) ds} \frac{\mathcal{T}(\Omega_0)}{V(\Omega_0)}.$$

- ▶ Let $(SU(2), g = \epsilon A_0 \Theta^1 \otimes \Theta^1 + B_0 \Theta^2 \otimes \Theta^2 + C_0 \Theta^3 \otimes \Theta^3)$ such that $\delta := \frac{B_0 - \epsilon A_0}{\epsilon A_0} < 1$.

$$\left[1 - \frac{4(1 + 6\delta)t}{B_0} \right]^{\frac{5(1+\delta)^3}{2(1+6\delta)}} \leq \frac{\mathcal{T}(\Omega_t)}{\mathcal{T}(\Omega_0)} \leq \left[1 - \frac{4t}{B_0} \right]^{\frac{5(1-\delta)}{2(1+\delta)}}$$

Theorem for IMCF

Let $\varphi : M \times [0, T_{\max}) \rightarrow \mathbb{R}^{n+1}$ by an inverse mean curvature flow from M to \mathbb{R}^{n+1} such that $\varphi_t(M) = \Sigma$ is strictly convex hypersurface of \mathbb{R}^{n+1} . Then the function

$$t \mapsto (V(\Omega_t))^{-3} \cdot \mathcal{T}(\Omega_t)$$

is non-increasing for $t \in [0, T_{\max})$, and the function

$$t \mapsto (V(\Omega_t))^{-1} \cdot \mathcal{T}(\Omega_t)$$

is non-decreasing for $t \in [0, T_{\max})$.

- ▶ Let Σ be a strictly convex free boundary disk-type hypersurface in the ball $\mathbb{B} \subset \mathbb{R}^{n+1}$. Then we have

$$\frac{\mathcal{T}(\Sigma)}{V^3(\Sigma)} \geq \frac{\mathcal{T}(\mathbb{D})}{V^3(\mathbb{D})}, \quad \frac{\mathcal{T}(\Sigma)}{V(\Sigma)} \leq \frac{\mathcal{T}(\mathbb{D})}{V(\mathbb{D})}$$

Bibliography

- ▶ Georg Polya and Gábo Szegő. *Isoperimetric Inequalities in Mathematical Physics*.
- ▶ Alexander Grigor'yan. *Analytic and geometric background of recurrence and non-explosion of the Brownian Motion on riemannian manifolds*.
- ▶ Vicent Gimeno i García and G.. *Torsional rigidity under geometric flows*.