

BULLET GROUPS AS A TEST OF Λ CDM

J. G. FERNÁNDEZ-TRINCADO^{1,2,3}, J. E. FORERO-ROMERO¹ AND T. VERDUGO³

¹ Departamento de Física, Universidad de los Andes, Cra. 1 No. 18A-10, Edificio Ip, Bogotá, Colombia

² Institute Utinam, CNRS UMR6213, Université de Franche-Comté, OSU THETA de Franche-Comté-Bourgogne, Besançon, France

³ Centro de Investigaciones de Astronomía, AP 264, Mérida 5101-A, Venezuela

Submitted for publication in *ApJL*

ABSTRACT

We estimate the expected distribution of displacements between the two dominant dark matter peaks in halos within a mass range corresponding to galaxy groups. We find that the probability of finding a system similar with displacements of $\sim 400 h^{-1} \text{kpc}$ is between 40% for $z=0$ and 60% for $z=1$ and Λ CDM standard model, which correspond to the observational constraint the object SL2S J08544-0121 that is a gravitational lens found in the SL2S and located at $z=0.35$. Given the larger abundance of groups with respect to clusters, finding multi-modal groups and baryonic-dark matter displacements.

Subject headings: cosmology: theory – dark matter

1. INTRODUCTION

The Bullet Cluster provided a new kind of observational evidence of the existence of dark matter. Quantifying the displacement between dark matter and the dominant baryonic component (hot X-ray emitting gas) has been used to test the CDM paradigm itself by quantifying the substructure velocity required to produce such displacement and also by estimating the expected abundance of such events in the Universe.

Since that time there are other observations of other Bullet-like systems [...].

Recently (Gastaldello et al. 2014) observed baryonic-DM displacement of $124 \pm 20 \text{ kpc}$ in a group-like system with a total mass $2.4 \pm 0.6 \times 10^{14} M_{\odot}$. Systems of this mass are ~ 10 times more massive than cluster systems in the mass range $> 10^{15} h^{-1} M_{\odot}$, this opens up the possibility of observationally finding bullet groups in a fair amount to impose constraints on Λ CDM. This greater abundance has to be weighted by the fraction of systems that present large displacements. Such study has been performed for clusters but not for lower mass systems.

In this Letter we present prediction for the abundance of group-like systems that might show a DM-baryon displacement. To this end we use a N-body cosmological simulation with such a resolution that allows us to identify multimodal dark matter clumps in the circular velocity range $300 - 1000 \text{ km s}^{-1}$.

This paper is organized as the follows. In Section 2 we present the simulation and the halo catalogs used in this work. We continue in Section 3 with the geometry of the problem at hand and the measurements setup. Next in Section 4 we present our results to finish with a discussion and conclusion in Sections 5 and 6.

2. SIMULATION, HALO CATALOGS AND PAIRS

We use the Bolshoi Run, a cosmological DM only simulation over a cubic volume of $250 \text{ comoving } h^{-1} \text{Mpc}$ on a side. The simulation uses the ART code to follow the evolution of a dark matter density field sampled with 2024^3 from $z=80$ to $z=0$ [...]. The cosmology used corresponds to the spatially flat concordance model with the following parameters: the density parameter for

matter (dark matter+baryons) $\Omega_m = 0.27$, the density parameter for baryonic matter $\Omega_b = 0.0469$, the density parameter for dark energy $\Omega_{\Lambda} = 0.73$, the Hubble parameter $h = 0.7$, the normalization of the Power spectrum $n = 0.95$ and the amplitude of mass density fluctuation (at redshift $z=0$) $\sigma_8 = 0.82$. The number of particles used for each of the DM component was 2048^3 , resulting in a mass resolution of $1.35 \times 10^8 M_{\odot} h^{-1}$. Klypin et al. (2011).

We use halo catalogs constructed using the BDM algorithm. [...]

In the snapshots at redshift $z = 0.0, 0.25, 0.5$ and 1.0 there are XX, XX, XX, XX host halos with circular velocities $V_c > 300 \text{ km s}^{-1}$ and XX, XX, XX, XX sub-halos with circular velocities $V_c > 75 \text{ km s}^{-1}$. These two sets of halos constitute the basis for our analysis.

For each host halo in the sample we find its most massive sub-halo. Each pair host/sub-halo is considered as a potential Bullet Group and is kept for the measurement and analysis described in the next section.

3. BULLET GEOMETRY AND MEASUREMENT SETUP

The Bullet groups is composed by two halos. the host halo and the sub halo. This configuration is described by the position and velocity of the sub-halo in a frame of reference where the main halo is at rest; thus $\vec{v} = \vec{v}_{sub} - \vec{v}_{halo}$ and $\vec{r} = \vec{r}_{sub} - \vec{r}_{halo}$, where the subscripts *host* and *sub* refer to the host and sub-halo in the frame of reference of the simulation, respectively.

The angle between these two vectors characterized by,

$$\mu \equiv \cos(\theta) = \frac{\vec{v} \cdot \vec{r}}{\|\vec{v}\| \|\vec{r}\|} \quad (1)$$

encodes the geometry of the collision, i.e. cases of $|\mu| \approx 1$ can be considered as head-on collisions while $|\mu| \approx 0$ describe a grazing trajectory.

The bullet-like encounter can be instantaneously described by quantities the following quantities the circular velocity of the host and the sub-halo, $V_{c,host}$ and $V_{c,sub}$; the size of the host halo R_{vir} ; the relative position and velocity of the substructure, \vec{v} and \vec{r} ; and the angle between the position and velocity μ .

As a first approximation there are two quantities that are available from observations of these Bullet-like systems. The projected distance between two dominant dark matter clumps and the ratio of the galaxies' luminosities associated to them. From the simulation point of view this can be translated into the 2D projected values of $||\vec{r}||$, its value relative to the virial radius $X_{2D} = ||\vec{r}||_{2D}/R_{vir}$ and the ratio of the circular velocities of the two clumps $V_{c,sub}/V_{c,host}$. In a higher degree of detail, in order to gain better insight we use the sub-structure velocity as a fraction of the host's circular velocity, $||\vec{r}||/V_{c,host}$, as a measure of the strength of the interaction. Finally, we also measure the geometry of each interaction through the values for μ .

All the physical quantities described above can be used to describe the three main stages in a bullet-like encounter. First, the sub-halo crosses the virial radius of the host halo starting a head on collision, $||\vec{r}||/R_{vir} \approx 1$ and $\mu < 0.0$. Second, as the sub-halo crosses for the first time the center of the host halo $||\vec{r}||/R_{vir} < 1.0$ and $\mu > 0.0$. Third, as the sub-halo reaches apogee and comes back to the center of the halo $||\vec{r}||/R_{vir} < 1.0$ and $\mu < 0.0$. We use these quantities in Section XXX to fully characterize the different kind of interactions observed in the Bolshoi simulation.

4. RESULTS

4.1. Displacements and Relative Circular Velocities

The main result of this paper is summarized in Figure 1, it presents the integrated probability distribution for the displacement between the center of the host halo and its dominant sub-halo. The left panel shows the displacement in physical units and the right panel as a fraction of the virial radius of the host halo.

Figure 1 shows the results for two different populations; groups with $300 \text{ km s}^{-1} < V_{c,host} < 700 \text{ km s}^{-1}$ and clusters with $V_{c,host} > 700 \text{ km s}^{-1}$. Additionally, this is presented for all redshifts $z = 0.0, 0.25, 0.5$ and 1.0 .

The panel with the physical displacements also shows a vertical stripe with the estimated displacement for the Bullet-group reported by Gastaldello et al. (2014). Considering this system as consistent with the groups sample we see that a fraction of $\sim 15 \pm 2\%$ of the groups should present a displacement equal or larger than the one estimated for SL2S J08544-0121, this fraction is close to $45 \pm 2\%$. The panel with the normalized displacements shows a distribution that can be considered close to universal in the sense that the two samples (groups and clusters) at all redshifts present a similar trend.[...]

Figure 2 shows 2D histograms in a plane composed by the ratio of the two circular velocities $V_{c,sub}/V_{c,host}$ and the physical displacements. This is revealing because these two quantities can be constrained by observations. The displacement is a direct observable, while the ratio of the circular velocities can be estimated from lensing studies or approximated by the ratio of the total galaxy luminosity associated with the galaxy peaks.

To construct this figure we co-add all the halos in the sample (left, groups; right, clusters) at all redshifts. We stack the data because we do not observe any strong time evolution, additionally this allows us to increase the signal in each bin. Overplotted there is a circle with

error bars that represents the observational estimates for the system SL2S J08544-0121.

4.2. Relative Velocities

In Figure 3 we present the integrated probability of the relative peculiar velocities of the sub-halos with respect to the host halo. The left panel presents this velocity in physical units while the right panel presents them as a fraction of the circular velocity of the host halo.

The panel with the normalized velocities shows that the distribution of sub-halo velocities is close to universal. Regardless of the mass of the host halo and the redshift the integrated distributions lie very close to each other.

The median of this distribution is located at $v/V_{c,host} = 1.1$. We also note the strong break at $v/V_{c,host} = 3.0$ that is present in the data from the group sample that allows us to probe fractions on the order of 10^{-4} . This break is located close to the escape velocity of $v/V_{c,host}$ for dark matter halos following a NFW profile with a concentration value $c \approx 6$ (Hayashi & White 2006).

4.3. Collision Geometries

Figure ?? presents the geometry of the bullet groups in our relative geometry of the collision using the variables μ and X_{off} .

5. DISCUSSION

Strictly speaking our results apply to multimodal groups and the expected separation between dark matter clumps. [...]

Expected displacement between dark and baryonic components are better described by systems where the collision already occurred.[...] (comparison with the old bullet clusters paper?)

The expectations to quantify this effects observationally are high. [...]

In order to facilitate the reproducibility and reuse of our results we have made available all the data and the source code available in a public repository. [...]

6. CONCLUSIONS

We selected four snapshots of the simulation for four different redshifts ($z = 0, z = 0.25, z = 0.5$ and $z = 1$) based in the distribution of redshift in the Figure 1 by Verdugo & Foex (2014), and for each sample in redshift we selected the host halo with circular velocities greater than 300 km s^{-1} and was split in two principal groups: The first group correspond to host halo with circular velocities between 300 km s^{-1} to 700 km s^{-1} with mass in the range of $10^{12} M_{\odot}$ to $10^{14} M_{\odot}$ in the range of mass of the Bullet Groups, and second group correspond to host halo with circular velocities greater than 700 km s^{-1} with mass $\geq 10^{14} M_{\odot}$, in the range of mass of the Bullet Clusters, in the Table ?? is shown the two groups selected in this work. For each group, we classified the corresponding substructures most massive and associated with the corresponding host halo. In this work, we estimate the expected distribution of displacements ($d_{real,(X,Y)}$) in the projection 2-D that can be estimated by observations, these displacements correspond to the separation between the




FIG. 1.— Integrated probability distribution for the displacement between the host halo and its dominant sub-halo. The left panel shows the results in terms of the physical displacements and the right panel the same displacement normalized by the virial radius of the host halo. The continuous line corresponds to the halos in the group sample $V_{\text{circ,host}} > 700 \text{ km s}^{-1}$ and the dashed lines to the sample $V_{\text{circ,host}} < 700 \text{ km s}^{-1}$. The vertical stripe corresponds to the estimate of the separation between the two dark matter clumps in the results reported by Gastaldello et al. (2014) for the group SL2S J08544-0121. The right panel shows the same results normalized by the virial radius of the host halo.




FIG. 2.— 2D histogram in the plane $V_{\text{c,sub}}/V_{\text{c,host}} - d_{2D}$. The left panel corresponds to groups and the right panel to clusters. The circle with error bars corresponds to SL2S J08544-0121 data reported from Muñoz et al. (2013) and Gastaldello et al. (2014). The data used to construct the histograms integrates the objects at all redshifts.




FIG. 3.— Integrated probability distribution for the relative velocity of the sub-halo with respect to its host. The left panel shows the results in physical units while the right panel show the same values normalized by the circular velocity of the host halo. The line coding follows the same structure as Figure ??

minimal potential of the host halo and the minimal potential of the substructure, both are dark matter distributions.

In order to explore the distribution of displacements expected in the observational, we define a new parameter given by:

$$\frac{v_{circ,sub}}{v_{circ,halo}} = 0.5 \quad (2)$$

Figure ?? shows the scatter plot of the parameter $\left(\frac{v_{circ,sub}}{v_{circ,halo}}\right)$ vs the displacement ($d_{2d,(X,Y)}$) between the host halo and the substructure and for different redshifts. This parameter is consistent with the observations. The red star symbol in the Figure ?? is equal to 0.54 corresponding to the fraction in velocity dispersion in the line-of-sight of the group SL2S SJ08544-0121 ($\sigma_{host,halo} = 341^{+43}_{-109} \text{ km s}^{-1}$ and $\sigma_{substructure} = 185^{+30}_{-62}$), reported by Muñoz et al. (2013). Based in the parameter $\left(\frac{v_{circ,sub}}{v_{circ,halo}}\right) > 0.5$, we estimate the expected distribution of displacements and this is shown in the Figure ?? for the two groups classified in this work.

ACKNOWLEDGEMENTS

The CosmoSim database used in this paper is a service by the Leibniz-Institute for Astrophysics Potsdam (AIP). The BolshoiP simulation was performed within the Bolshoi project of the University of California High-Performance AstroComputing Center (UC-HIPACC) and was run at the NASA Ames Research Center.

REFERENCES

- Gastaldello, F., Limousin, M., & Foex, G. 2014, MNRAS submitted
 Hayashi, E., & White, S. D. M. 2006, MNRAS, 370, L38
 Klypin, A. A., Trujillo-Gomez, S., & Primack, J. 2011, ApJ, 740, 102
 Muñoz, R. P., Motta, V., Verdugo, T., Garrido, F., Limousin, M., Padilla, N., Foëx, G., Cabanac, R., Gavazzi, R., Barrientos, L. F., & Richard, J. 2013, A&A, 552, A80
 Verdugo, T., & Foex, G. 2014, a submitted




FIG. 4.— x