

Table 1. Characteristics of the Total Sample and Stratified by Sedentary Behavior

Variables	Sedentary Behavior			
	All Participants N = 1711711851.64708 ¹	1st Quintile N = 29958395.1796981 ¹	2nd Quintile N = 33670734.6019899 ¹	3rd Quintile N = 32151605.6561111 ¹
Age (yr)	46 ± 17	43 ± 16	47 ± 16	47 ± 18
Sex (M/F)				
1	87,415,471 (51%)	15,630,783 (52%)	17,088,013 (51%)	14,829,748 (46%)
2	84,296,381 (49%)	14,327,612 (48%)	16,582,722 (49%)	17,321,858 (54%)
Race				
1	28,052,894 (16%)	9,465,830 (32%)	5,874,925 (17%)	4,676,042 (14%)
3	107,984,325 (63%)	14,000,697 (47%)	21,266,629 (63%)	19,746,147 (61%)
4	19,041,063 (11%)	3,606,894 (12%)	3,821,580 (11%)	4,271,914 (13%)
5	16,633,570 (9.7%)	2,884,975 (9.6%)	2,707,601 (8.0%)	3,457,503 (11%)
Education				
1	15,793,210 (9.2%)	5,694,517 (19%)	3,127,551 (9.3%)	2,418,206 (7.5%)
2	42,909,327 (25%)	10,319,766 (34%)	9,514,406 (28%)	9,701,677 (30%)
3	52,614,771 (31%)	8,550,189 (29%)	10,832,151 (32%)	9,055,739 (28%)
4	60,380,152 (35%)	5,393,923 (18%)	10,196,626 (30%)	10,975,984 (34%)
Poverty Income Ratio	3.31 ± 1.60	2.60 ± 1.61	3.20 ± 1.57	3.15 ± 1.61
BMI (Kg/m ²)	29 ± 7	28 ± 6	29 ± 7	30 ± 8
Waist Circumference (cm)	99 ± 16	96 ± 15	99 ± 16	100 ± 16
Overweight/Obesity				
0	45,124,061 (26%)	9,729,369 (33%)	7,778,540 (23%)	9,059,969 (28%)

¹ Mean ± SD; n (%)

Variables	Sedentary Behavior			
	All Participants N = 171711851.64708 ¹	1st Quintile N = 29958395.1796981 ¹	2nd Quintile N = 33670734.6019899 ¹	3rd Quintile N = 32151605.6561 ¹
1	125,942,783 (74%)	20,045,684 (67%)	25,743,007 (77%)	23,015,467 (73%)
SBP (mmHg)	120 ± 16	119 ± 16	121 ± 16	120 ± 16
DBP (mmHg)	74 ± 11	73 ± 11	74 ± 11	74 ± 11
Hypertension				
0	100,275,906 (58%)	18,651,392 (62%)	19,541,235 (58%)	18,705,947 (58%)
1	71,435,945 (42%)	11,307,004 (38%)	14,129,500 (42%)	13,445,659 (42%)
Total Cholesterol (mg/dL)	186 ± 40	183 ± 38	192 ± 43	186 ± 40
LDL-Cholesterol (mg/dL)	110 ± 35	108 ± 33	114 ± 35	111 ± 35
HDL-Cholesterol (mg/dL)	54 ± 16	55 ± 18	56 ± 16	54 ± 17
Fasting Triglycerides (mg/dL)	110 ± 95	107 ± 101	112 ± 74	108 ± 74
Low-HDL				
0	124,174,300 (73%)	21,665,620 (73%)	25,334,148 (76%)	22,188,464 (73%)
1	45,407,558 (27%)	7,963,962 (27%)	8,097,475 (24%)	9,705,718 (30%)
Hypertriglyceridemia				
0	114,348,764 (67%)	21,956,064 (74%)	22,771,728 (68%)	20,026,585 (63%)
1	55,465,026 (33%)	7,801,275 (26%)	10,659,896 (32%)	11,867,597 (37%)
Fasting Glucose (mg/dL)	108 ± 30	108 ± 32	111 ± 41	108 ± 29
Insulin (μU/mL)	13 ± 22	12 ± 11	12 ± 12	13 ± 12
¹ Mean ± SD; n (%)				

Variables	Sedentary Behavior			
	All Participants N = 171711851.64708 ¹	1st Quintile N = 29958395.1796981 ¹	2nd Quintile N = 33670734.6019899 ¹	3rd Quintile N = 32151605.65611 ¹
HbA1C (%)	5.61 ± 0.88	5.64 ± 0.87	5.70 ± 1.03	5.59 ± 0.87
HOMA-IR	3.82 ± 8.33	3.23 ± 3.55	3.45 ± 4.82	3.65 ± 4.10
Dysglycaemia or DM2				
0	59,970,981 (35%)	10,881,706 (36%)	10,988,867 (33%)	11,112,780 (34%)
1	111,740,870 (65%)	19,076,690 (64%)	22,681,868 (67%)	21,038,826 (66%)
GGT (U/L)	28 ± 29	28 ± 30	26 ± 25	29 ± 29
ALT (U/L)	22 ± 15	21 ± 15	22 ± 15	23 ± 15
AST (U/L)	21 ± 10	21 ± 10	22 ± 10	21 ± 10
ALT/AST	1.02 ± 0.35	1.00 ± 0.35	1.01 ± 0.33	1.04 ± 0.34
Total Energy Intake (kcal/d)	2,115 ± 799	2,078 ± 878	2,121 ± 812	2,078 ± 878
Smoking Habit				
0	100,647,680 (59%)	17,546,251 (59%)	17,618,416 (52%)	18,910,562 (59%)
1	44,469,963 (26%)	6,208,156 (21%)	10,113,679 (30%)	7,516,717 (23%)
2	26,557,204 (15%)	6,203,988 (21%)	5,938,640 (18%)	5,724,326 (18%)
Alcohol Intake				
0	13,475,724 (7.8%)	2,956,561 (9.9%)	3,127,950 (9.3%)	2,394,443 (7.5%)
1	24,068,933 (14%)	3,951,795 (13%)	4,801,029 (14%)	4,840,003 (15%)
2	134,167,195 (78%)	23,050,039 (77%)	25,741,756 (76%)	24,917,160 (77%)
Sedentary Behavior (h)	354 ± 204	106 ± 40	204 ± 29	294 ± 40

¹ Mean ± SD; n (%)

Variables	Sedentary Behavior			
	All Participants N = 171711851.64708 ¹	1st Quintile N = 29958395.1796981 ¹	2nd Quintile N = 33670734.6019899 ¹	3rd Quintile N = 32151605.65611 ¹
Median CAP (dB/m)	261 ± 61	250 ± 59	263 ± 62	261 ± 61
Liver Stifness (Kpa)	5.47 ± 3.13	5.35 ± 3.29	5.75 ± 4.00	5.57 ± 2.91
MASLD (CAP ≥ 248 db/m)				
0	79,667,134 (46%)	16,202,254 (54%)	14,535,122 (43%)	15,310,807 (47%)
1	92,044,718 (54%)	13,756,141 (46%)	19,135,612 (57%)	16,840,798 (53%)
Hepatic Advanced Fibrosis (LSM ≥ 8 kPa)				
0	158,092,753 (92%)	28,007,872 (93%)	30,710,977 (91%)	29,060,813 (91%)
1	13,619,099 (7.9%)	1,950,523 (6.5%)	2,959,757 (8.8%)	3,090,792 (9.6%)
NAFLD (CAP ≥ 248 db/m)				
0	74,370,555 (43%)	15,157,393 (51%)	13,564,111 (40%)	13,889,200 (43%)
1	97,341,297 (57%)	14,801,002 (49%)	20,106,623 (60%)	18,262,406 (57%)
HSI	38 ± 8	37 ± 8	38 ± 8	39 ± 9
FLI	0.50 ± 0.33	0.45 ± 0.32	0.49 ± 0.32	0.51 ± 0.33
MVPA	1,232 ± 1,810	2,114 ± 2,481	1,653 ± 1,946	1,139 ± 1,710
Work-Related Activity	668 ± 1,143	1,201 ± 1,535	978 ± 1,282	610 ± 1,040
Walking or Bicycling for Transportation	59 ± 239	87 ± 318	74 ± 240	73 ± 330
Leisure-Time Physical Activity	176 ± 263	188 ± 300	179 ± 259	174 ± 266

¹ Mean ± SD; n (%)