

Part III — Mechanics

Based on lectures by Brian
Notes taken by Dexter Chua

Lent 2017-2018

These notes are not endorsed by the lecturers, and I have modified them (often significantly) after lectures. They are nowhere near accurate representations of what was actually lectured, and in particular, all errors are almost surely mine.

Contents

1 Kinematics of a particles moving in a straight line	4
2 Dynamics of a particle moving in a straight line	5
3 Statics of a particle	6
4 Moments	7
5 Vectors	8
6 Kinematics of a particle moving in a straight line or plane	9
7 Centres of mass	10
8 Work , energy and power	11
9 Collisions	12
10 Statics of rigid bodies 1	13
11 Further kinematics	14
11.1 Forces which vary with speed	14
12 Elastic strings and springs	15
12.1 Hooke's Law	15
12.2 Energy stored in an elastic string or spring	15
13 Further dynamics	16
13.1 Impulse of a variable force	16
13.2 Work done by a variable force	16
13.3 Newton's Law of Gravitation	16
13.4 Finding k in $F = \frac{k}{x^2}$	16
13.5 Simple harmonic motion S.H.M.	16
14 Motion in a circle	17
14.1 Angular velocity	17
14.2 Acceleration	17
14.3 Motion in a vertical circle	17
15 Statics of rigid bodies 2	18
15.1 Centre of mass	18
16 Relative motion	19
17 Elastic collisions in two dimensions	20
18 Resisted motion of a particle moving in a straight line	21
19 Damped and forced harmonic motion	22
20 Stability	23

21 Applications of vectors in mechanics	24
22 Variable mass	25
23 Moments of inertia of a rigid body	26
24 Rotation of a rigid body about a fixed smooth axis	27

1 Kinematics of a particles moving in a straight line

2 Dynamics of a particle moving in a straight line

3 Statics of a particle

4 Moments

5 Vectors

6 Kinematics of a particle moving in a straight line or plane

7 Centres of mass

8 Work , energy and power

9 Collisions

10 Statics of rigid bodies 1

11 Further kinematics

11.1 Forces which vary with speed

Proposition.

$$\mathbf{a} = \mathbf{v} \frac{dv}{dx}$$

Proof.

$$\mathbf{a} = \frac{d\mathbf{x}}{dt} \times \frac{d\mathbf{v}}{dx} = \mathbf{v} \frac{dv}{dx}$$

□

12 Elastic strings and springs

12.1 Hooke's Law

Law (Hooke's Law). There are two cases for using Hooke's Law

- (i) Elastic strings: The tension T in an elastic string is


$$T = \frac{\lambda x}{l}$$

where

l is the natural (unstretched) length of the string,

x is the extension and

λ is the modulus of elasticity



When the string is slack there is no tension.

- (ii) Elastic springs: The tension, or thrust, T in an elastic spring is



$$T = \frac{\lambda x}{l}$$

where

l is the natural (unstretched) length of the string,

x is the extension or compression and

λ is the modulus of elasticity



12.2 Energy stored in an elastic string or spring

Like kinematics, If there is force F and displacement traveled δs , the Work done is $\delta W = F\delta s$. Similarly, If the tension force is T and string/spring extended/stretched, then

$$\delta W \approx T\delta x$$

Total work done in extending from $x = 0$ to $x = X$ is approximately

$$\sum_0^X T\delta x$$

and, as $\delta x \rightarrow 0$, the total work done:

$$W = \int_0^X T dx = \int_0^X \frac{\lambda x}{l} dx = \frac{\lambda x^2}{2l}$$

The expression of Total work done is also called the Elastic Potential Energy

13 Further dynamics

13.1 Impulse of a variable force

$$\delta I \approx F(t)\delta t$$

The total impulse from time t_1 to t_2 is

$$I \approx \sum_{t_1}^{t_2} F(t)\delta t$$

and as $\delta t \rightarrow 0$, the total impulse is

$$I = \int_{t_1}^{t_2} F(t)dt$$

Also, as $F(t) = ma = m \frac{dv}{dt}$

$$\int_{t_1}^{t_2} F(t)dt = \int_U^V m dv = mV - mU$$

13.2 Work done by a variable force

$$\delta W \approx G(x)\delta x$$

and the total work done in moving from a displacement x_1 to x_2 is

$$W \approx \sum_{x_1}^{x_2} G(x)\delta x$$

and as $\delta x \rightarrow 0$, the total work done is

$$W = \int_{x_1}^{x_2} G(x)dx$$

Also $G(x) = ma = m \frac{dv}{dx} = m \frac{dx}{dt} \times \frac{dv}{dx} = mv \frac{dv}{dx}$

$$\int_{x_1}^{x_2} G(x)dx = \int_U^V mv dv = \frac{1}{2}mV^2 - \frac{1}{2}mU^2$$

13.3 Newton's Law of Gravitation

Law. The force of attraction between two bodies of masses M_1 and M_2 is directly proportional to the product of their masses and inversely proportional to the square of the distance, d , between them:

$$F = \frac{GM_1M_2}{d^2}$$

where G is a constant known as the constant of Gravitation

13.4 Finding k in $F = \frac{k}{x^2}$

$$F = ma = \frac{k}{d^2}$$

13.5 Simple harmonic motion S.H.M.

14 Motion in a circle

14.1 Angular velocity

14.2 Acceleration

Types of problems

14.3 Motion in a vertical circle

Types of problems

15 Statics of rigid bodies 2

15.1 Centre of mass

16 Relative motion

17 Elastic collisions in two dimensions

18 Resisted motion of a particle moving in a straight line

19 Damped and forced harmonic motion

20 Stability

21 Applications of vectors in mechanics

22 Variable mass

23 Moments of inertia of a rigid body

24 Rotation of a rigid body about a fixed smooth axis