

Mathematical Preliminaries

- Sets
- Graphs
- Proof Techniques

SETS

A set is a collection of elements

$$A = \{1, 2, 3\}$$

$$B = \{train, bus, bicycle, airplane\}$$

We write

$$1 \in A$$

$$ship \notin B$$

Set Representations


$$C = \{ a, b, c, d, e, f, g, h, i, j, k \}$$

$$C = \{ a, b, \dots, k \} \longrightarrow \textit{finite set}$$

$$S = \{ 2, 4, 6, \dots \} \longrightarrow \textit{infinite set}$$

$$S = \{ j : j > 0, \text{ and } j = 2k \text{ for some } k > 0 \}$$

$$A = \{1, 2, 3, 4, 5\}$$



Universal Set: all possible elements

$$U = \{1, \dots, 10\}$$


Set Operations

$$A = \{1, 2, 3\}$$

$$B = \{2, 3, 4, 5\}$$


- Union

$$A \cup B = \{1, 2, 3, 4, 5\}$$



- Intersection


$$A \cap B = \{2, 3\}$$



- Difference

$$A - B = \{1\}$$

$$B - A = \{4, 5\}$$




Venn diagrams

- Complement

Universal set = $\{1, \dots, 7\}$


$$A = \{1, 2, 3\} \longrightarrow \overline{A} = \{4, 5, 6, 7\}$$



$$\overline{\overline{A}} = A$$

$$\{ \text{even integers} \} = \{ \text{odd integers} \}$$

Integers



DeMorgan's Laws

$$\overline{A \cup B} = \bar{A} \cap \bar{B}$$

$$\overline{A \cap B} = \bar{A} \cup \bar{B}$$

Empty, Null Set: \emptyset

$$\emptyset = \{\}$$

$$S \cup \emptyset = S$$

$$S \cap \emptyset = \emptyset$$

$$S - \emptyset = S$$

$$\emptyset - S = \emptyset$$

$$\overline{\emptyset} = \text{Universal Set}$$


Subset

$$A = \{1, 2, 3\}$$

$$B = \{1, 2, 3, 4, 5\}$$

$$A \subseteq B$$

Proper Subset: $A \subset B$




Disjoint Sets

$$A = \{ 1, 2, 3 \}$$

$$B = \{ 5, 6 \}$$

$$A \cap B = \emptyset$$



Set Cardinality

- For finite sets

$$A = \{ 2, 5, 7 \}$$

$$|A| = 3$$

(set size)

Powersets

A powerset is a set of sets

$$S = \{ a, b, c \}$$

Powerset of S = the set of all the subsets of S

$$2^S = \{ \emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\} \}$$

Observation: $|2^S| = 2^{|S|} \quad (8 = 2^3)$

Cartesian Product

$$A = \{ 2, 4 \}$$

$$B = \{ 2, 3, 5 \}$$

$$A \times B = \{ (2, 2), (2, 3), (2, 5), (4, 2), (4, 3), (4, 5) \}$$


$$|A \times B| = |A| |B|$$

Generalizes to more than two sets

$$A \times B \times \dots \times Z$$

GRAPHS

A directed graph




- Nodes (Vertices)

$$V = \{ a, b, c, d, e \}$$


- Edges

$$E = \{ (a,b), (b,c), (b,e), (c,a), (c,e), (d,c), (e,b), (e,d) \}$$

Labeled Graph




Walk



Walk is a sequence of adjacent edges

$(e, d), (d, c), (c, a)$


Path



Path is a walk where no edge is repeated

Simple path: no node is repeated


Cycle



Cycle: a walk from a node (base) to itself


Simple cycle: only the base node is repeated

Euler Tour




A cycle that contains each edge once

Hamiltonian Cycle




A simple cycle that contains all nodes


Trees



Trees have no cycles



Binary Trees



PROOF TECHNIQUES

- Proof by induction
- Proof by contradiction

Induction

We have statements P_1, P_2, P_3, \dots

If we know

- for some b that P_1, P_2, \dots, P_b are true
- for any $k \geq b$ that

$$P_1, P_2, \dots, P_k \text{ imply } P_{k+1}$$

Then

Every P_i is true

Proof by Induction

- Inductive basis

Find P_1, P_2, \dots, P_b which are true

- Inductive hypothesis

Let's assume P_1, P_2, \dots, P_k are true,
for any $k \geq b$

- Inductive step


Show that P_{k+1} is true

Example

Theorem: A binary tree of height n
has at most 2^n leaves.

Proof by induction:

let $L(i)$ be the maximum number of
leaves of any subtree at height i



We want to show: $L(i) \leq 2^i$

- Inductive basis

$$L(0) = 1 \quad (\text{the root node}) \quad \bigcirc$$


- Inductive hypothesis

Let's assume $L(i) \leq 2^i$ for all $i = 0, 1, \dots, k$

- Induction step


we need to show that $L(k + 1) \leq 2^{k+1}$

Induction Step



From Inductive hypothesis: $L(k) \leq 2^k$

Induction Step



$$L(k+1) \leq 2 * L(k) \leq 2 * 2^k = 2^{k+1}$$

(we add at most two nodes for every leaf of level k)

Proof by Contradiction

We want to prove that a statement P is true

- we assume that P is false
- then we arrive at an incorrect conclusion
- therefore, statement P must be true

Example

Theorem: $\sqrt{2}$ is not rational

Proof:

Assume by contradiction that it is rational

$$\sqrt{2} = n/m$$

n and m have no common factors

We will show that this is impossible

$$\sqrt{2} = n/m \quad \longrightarrow \quad 2 m^2 = n^2$$

Therefore, n^2 is even \longrightarrow n is even
 $n = 2 k$

$$2 m^2 = 4 k^2 \quad \longrightarrow \quad m^2 = 2 k^2 \quad \longrightarrow \quad m \text{ is even} \\ m = 2 p$$

Thus, m and n have common factor 2

Contradiction!

Languages

A language is a set of strings

String: A sequence of letters

Examples: "cat", "dog", "house", ...

Defined over an alphabet:

$$\Sigma = \{a, b, c, \dots, z\}$$

Alphabets and Strings

We will use small alphabets: $\Sigma = \{a, b\}$

Strings

a

ab

abba

baba

aaabbbbaabab

$u = ab$

$v = bbbaaa$

$w = abba$

String Operations

$$w = a_1a_2 \cdots a_n$$

abba

$$v = b_1b_2 \cdots b_m$$

bbbbaaa

Concatenation

$$wv = a_1a_2 \cdots a_nb_1b_2 \cdots b_m$$

abbabbbbaaa

$$w = a_1 a_2 \cdots a_n$$

ababaaaabbb

Reverse

$$w^R = a_n \cdots a_2 a_1$$

bbbaaababab

String Length

$$w = a_1 a_2 \cdots a_n$$

Length: $|w| = n$

Examples: $|abba| = 4$

$$|aa| = 2$$

$$|a| = 1$$

Length of Concatenation

$$|uv| = |u| + |v|$$

Example: $u = aab, |u| = 3$

$v = abaab, |v| = 5$

$$|uv| = |aababaab| = 8$$

$$|uv| = |u| + |v| = 3 + 5 = 8$$

Empty String

A string with no letters: λ

Observations: $|\lambda| = 0$

$$\lambda w = w \lambda = w$$

$$\lambda abba = abba \lambda = abba$$

Substring

Substring of string:

a subsequence of consecutive characters

String

abbab

abbab

abbab

abbab

Substring

ab

abba

b

bbab

Prefix and Suffix

abbab

Prefixes

Suffixes

λ

abbab

a

bbab

ab

bab

abb


ab

abba

b

abbab

λ



Another Operation

$$w^n = \underbrace{ww \cdots w}_n$$

Example: $(abba)^2 = abbaabba$

Definition: $w^0 = \lambda$

$$(abba)^0 = \lambda$$

The * Operation

Σ^* : the set of all possible strings from
alphabet Σ

$$\Sigma = \{a, b\}$$

$$\Sigma^* = \{\lambda, a, b, aa, ab, ba, bb, aaa, aab, \dots\}$$

The + Operation

Σ^+ : the set of all possible strings from alphabet Σ except λ

$$\Sigma = \{a, b\}$$

$$\Sigma^* = \{\lambda, a, b, aa, ab, ba, bb, aaa, aab, \dots\}$$

$$\Sigma^+ = \Sigma^* - \lambda$$

$$\Sigma^+ = \{a, b, aa, ab, ba, bb, aaa, aab, \dots\}$$

Languages

A language is any subset of Σ^*

Example: $\Sigma = \{a, b\}$

$$\Sigma^* = \{\lambda, a, b, aa, ab, ba, bb, aaa, \dots\}$$

Languages: $\{\lambda\}$

$$\{a, aa, aab\}$$

$$\{\lambda, abba, baba, aa, ab, aaaaaa\}$$

Note that:

Sets

$$\emptyset = \{ \} \neq \{ \lambda \}$$

Set size

$$|\{ \}| = |\emptyset| = 0$$

Set size

$$|\{ \lambda \}| = 1$$

String length

$$|\lambda| = 0$$

Another Example

An infinite language $L = \{a^n b^n : n \geq 0\}$

λ
 ab
 $aabb$
 $aaaaabbbbb$

} $\in L$ $abb \notin L$

Operations on Languages

The usual set operations

$$\{a, ab, aaaa\} \cup \{bb, ab\} = \{a, ab, bb, aaaa\}$$

$$\{a, ab, aaaa\} \cap \{bb, ab\} = \{ab\}$$

$$\{a, ab, aaaa\} - \{bb, ab\} = \{a, aaaa\}$$

Complement: $\bar{L} = \Sigma^* - L$

$$\overline{\{a, ba\}} = \{\lambda, b, aa, ab, bb, aaaa, \dots\}$$

Reverse

Definition: $L^R = \{w^R : w \in L\}$

Examples: $\{ab, aab, baba\}^R = \{ba, baa, abab\}$

$$L = \{a^n b^n : n \geq 0\}$$

$$L^R = \{b^n a^n : n \geq 0\}$$

Concatenation

Definition: $L_1L_2 = \{xy : x \in L_1, y \in L_2\}$

Example: $\{a, ab, ba\}\{b, aa\}$

$$= \{ab, aaa, abb, abaa, bab, baaa\}$$

Another Operation

Definition: $L^n = \underbrace{LL \cdots L}_n$

$$\{a,b\}^3 = \{a,b\}\{a,b\}\{a,b\} = \\ \{aaa, aab, aba, abb, baa, bab, bba, bbb\}$$

Special case: $L^0 = \{\lambda\}$

$$\{a, bba, aaa\}^0 = \{\lambda\}$$

More Examples

$$L = \{a^n b^n : n \geq 0\}$$

$$L^2 = \{a^n b^n a^m b^m : n, m \geq 0\}$$

$$aabbbaaabb \in L^2$$

Star-Closure (Kleene *)

Definition: $L^* = L^0 \cup L^1 \cup L^2 \dots$

Example:

$$\{a, bb\}^* = \left\{ \begin{array}{l} \lambda, \\ a, bb, \\ aa, abb, bba, bbbb, \\ aaa, aabb, abba, abbbb, \dots \end{array} \right\}$$

Positive Closure

Definition: $L^+ = L^1 \cup L^2 \cup \dots$
 $= L^* - \{\lambda\}$

$$\{a, bb\}^+ = \left\{ \begin{array}{l} a, bb, \\ aa, abb, bba, bbbb, \\ aaa, aabb, abba, abbbb, \dots \end{array} \right\}$$