


Standard Representations of Regular Languages



When we say: We are given
a Regular Language L

We mean: Language L is in a standard
representation

Elementary Questions

about


Regular Languages

Membership Question

Question: Given regular language L
and string w
how can we check if $w \in L$?


Answer: Take the DFA that accepts L
and check if w is accepted

DFA



$$w \in L$$

DFA




$$w \notin L$$

Question: Given regular language L
how can we check
if L is empty: $(L = \emptyset)$?

Answer: Take the DFA that accepts L


Check if there is any path from
the initial state to a final state

DFA



$$L \neq \emptyset$$

DFA




$$L = \emptyset$$

Question: Given regular language L
how can we check
if L is finite?

Answer: Take the DFA that accepts L


Check if there is a walk with cycle
from the initial state to a final state

DFA



L is infinite

DFA



L is finite


Question: Given regular languages L_1 and L_2
how can we check if $L_1 = L_2$?

Answer: Find if $(L_1 \cap \overline{L_2}) \cup (\overline{L_1} \cap L_2) = \emptyset$


$$(L_1 \cap \overline{L_2}) \cup (\overline{L_1} \cap L_2) = \emptyset$$



$$L_1 \cap \overline{L_2} = \emptyset \quad \text{and} \quad \overline{L_1} \cap L_2 = \emptyset$$



$$L_1 \subseteq L_2$$



$$L_2 \subseteq L_1$$



$$L_1 = L_2$$


$$(L_1 \cap \overline{L_2}) \cup (\overline{L_1} \cap L_2) \neq \emptyset$$




$$L_1 \cap \overline{L_2} \neq \emptyset$$

or

$$\overline{L_1} \cap L_2 \neq \emptyset$$



$$L_1 \not\subseteq L_2$$



$$L_2 \not\subseteq L_1$$



$$L_1 \neq L_2$$

Non-regular languages

Non-regular languages

$$\{a^n b^n : n \geq 0\}$$

$$\{vv^R : v \in \{a,b\}^*\}$$

Regular languages

$$a^*b$$

$$b^*c + a$$

$$b + c(a + b)^*$$


etc...

How can we prove that a language L is not regular?

Prove that there is no DFA that accepts L





Problem: this is not easy to prove

Solution: the Pumping Lemma !!!






The Pigeonhole Principle




4 pigeons



3 pigeonholes



A pigeonhole must
contain at least two pigeons



n pigeons





.....




m pigeonholes

$n > m$



.....





The Pigeonhole Principle

n pigeons


m pigeonholes

$$n > m$$

There is a pigeonhole
with at least 2 pigeons



.....




The Pigeonhole Principle

and

DFAs

DFA with 4 states



In walks of strings:


a

no state

aa

is repeated

aab




In walks of strings: $aabb$

$bbaa$

$abbabb$

$abbbabbabb...$


a state
is repeated



If string w has length $|w| \geq 4$:

Then the transitions of string w
are more than the states of the DFA

Thus, a state must be repeated




In general, for any DFA:

String w has length \geq number of states



A state q must be repeated in the walk of w





In other words for a string w :

\xrightarrow{a} transitions are pigeons




(q) states are pigeonholes



The Pumping Lemma

Take an **infinite** regular language L

There exists a DFA that accepts L



m
states

Take string w with $w \in L$


There is a walk with label w :




If string w has length $|w| \geq m$ (number of states of DFA)

then, from the pigeonhole principle:


a state is repeated in the walk w



Let q be the first state repeated in the walk of w




Write $w = x y z$




Observations: length $|x y| \leq m$ number
 of states
 of DFA


length $|y| \geq 1$




Observation: The string xzy is accepted



Observation: The string $x y y z$ is accepted




Observation: The string $x y y y z$ is accepted




In General:

The string $x y^i z$
is accepted $i = 0, 1, 2, \dots$






In General: $x y^i z \in L \quad i = 0, 1, 2, \dots$




Language accepted by the DFA



In other words, we described:



The Pumping Lemma !!!



The Pumping Lemma:

- Given a infinite regular language L
- there exists an integer m
- for any string $w \in L$ with length $|w| \geq m$
- we can write $w = x y z$
- with $|x y| \leq m$ and $|y| \geq 1$
- such that: $x y^i z \in L \quad i = 0, 1, 2, \dots$

Applications of the Pumping Lemma

Theorem: The language $L = \{a^n b^n : n \geq 0\}$
is not regular

Proof: Use the Pumping Lemma

$$L = \{a^n b^n : n \geq 0\}$$

Assume for contradiction
that L is a regular language

Since L is infinite
we can apply the Pumping Lemma

$$L = \{a^n b^n : n \geq 0\}$$

Let m be the integer in the Pumping Lemma

Pick a string w such that: $w \in L$

$$\text{length } |w| \geq m$$

We pick $w = a^m b^m$

Write: $a^m b^m = x y z$

From the Pumping Lemma

it must be that length $|x y| \leq m, \quad |y| \geq 1$

$$xyz = a^m b^m = \overbrace{a \dots a}^m \overbrace{a \dots a b \dots b}^m$$

$x \quad y \quad z$

Thus: $y = a^k, \quad k \geq 1$

$$x y z = a^m b^m$$

$$y = a^k, \quad k \geq 1$$

From the Pumping Lemma: $x y^i z \in L$

$$i = 0, 1, 2, \dots$$

Thus: $x y^2 z \in L$

$$x y z = a^m b^m$$

$$y = a^k, \quad k \geq 1$$

From the Pumping Lemma: $x y^2 z \in L$


$$xy^2z = \overbrace{a \dots a a \dots a a \dots a a \dots a}^{m+k} \overbrace{b \dots b}^m \in L$$

$\underbrace{\hspace{1.5cm}}_x \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{2.5cm}}_z$

Thus: $a^{m+k} b^m \in L$

$$a^{m+k}b^m \in L \qquad k \geq 1$$

BUT: $L = \{a^n b^n : n \geq 0\}$



$$a^{m+k}b^m \notin L$$

CONTRADICTION!!!

Therefore: Our assumption that L
is a regular language is not true

Conclusion: L is not a regular language

Non-regular languages $\{a^n b^n : n \geq 0\}$



Regular languages