

→ **[plot_format,gnuplot]****(%i1) solve(2·x^3-7·x^2-7·x+30=0,x);****(%o1)** $\left[x = -2, x = -\frac{5}{2}, x = 3 \right]$ **(%i3) allroots(2·x^3-7·x^2-7·x+30=0);****(%o3)** $[x = 2.5000000000000001, x = -2.0, x = 3.0]$ **(%i5) find_root(sin(x)-x-2,x,-4,0);****(%o5)** -2.554195952837043→ **load(newton)\$****(%i14) newton(x-2-log(x),x,0.1,1E-8);****(%o14)** 0.1585943395480817**(%i15) newton(x-2-log(x),x,3,1E-8);****(%o15)** 3.146193220620587**(%i20) plot2d(x-2-log(x),[x,0,3]);**


log: encountered log(0).

plot2d: expression evaluates to non-numeric value somewhere in plotting range.

(%o20) false**(%i24) allroots(x^5-x^4+2·x^3+x^2-x+5);****(%o24)** $[x = 1.074792179417892 \text{ %i} + 0.9706128683007163, x = 0.9706128683007163 - 1.074792179417892 \text{ %i}, x = -1.182801347552513, x = 1.41457071898534 \text{ %i} + 0.1207878054755404, x = 0.1207878054755404 - 1.41457071898534 \text{ %i}]$

Chercher la racine de

(%i30) f(x):=if x < 2 then x/(x-3) else (x-2)^2 -2;**(%o30)** $f(x) := \text{if } x < 2 \text{ then } \frac{x}{x-3} \text{ else } (x-2)^2 - 2$ **(%i36) solve(f(x)=0,x);****(%o36)** $\left[\left(\text{if } x < 2 \text{ then } \frac{x}{x-3} \text{ else } x^2 - 4x + 2 \right) = 0 \right]$ **(%i37) allroots(2·x^3-7·x^2-7·x+30=0);****(%o37)** $[x = 2.5000000000000001, x = -2.0, x = 3.0]$ **(%i44) find_root(f(x),x,-4,0);****(%o44)** 0.0**(%i49) wxplot2d(f(x),[x,0,3]);****(%t49)**



(%o49)