




Contents lists available at ScienceDirect

Contemporary Clinical Trials

journal homepage: www.elsevier.com/locate/conclintrial

A decade of individual participant data meta-analyses: A review of current practice

Mark Simmonds^{a,*}, Gavin Stewart^b, Lesley Stewart^a^a Centre for Reviews and Dissemination, University of York, UK^b School of Agriculture, Food and Rural Development, Newcastle University, UK

ARTICLE INFO

Article history:

Received 2 April 2015

Received in revised form 5 June 2015

Accepted 15 June 2015

Available online xxxx

Keywords:

Individual participant data

Meta-analysis

Systematic review

ABSTRACT

Introduction: Individual participant data (IPD) systematic reviews and meta-analyses are often considered to be the gold standard for meta-analysis. In the ten years since the first review into the methodology and reporting practice of IPD reviews was published much has changed in the field. This paper investigates current reporting and statistical practice in IPD systematic reviews.

Methods: A systematic review was performed to identify systematic reviews that collected and analysed IPD. Data were extracted from each included publication on a variety of issues related to the reporting of IPD review process, and the statistical methods used.

Results: There has been considerable growth in the use of “one-stage” methods to perform IPD meta-analyses. The majority of reviews consider at least one covariate other than the primary intervention, either using subgroup analysis or including covariates in one-stage regression models. Random-effects analyses, however, are not often used.

Reporting of review methods was often limited, with few reviews presenting a risk-of-bias assessment. Details on issues specific to the use of IPD were little reported, including how IPD were obtained; how data was managed and checked for consistency and errors; and for how many studies and participants IPD were sought and obtained.

Conclusion: While the last ten years have seen substantial changes in how IPD meta-analyses are performed there remains considerable scope for improving the quality of reporting for both the process of IPD systematic reviews, and the statistical methods employed in them. It is to be hoped that the publication of the PRISMA-IPD guidelines specific to IPD reviews will improve reporting in this area.

© 2015 Elsevier Inc. All rights reserved.

Contents

1. Introduction	0
2. Methods	0
3. Results	0
4. Quality of reporting	0
5. Statistical methods	0
6. Discussion	0
7. Conclusion	0
Funding	0
Acknowledgements	0
References	0

1. Introduction

The aim of individual participant data (IPD) systematic reviews and meta-analyses is to obtain all the original, raw participant data from all studies on a specified topic, in order to reanalyse the data and pool it across studies. While obtaining all the original data from all relevant

* Corresponding author.

E-mail address: mark.simmonds@york.ac.uk (M. Simmonds).

studies may be time consuming and difficult, IPD meta-analysis is recognised as having many advantages over meta-analysis based on data reported in publications, and is considered the “gold standard” for meta-analysis [1–3].

Ten years ago the first review of the practice and reporting of individual participant data meta-analyses was published in *Clinical Trials* [4]. At that time IPD meta-analysis was still in its early stages with numbers of publications being limited before the late 1990s. The development of statistical methods in the area was also in its infancy, with key methods papers having been published only a few years before [5–7]. The review found that reporting of the processes of IPD meta-analyses was generally poor, particularly with regard to how much of the total IPD was obtained for analysis, and with poor reporting of statistical methods. Statistical analysis was also limited, with little investigation of heterogeneity, and most meta-analyses focussing on calculating overall treatment effects, with little consideration given to subgroup analyses, or how factors such as a participant's age might modify the effectiveness of a treatment. Another review performed in 2008 and published in 2010 [8] came to broadly similar conclusions.

In the ten years since that first review was published much has changed. The number of systematic reviews and meta-analyses published continues to grow, and consequently IPD meta-analysis has also grown in popularity. Advances in computing have also made meta-analyses easier to perform; most methods for IPD meta-analysis can now be implemented in all major statistical software packages. As the number of IPD meta-analyses grows there is an increased need to ensure high quality of conduct reporting of these analyses, to avoid some of the problems identified in the original review.

The Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) statement [9] is widely used as guidance on how to report a systematic review and meta-analysis. Recently a new PRISMA-IPD statement has been released specifically to guide the conduct and reporting of IPD reviews and meta-analyses [10]. As part of the process of creating this statement, a review of current practice in reporting of IPD reviews was conducted to identify areas where reporting was poor to inform development of the guidance. This paper presents the results of this review, updated to 2014, and expanded to consider statistical methods used in IPD meta-analyses.

2. Methods

The aim of this review was to identify published systematic reviews of medical interventions that sought to obtain and meta-analyse IPD. It was intended to obtain a representative sample of recent reviews for analysis, rather than find all such reviews, so an exhaustive database search was not performed. A MEDLINE search was performed including terms “individual participant/patient data”, “meta-analysis” and “systematic review”. This search was originally performed in January 2013 and was updated in February 2015. Papers published from 2008 (when the previous review in this field by Riley et al. was performed (8)) up to the end of 2014 were sought.

Only systematic reviews of IPD were included, so we did not include “opportunistic” analyses (where the reviewers combined data to which they had access) or collaborative reviews (where data from several collaborators was combined without performing a formal systematic review). Other reasons for exclusion were: papers that did not consider a medical treatment (e.g. epidemiologic reviews of the causes of a disease, diagnostic test accuracy reviews and reviews of disease prognosis), and papers not published in English. One reviewer reviewed titles and abstracts. For those considered potentially eligible the full text was sought. Where full text could be obtained the paper was further checked for eligibility.

For all eligible papers one reviewer extracted a range of data covering the reporting of the review (such as reporting of medical field, numbers of included studies, whether quality assessment was performed); the statistical methods used (such as outcomes considered, whether

one or two stage methods were used, descriptions of methods, and assessment of heterogeneity); reporting of results (how many outcomes reported, how many were statistically significant, reporting of subgroup analyses or meta-regression, use of forest plots). The data extracted from the publications was analysed by creating suitable summary tables and graphs.

3. Results

The original and updated search together identified 1371 potentially relevant records. After checking titles and abstracts 184 papers were considered for inclusion. After obtaining full texts (where readily available) and further checking, 100 systematic reviews of IPD were included in this review [11–110]. Because a specific rather than a sensitive search was used, and because we include only systematic reviews, these represent only a minority of all IPD meta-analyses performed. For comparison, in 2014 there were 68 references that matched “Individual participant/patient data meta-analysis” in MEDLINE, not all of which will be IPD meta-analyses. These remain a small minority of meta-analyses as a whole; 6020 papers were tagged as meta-analyses in MEDLINE in 2014.

In the sample considered here, the numbers of reviews per year varied from 10 to 22, but there was no evidence of a trend over time, suggesting that the use of IPD analysis has stabilised in recent years. Fig. 1 shows the medical field in which the reviews were conducted. As in the review published in 2005, cancer and cardiovascular disease predominate. IPD methods appear to be gaining popularity in other fields such as paediatrics and mental health, but numbers remain small. Fig. 2a shows the number of studies for which IPD was obtained, and Fig. 2b the total number of participants providing IPD, across the reviews. Sizes vary considerably from two studies to 78 (median 8), and from 64 to 70,528 participants (median 1879).

4. Quality of reporting

The included reviews varied considerably in how they reported aspects of the review process, such as the search strategy, and quality assessment process. The numbers of reviews reporting key aspects of the IPD review process are summarised in Table 1. In general, aspects of the review process common to all systematic reviews were well reported, with, for example, 80% of reviews reporting details of the search strategy and search process. Inclusion and exclusion criteria were also generally clearly described. One area that was not widely reported was risk-of-bias or study quality assessments, despite this being a component included in the PRISMA statement. Only 34% of reviews reported performing any risk-of-bias assessment, and only 65% (22 of 34) of those reported the findings of the assessment in any detail.

Aspects of reporting specific to IPD reviews were generally rather poorly reported. Only 48% of reviews explained why an IPD review was performed, rather than a review based on published data. Only 52% described how the IPD was obtained, and for most that did it was described simply as “by contacting the study authors” or with a similar phrase. Only 33% of reviews reported that any checking of the IPD was performed to ensure consistency or identify data errors. Just 23% of reviews described why IPD was unavailable where they could not obtain IPD from all relevant studies. The most common reasons given were inability to contact authors, non-cooperation of study authors, and loss of original data.

One aspect of IPD reviews identified as being particularly poorly reported in the 2005 paper was the numbers of studies and participants from which IPD was sought, but not necessarily obtained. As in 2005 this still appears to be poorly reported: 24% of reviews did not report how many studies were sought, and 56% did not report the total number of participants sought. Of those that did report these data, most reported that they obtained all the studies (45% of such reviews) and/or all participants (58%).




Fig. 1. Distribution across medical fields in the IPD reviews.

5. Statistical methods

The review of 2005 identified two distinct statistical approaches for IPD meta-analysis [4]. In a two-stage approach studies are analysed individually to obtain summary results for each study in the first stage, and the summary results pooled across studies as in a conventional meta-analysis in the second stage. In a one-stage approach all data from all studies are analysed simultaneously, in a single model, usually a regression model. This distinction in approaches remains in current practice.

Table 2 shows the distribution of methods used according to the type of primary outcome in each review. In the 2005 review two-stage methods predominated, being the method used in over two-thirds of reviews. The situation has changed dramatically, with just 26% of reviews using only two-stage methods. One-stage methods are now more popular, being used exclusively in 37% of reviews. This is particularly true of reviews of dichotomous or continuous outcomes, where 51% of reviews used only one-stage methods. The most commonly used one-stage approaches are the use of logistic and linear regression models to pool data across studies. This change is probably explained

by growing familiarity with one-stage methods among reviewers, and improved software to perform them. Two-stage methods remain more popular in analyses of survival data; 76% of survival analyses used two-stage methods. This may be because statistical methods for one-stage analyses of survival data are less well developed and little software is available. Two-stage methods are particularly common in reviews of cancer treatment; 79% of such reviews used two stage methods. In IPD reviews of cancer trials two-stage methods based on the log rank test are well-established and remain the most commonly used statistical method [111].

Description of statistical methods was often poor, with 36% of papers deemed not to have described the statistical methods used (see Table 3); six were described so poorly it was not possible to tell what type of outcome was used or whether analyses were one or two-stage. One-stage methods were in general more poorly described, perhaps because of the greater complexity involved in describing properties of regression models. Even where the methods were judged to have been described this description may have been limited. For example, one-stage regression models were often described simply as “a logistic regression model was used...”, with no further clarification as to the




Fig. 2. Distribution of numbers of studies and participants.

Table 1
Reporting quality in the IPD reviews.

	Number reporting		Number reporting
Rationale for using IPD	48	IPD checking process	33
Details of search	80	Results of IPD checking	8
How IPD was obtained	52	Whether missing or incomplete IPD was sought	11
Risk-of-bias or quality assessment	34	Reasons for non-availability of IPD	23
Results of quality assessment	22		

structure of the model. Only in a minority of cases did we consider that a reader could readily replicate the one-stage analyses performed in the reviews. The use of the actual terms “one-stage” and “two-stage” to describe IPD methods appears to be growing in popularity, but these terms were used in only a minority of reviews.

Table 4 summarises the use of fixed and random effects and reporting of heterogeneity. Only 33% of all reviews used random-effects analyses. This is a greater proportion than in 2005, but small compared to the widespread use of random-effects methods in published data meta-analyses. Most analyses of survival data used fixed effect models (84%), because this remains conventional in analyses of cancer survival. The choice between fixed and random effects was more evenly balanced for other outcomes, across one-and two-stage methods and in medical areas other than cancer. Assessments of heterogeneity were rarely reported. Only 18% of studies reported I^2 , and 24% Cochran's Q test. Heterogeneity estimates were rarely reported in papers using one-stage regression models; just four reviews using one-stage models reported the heterogeneity in these models.

IPD meta-analysis, because it provides access to full data on every participant, is often used to go beyond estimating the overall treatment effect to investigate how covariates such as participant age might modify the effect. A summary of how covariates were considered across reviews is given in Table 5. Unlike in 2005, the majority of reviews (72%) examined how covariates might influence the effect of the intervention. In two-stage analyses subgroup analysis was most commonly used, either by dividing trials into groups or dividing participants within trials into subgroups. Only three reviews used meta-regression. One review estimated interactions between treatment and covariates within each study and pooled these results across studies. In one-stage analyses, most papers reported including covariates in the one-stage regression model (21 reviews), although exactly how this was done was rarely reported.

As IPD may not be obtained for all studies combining IPD with published results may be important. However only 14 of the 66 reviews that were missing at least some IPD made any attempt to combine IPD and published data. Nearly all of these used a two-stage approach (11 reviews), pooling summary estimates from publications with summary results derived from the IPD.

When reporting results and outcomes, most papers (77%) reported more than one outcome, and of those 68% of reviews reported at least one result that was not statistically significant. Similarly, when investigating the impact covariates such as age had on the treatment effect, most papers (56%) reported results of multiple subgroup analyses, or findings from multiple covariates in one-stage models. The subgroup analyses or interaction terms were rarely statistically significant; only

25 reviews identified any statistically significant covariate effects. This suggests that there is little selective reporting bias in IPD reviews. Most reviews presented some results in forest plots, with 75% of reviews using a two-stage analysis presenting a forest plot for at least one outcome, although most reviews presented forest plots for only a subset of all the outcomes reported. Similarly 64% of reviews that used a two-stage subgroup analysis to assess the impact of covariates presented a forest plot of the subgroup analysis for at least one covariate.

6. Discussion

Much has changed in IPD systematic reviews and meta-analyses in the last ten years. The most notable change is the switch from a predominance of two-stage methods, where studies are analysed separately and summary results pooled across studies; to one-stage methods where all IPD are analysed simultaneously. This is probably driven by growing familiarity with these methods, improved software, and recognition that regression models offer the greatest flexibility for IPD analysis. Another key change is the growth in analyses of covariates other than treatment. The ability to examine how factors such as age can impact on treatment effectiveness has long been recognised as a benefit of IPD analysis, but up to 2005 it was little used. Now the majority of reviews consider at least one covariate other than treatment, either using subgroup analysis or including covariates in one-stage regression models.

In other areas there has been less change. IPD reviews remain a tiny minority of systematic reviews and meta-analyses, and most are still performed on cancer or cardiovascular disease trials. This might change if IPD become more readily available as a result of current initiatives to increase access to, and sharing of, clinical trial data. Poor quality of reporting, particularly of statistical methods, remains an issue, although there have been some improvements in this.

This review found that IPD systematic reviews were generally good at reporting the basic processes of a systematic review such as searching and inclusion criteria. One area of weakness was a lack of risk-of-bias assessment, despite this being explicitly required in the PRISMA guidance. Although the analysis of IPD can reduce some risks of bias, such as selective reporting (because data on all outcomes are collected from included trials not just those that were reported in the trial publication) or poor randomisation (by adjusting for covariates which were not balanced across groups), it cannot remove all potential biases, such as a lack of blinding. Hence considering risk of bias in an IPD review is important and should be performed in all such reviews. Much of a risk-of-bias

Table 2
Type of outcomes and main statistical approach used.

Statistical approach	Primary outcome				Total
	Dichotomous	Continuous	Survival	Other	
One-stage	14	13	10	0	37
Two-stage	6	5	15	0	26
One-stage and two-stage	9	3	19	0	31
Unclear	2	1	1	2	6
Total	31	22	45	2	

Table 3
Summary of quality of description of statistical methods.

Statistical approach	Quality of description of methods	Number of reviews
One-stage	Methods described	27
	Not described	10
Two-stage	Methods described	20
	Not described	6
One-stage and two-stage	Both described	17
	One-stage not described	8
	Two-stage not described	4
	Neither described	2
Unclear		6

Table 4
Summary of heterogeneity assessment across reviews.

	Statistical approach			Total
	One-stage	Two-stage	Both	
<i>Fixed vs random effects</i>				
Fixed effect	23	17	21	61
Random effects	14	9	10	33
<i>Reporting of heterogeneity</i>				
I ²	7	6	5	18
Cochran's Q (without I ²)	4	11	9	24
From one-stage model	3	–	1	4
Other	0	1	1	2
Not reported	23	7	15	45

assessment can be performed or supplemented by examining the IPD itself, but this has not been done in practice. The IPD reviews identified often did not report much detail on issues specific to the use of IPD, including explaining the benefits of using IPD; how IPD was obtained, managed and checked for consistency and errors; and how for how many studies and participants IPD were sought and obtained. These are all issues which should be discussed in any IPD review.

Reporting of statistical methods could also be improved in many reviews. Where two-stage methods were used they were mostly described adequately, as they generally used methods familiar from meta-analyses of published data. One-stage models, however, were generally poorly described, and simply noting that a logistic regression model, or similar, was used should not be considered sufficient. Ideally, all descriptions of one-stage models should clearly state the following: how models were stratified by study to preserve distinctions between studies and to preserve randomisation; whether models were adjusted for baseline factors, and if so, which ones; how treatment was modelled; whether any interactions between treatment and other covariates were included; whether random-effects were included and, if so, on which model terms. The presentation of a formal model specification, or sample computer code, may aid substantially in understanding how one-stage models are used.

Reporting of the results of one-stage analyses could also be improved. Reporting regression coefficients from the model, as was common in the included reviews, may be difficult to interpret. Using two-stage methods for part of the analysis, for example, in order to present a forest plot of the primary outcome with an assessment of heterogeneity, may aid understanding in IPD reviews. One-stage models could then be used for the more complex modelling processes, such as for investigating the impact of covariates on efficacy.

Assessing heterogeneity and performing random-effects meta-analyses have become commonplace in meta-analyses of published results, so it perhaps surprising that they are so little used in IPD analyses, with fixed effect analyses being in the majority, although some reviews may have considered random effects methods but not reported them because no heterogeneity was identified. Where IPD reviews intend to investigate how covariates might alter the effectiveness of an intervention, assessing heterogeneity to identify differences across studies and participants should be considered a priority.

Table 5
Summary of analyses of covariates.

	Number of reviews
Any analysis of covariates	72
Any two-stage analysis	50
Subgroups of studies	29
Subgroups of participants	36
Meta-regression	3
Within-study models including covariates	1
Covariates included in one-stage models	38

There remains considerable scope for the use of more advanced statistical modelling methods in IPD meta-analysis. Multivariate meta-analysis models, for example, are widely employed in other disciplines but appear to have no uptake in the world of clinical medicine. Comparing multiple models using stepwise regression, AIC or other model fit diagnostics is not widely used. Multiple imputation methods could assist in fitting models where there is substantial missing covariate or outcome data [112].

This review has focused on IPD systematic reviews of medical treatments. IPD meta-analysis is also popular for “opportunistic” and collaborative meta-analyses where researchers analyse IPD without performing a full systematic review. We suspect that many of the issues around reporting and use of statistical methods discussed in this paper will apply equally to these non-systematic analyses. Our database searches also found that there has been growth in the use of IPD meta-analysis in other medical areas, such as in reviews of diagnostic tests, network meta-analyses, epidemiological reviews examining the causes of disease, and reviews of disease prognosis and progression. In these last two cases one-stage regression models appear to be particularly widely used as they offer considerable flexibility when analysing non-randomised data such as data from cohort studies.

7. Conclusion

The last ten years have seen substantial changes in how IPD meta-analyses are performed, particularly with the growth of the use of one-stage regression models. There remains considerable scope for improving the quality of reporting both the process of IPD systematic reviews, and the statistical methods employed in them. It is to be hoped that the publication of the PRISMA-IPD guidelines specific to IPD reviews will improve reporting in this area [10].

Funding

The systematic review described in this paper was performed, in part, to inform the development of PRISMA-IPD. That project was supported, in part, through a National Institute for Health Research (NIHR) Senior Investigator personal award (Dr L. Stewart), and as part of the MRC–National Institute for Health Research Methodology Research Programme (grant ID 88053). The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation review, or approval of the manuscript; and decision to submit the manuscript for publication. The views expressed are those of the authors and not necessarily those of the NIHR or MRC.

Acknowledgements

We would like to thank Kath Wright for performing the literature searches for this review, and Dean Langan and Alison Smith for their assistance in the study selection and data extraction processes. We also thank the members of the PRISMA-IPD development group for their comments on early versions of this review.

References

- [1] L. Stewart, J.F. Tierney, To IPD or not to IPD? Eval. Health Prof. 25 (2002) 76–97.
- [2] L.A. Stewart, M.J. Clarke, Practical methodology of meta-analyses (overviews) using updated individual patient data, Stat. Med. 14 (1995) 2057–2079.
- [3] L.A. Stewart, M.K.B. Parmar, Meta-analysis of the literature or of individual patient data: is there a difference? Lancet 341 (1993) 418–422.
- [4] M.C. Simmonds, J.P.T. Higgins, L.A. Stewart, L.F. Tierney, M.J. Clarke, S.G. Thompson, Meta-analysis of individual patient data from randomized trials; a review of methods used in practice, Clin Trials 2 (3) (2005) 209–217.
- [5] J.P.T. Higgins, A. Whitehead, et al., Meta-analysis of continuous outcome data from individual patients, Stat. Med. 20 (2001) 2219–2241.
- [6] R.M. Turner, R.Z. Omar, M. Yang, H. Goldstein, S.G. Thompson, A multilevel model framework for meta-analysis of clinical trials with binary outcomes, Stat. Med. 19 (2000) 3417–3432.

- [7] A. Whitehead, R.Z. Omar, et al., Meta-analysis of ordinal outcomes using individual patient data, *Stat. Med.* 20 (2001) 2243–2260.
- [8] R.D. Riley, P.C. Lambert, G. Abo-Zaid, Meta-analysis of individual participant data: rationale, conduct, and reporting, *BMJ* 340 (2010).
- [9] D. Moher, A. Liberati, J. Tetzlaff, D.G. Altman, Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement, *BMJ* 339 (2009).
- [10] L.A. Stewart, M. Clarke, M. Rovers, et al., Preferred reporting items for a systematic review and meta-analysis of individual participant data: the PRISMA-IPD statement, *JAMA* 313 (16) (2015) 1657–1665.
- [11] Antithrombotic Trialists Collaboration, C. Baigent, L. Blackwell, R. Collins, J. Emberson, J. Godwin, et al., Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials, *Lancet* 373 (9678) (2009) 1849–1860.
- [12] A. Aupein, C. Le Pechoux, E. Rolland, W.J. Curran, K. Furuse, P. Fournel, et al., Meta-analysis of concomitant versus sequential radiochemotherapy in locally advanced non-small-cell lung cancer, *J. Clin. Oncol.* 28 (13) (2010) 2181–2190.
- [13] J.M. Bae, Y.Y. Choi, H. Kim, B.H. Oh, M.R. Roh, K. Nam, et al., Mohs micrographic surgery for extramammary Paget disease: a pooled analysis of individual patient data, *J. Am. Acad. Dermatol.* 68 (4) (2013) 632–637.
- [14] T.S. Baman, P. Meier, J. Romero, L. Gakenheimer, J.N. Kirkpatrick, P. Sovitch, et al., Safety of pacemaker reuse: a meta-analysis with implications for underserved nations, *Circ.* 4 (3) (2011) 318–323.
- [15] K.J.L. Bell, A. Hayen, P. Macaskill, J.C. Craig, B.C. Neal, K.M. Fox, et al., Monitoring initial response to angiotensin-converting enzyme inhibitor-based regimens: an individual patient data meta-analysis from randomized, placebo-controlled trials, *Hypertension* 56 (3) (2010) 533–539.
- [16] K. Bennett, K. Manassis, S.D. Walter, A. Cheung, P. Wilansky-Traynor, N. Diaz-Granados, et al., Cognitive behavioral therapy age effects in child and adolescent anxiety: an individual patient data meta-analysis, *Depress. Anxiety* 30 (9) (2013) 829–841.
- [17] Y. Ben-Shlomo, M. Spears, C. Boustred, M. May, S.G. Anderson, E.J. Benjamin, et al., Aortic pulse wave velocity improves cardiovascular event prediction: an individual participant meta-analysis of prospective observational data from 17,635 subjects, *J. Am. Coll. Cardiol.* 63 (7) (2014) 636–646.
- [18] D.A. Berry, N.T. Ueno, M.M. Johnson, X. Lei, J. Caputo, S. Rodenhuis, et al., High-dose chemotherapy with autologous stem-cell support as adjuvant therapy in breast cancer: overview of 15 randomized trials, *J. Clin. Oncol.* 29 (24) (2011) 3214–3223.
- [19] A. Beyerlein, M. Hadders-Algra, K. Kennedy, M. Fewtrell, A. Singhal, E. Rosenfeld, et al., Infant formula supplementation with long-chain polyunsaturated fatty acids has no effect on Bayley developmental scores at 18 months of age—IPD meta-analysis of 4 large clinical trials, *J. Pediatr. Gastroenterol. Nutr.* 50 (1) (2010) 79–84.
- [20] T. Bongartz, F.C. Warren, D. Mines, E.L. Matteson, K.R. Abrams, A.J. Sutton, Etanercept therapy in rheumatoid arthritis and the risk of malignancies: a systematic review and individual patient data meta-analysis of randomised controlled trials, *Ann. Rheum. Dis.* 68 (7) (2009) 1177–1183.
- [21] J. Bourhis, P. Blanchard, E. Maillard, D.M. Brizel, B. Movsas, J. Buentzel, et al., Effect of amifostine on survival among patients treated with radiotherapy: a meta-analysis of individual patient data, *J. Clin. Oncol.* 29 (18) (2011) 2590–2597.
- [22] S.S. Brar, J. ten Berg, R. Marcucci, M.J. Price, M. Valgimigli, H.-S. Kim, et al., Impact of platelet reactivity on clinical outcomes after percutaneous coronary intervention. A collaborative meta-analysis of individual participant data, *J. Am. Coll. Cardiol.* 58 (19) (2011) 1945–1954.
- [23] A.J. Breugnot, M. Swets, J.F. Bosset, L. Collette, A. Sainato, L. Cionini, et al., Adjuvant chemotherapy after preoperative (chemo)radiotherapy and surgery for patients with rectal cancer: a systematic review and meta-analysis of individual patient data, *Lancet Oncol.* 16 (2) (2015) 200–207.
- [24] M.F. Brin, T.I. Boodhoo, J.M. Pogoda, L.M. James, G. Demos, Y. Terashima, et al., Safety and tolerability of onabotulinumtoxinA in the treatment of facial lines: a meta-analysis of individual patient data from global clinical registration studies in 1678 participants, *J. Am. Acad. Dermatol.* 61 (6) (2009) 961–970 (e1–11).
- [25] F. Burzotta, M. De Vita, Y.L. Gu, T. Isshiki, T. Lefevre, A. Kaltoft, et al., Clinical impact of thrombectomy in acute ST-elevation myocardial infarction: an individual patient-data pooled analysis of 11 trials, *Eur. Heart J.* 30 (18) (2009) 2193–2203.
- [26] M. Buyse, P. Squifflet, B.J. Lange, T.A. Alonzo, R.A. Larson, J.E. Koltz, et al., Individual patient data meta-analysis of randomized trials evaluating IL-2 monotherapy as remission maintenance therapy in acute myeloid leukemia, *Blood* 117 (26) (2011) 7007–7013.
- [27] C. L. L. Trialists' Collaborative Group, Systematic review of purine analog treatment for chronic lymphocytic leukemia: lessons for future trials, *Haematologica* 97 (3) (2012) 428–436.
- [28] L. Chambrone, P.M. Preshaw, E.F. Rosa, P.A. Heasman, G.A. Romito, C.M. Pannuti, et al., Effects of smoking cessation on the outcomes of non-surgical periodontal therapy: a systematic review and individual patient data meta-analysis, *J. Clin. Periodontol.* 40 (6) (2013) 607–615.
- [29] Chemoradiotherapy for Cervical Cancer Meta-Analysis Collaboration, Reducing uncertainties about the effects of chemoradiotherapy for cervical cancer: a systematic review and meta-analysis of individual patient data from 18 randomized trials, *J. Clin. Oncol.* 26 (35) (2008) 5802–5812.
- [30] Childhood Acute Lymphoblastic Leukaemia Collaborative Group, Beneficial and harmful effects of anthracyclines in the treatment of childhood acute lymphoblastic leukaemia: a systematic review and meta-analysis, *Br. J. Haematol.* 145 (3) (2009) 376–388.
- [31] R. Clarke, J. Halsey, S. Lewington, E. Lonn, J. Armitage, J.E. Manson, et al., Effects of lowering homocysteine levels with B vitamins on cardiovascular disease, cancer, and cause-specific mortality: meta-analysis of 8 randomized trials involving 37,485 individuals, *Arch. Intern. Med.* 170 (18) (2010) 1622–1631.
- [32] J.P. Daniels, L. Middleton, T. Xiong, R. Champaer, N.P. Johnson, E.M. Lichten, et al., Individual patient data meta-analysis of randomized evidence to assess the effectiveness of laparoscopic uterosacral nerve ablation in chronic pelvic pain, *Hum. Reprod. Update* 16 (6) (2010) 568–576.
- [33] T. De Backer, R. Vander Stichele, P. Lebert, L. Van Bortel, Naftidrofuryl for intermittent claudication: meta-analysis based on individual patient data, *BMJ* 338 (2009) b603.
- [34] G. De Luca, M.T. Dirksen, C. Spaulding, H. Kelbaek, M. Schali, L. Thuesen, et al., Drug-eluting vs bare-metal stents in primary angioplasty: a pooled patient-level meta-analysis of randomized trials, *Arch. Intern. Med.* 172 (8) (2012) 611–621 (discussion 21–2).
- [35] G. De Luca, C.M. Gibson, F. Bellandi, S. Murphy, M. Maioli, M. Noc, et al., Early glycoprotein IIb/IIIa inhibitors in primary angioplasty (EGYPT) cooperation: an individual patient data meta-analysis, *Heart* 94 (12) (2008) 1548–1558.
- [36] A. de Waha, A. Dibra, S. Kufner, D. Baumgart, M. Sabate, A. Maresta, et al., Long-term outcome after sirolimus-eluting stents versus bare metal stents in patients with diabetes mellitus: a patient-level meta-analysis of randomized trials [Erratum (Erratum appears in Clin Res Cardiol. 2011 Sep;100(9):811–3)] *Clin. Res. Cardiol.* 100 (7) (2011) 561–570.
- [37] A.K. Devaiah, M.T. Andreoli, Treatment of esthesioneuroblastoma: a 16-year meta-analysis of 361 patients, *Laryngoscope* 119 (7) (2009) 1412–1416.
- [38] M. Di Maio, N. Lama, A. Morabito, E.F. Smit, V. Georgoulas, K. Takeda, et al., Clinical assessment of patients with advanced non-small-cell lung cancer eligible for second-line chemotherapy: a prognostic score from individual data of nine randomised trials, *Eur. J. Cancer* 46 (4) (2009) 735–743.
- [39] Dipart Group, Patient level pooled analysis of 68,500 patients from seven major vitamin D fracture trials in US and Europe, *BMJ* 340 (2010) b5463.
- [40] J.W. Eikelboom, D.J. Quinlan, M. O'Donnell, Major bleeding, mortality, and efficacy of fondaparinux in venous thromboembolism prevention trials, *Circulation* 120 (20) (2009) 2006–2011.
- [41] A.J. Farmer, R. Perera, A. Ward, C. Heneghan, J. Oke, A.H. Barnett, et al., Meta-analysis of individual patient data in randomised trials of self monitoring of blood glucose in people with non-insulin treated type 2 diabetes, *BMJ* 344 (2012) e486.
- [42] M. Flather, J.-W. Rhee, D.B. Boothroyd, E. Boersma, M.M. Brooks, D. Carrie, et al., The effect of age on outcomes of coronary artery bypass surgery compared with balloon angioplasty or bare-metal stent implantation among patients with multivessel coronary disease: a collaborative analysis of individual patient data from 10 randomized trials, *J. Am. Coll. Cardiol.* 60 (21) (2012) 2150–2157.
- [43] K.A.A. Fox, T.C. Clayton, P. Damman, S.J. Pocock, R.J. de Winter, J.G.P. Tijssen, et al., Long-term outcome of a routine versus selective invasive strategy in patients with non-ST-segment elevation acute coronary syndrome: a meta-analysis of individual patient data, *J. Am. Coll. Cardiol.* 55 (22) (2010) 2435–2445.
- [44] M. Fruh, E. Rolland, J.-P. Pignon, L. Seymour, K. Ding, H. Tribodet, et al., Pooled analysis of the effect of age on adjuvant cisplatin-based chemotherapy for completely resected non-small-cell lung cancer, *J. Clin. Oncol.* 26 (21) (2008) 3573–3581.
- [45] R. Fu, S. Selph, M. McDonagh, K. Peterson, A. Tiwari, R. Chou, et al., Effectiveness and harms of recombinant human bone morphogenetic protein-2 in spine fusion: a systematic review and meta-analysis, *Ann. Intern. Med.* 158 (12) (2013) 890–902.
- [46] T. Garcia-Berrococo, D. Giral, A. Bustamante, T. Etgen, J.K. Jensen, J.C. Sharma, et al., B-type natriuretic peptides and mortality after stroke: a systematic review and meta-analysis, *Neurology* 81 (23) (2013) 1976–1985.
- [47] Gastric Group, K. Oba, X. Paoletti, Y.J. Bang, H. Bleiberg, T. Burzykowski, et al., Role of chemotherapy for advanced/recurrent gastric cancer: an individual-patient-data meta-analysis, *Eur. J. Cancer* 49 (7) (2013) 1565–1577.
- [48] Gastric Group, X. Paoletti, K. Oba, T. Burzykowski, S. Michiels, Y. Ohashi, et al., Benefit of adjuvant chemotherapy for resectable gastric cancer: a meta-analysis, *JAMA* 303 (17) (2010) 1729–1737.
- [49] J.R. Geddes, J.R. Calabrese, G.M. Goodwin, Lamotrigine for treatment of bipolar depression: independent meta-analysis and meta-regression of individual patient data from five randomised trials, *Br. J. Psychiatry* 194 (1) (2009) 4–9.
- [50] F.L. Giorgino, C.G. Egan, Use of isoxsuprine hydrochloride as a tocolytic agent in the treatment of preterm labour: a systematic review of previous literature, *Arzneimittelforschung* 60 (7) (2010) 415–420.
- [51] L.L. Gluud, G. Dam, M. Borre, I. Les, J. Cordoba, G. Marchesini, et al., Oral branched-chain amino acids have a beneficial effect on manifestations of hepatic encephalopathy in a systematic review with meta-analyses of randomized controlled trials, *J. Nutr.* 143 (8) (2013) 1263–1268.
- [52] L.L. Gluud, F.K. Knop, T. Vilsbøll, Effects of lixisenatide on elevated liver transaminases: systematic review with individual patient data meta-analysis of randomised controlled trials on patients with type 2 diabetes, *BMJ Open* 4 (12) (2014) e005325.
- [53] M. Gomes, I. Ribeiro, M. Warsame, H. Karunajeewa, M. Petzold, Rectal artemisinins for malaria: a review of efficacy and safety from individual patient data in clinical studies, *BMC Infect. Dis.* 8 (2008) 39.
- [54] S. Goodacre, J.W. Stevens, A. Pandor, E. Poku, S. Ren, A. Cantrell, et al., Prehospital noninvasive ventilation for acute respiratory failure: systematic review, network meta-analysis, and individual patient data meta-analysis, *Acad. Emerg. Med.* 21 (9) (2014) 960–970.
- [55] E. Groeneweld, K.A. Broeze, M.J. Lambers, M. Haapsamo, K. Dirckx, B.C. Schoot, et al., Is aspirin effective in women undergoing in vitro fertilization (IVF)? Results from an individual patient data meta-analysis (IPD MA), *Hum. Reprod. Update* 17 (4) (2011) 501–509.

- [56] M. Haasova, F.C. Warren, M. Ussher, K. Janse Van Rensburg, G. Faulkner, M. Croyley, et al., The acute effects of physical activity on cigarette cravings: systematic review and meta-analysis with individual participant data, *Addiction* 108 (1) (2013) 26–37.
- [57] R.A. Hansen, L. Song, C.G. Moore, A.W. Gilseman, M.M. Kim, M.O. Calloway, et al., Effect of ropinirole on sleep outcomes in patients with restless legs syndrome: meta-analysis of pooled individual patient data from randomized controlled trials, *Pharmacother. J. Hum. Pharmacol. Drug Ther.* 29 (3) (2009) 255–262.
- [58] M.A. Hlatky, D.B. Boothroyd, D.M. Bravata, E. Boersma, J. Booth, M.M. Brooks, et al., Coronary artery bypass surgery compared with percutaneous coronary interventions for multivessel disease: a collaborative analysis of individual patient data from ten randomised trials, *Lancet* 373 (9670) (2009) 1190–1197.
- [59] P.D. Home, A. Fritsche, S. Schinzel, M. Massi-Benedetti, Meta-analysis of individual patient data to assess the risk of hypoglycaemia in people with type 2 diabetes using NPH insulin or insulin glargine, *Diabetes Obes. Metab.* 12 (9) (2010) 772–779.
- [60] J. Jefferis, R. Perera, H. Everitt, H. van Weert, R. Rietveld, P. Glasziou, et al., Acute infective conjunctivitis in primary care: who needs antibiotics? An individual patient data meta-analysis, *Br. J. Gen. Pract.* 61 (590) (2011) e542–e548.
- [61] R.V. Jeger, P. Urban, S.M. Harkness, C.-H. Tseng, J.-C. Stauffer, T.H. Lejemtel, et al., Early revascularization is beneficial across all ages and a wide spectrum of cardiogenic shock severity: a pooled analysis of trials, *Acute Card. Care* 13 (1) (2011) 14–20.
- [62] B. Kassai, C. Chiron, S. Augier, M. Cucherat, E. Rey, F. Gueyffier, et al., Severe myoclonic epilepsy in infancy: a systematic review and a meta-analysis of individual patient data, *Epilepsia* 49 (2) (2008) 343–348.
- [63] D. Kendrick, M.C. Watson, C.A. Mulvaney, S.J. Smith, A.J. Sutton, C.A.C. Coupland, et al., Preventing childhood falls at home: meta-analysis and meta-regression, *Am. J. Prev. Med.* 35 (4) (2008) 370–379.
- [64] S.H. Kennedy, H.F. Andersen, M.E. Thase, Escitalopram in the treatment of major depressive disorder: a meta-analysis, *Curr. Med. Res. Opin.* 25 (1) (2009) 161–175.
- [65] M.W.J. Koeter, W. van den Brink, P. Lehart, Effect of early and late compliance on the effectiveness of acamprosate in the treatment of alcohol dependence, *J. Subst. Abuse Treat.* 39 (3) (2010) 218–226.
- [66] L. Koopman, A.W. Hoes, P.P. Glasziou, C.L. Appelman, P. Burke, D.P. McCormick, et al., Antibiotic therapy to prevent the development of asymptomatic middle ear effusion in children with acute otitis media: a meta-analysis of individual patient data, *Arch. Otolaryngol. Head Neck Surg.* 134 (2) (2008) 128–132.
- [67] S. Kufner, A. de Waha, F. Tomai, S.-W. Park, S.-W. Lee, D.-S. Lim, et al., A meta-analysis of specifically designed randomized trials of sirolimus-eluting versus paclitaxel-eluting stents in diabetic patients with coronary artery disease, *Am. Heart J.* 162 (4) (2011) 740–747.
- [68] S. Laporte, J. Liotier, L. Bertolotti, F.X. Kleber, G.F. Pineo, C. Chapelle, et al., Individual patient data meta-analysis of enoxaparin vs. unfractionated heparin for venous thromboembolism prevention in medical patients, *J. Thromb. Haemost.* 9 (3) (2011) 464–472.
- [69] P. Leheret, G. Cheron, G.A. Calatayud, J.-P. Cezard, P.G. Castrellon, J.-M.M. Garcia, et al., Racecadotril for childhood gastroenteritis: an individual patient data meta-analysis, *Dig. Liver Dis.* 43 (9) (2011) 707–713.
- [70] S. Leroy, A. Fernandez-Lopez, R. Nikfar, C. Romanello, F. Bouissou, A. Gervais, et al., Association of procalcitonin with acute pyelonephritis and renal scars in pediatric UTI, *Pediatrics* 131 (5) (2013) 870–879.
- [71] C.L. Loprinzi, J. Sloan, V. Stearns, R. Slack, M. Iyengar, B. Diekmann, et al., Newer antidepressants and gabapentin for hot flashes: an individual patient pooled analysis, *J. Clin. Oncol.* 27 (17) (2009) 2831–2837.
- [72] H. Ludwig, J. Crawford, A. Osterborg, J. Vansteenkiste, D.H. Henry, A. Fleishman, et al., Pooled analysis of individual patient-level data from all randomized, double-blind, placebo-controlled trials of darbepoetin alfa in the treatment of patients with chemotherapy-induced anemia, *J. Clin. Oncol.* 27 (17) (2009) 2838–2847.
- [73] P. Mathurin, J. O'Grady, R.L. Carithers, M. Phillips, A. Louvet, C.L. Mendenhall, et al., Corticosteroids improve short-term survival in patients with severe alcoholic hepatitis: meta-analysis of individual patient data, *Gut* 60 (2) (2011) 255–260.
- [74] A. Mauguen, C. Le Pechoux, M.I. Saunders, S.E. Schild, A.T. Turrissi, M. Baumann, et al., Hyperfractionated or accelerated radiotherapy in lung cancer: an individual patient data meta-analysis, *J. Clin. Oncol.* 30 (22) (2012) 2788–2797.
- [75] M.N. Mavros, S.C. Mayo, O. Hyder, T.M. Pawlik, A systematic review: treatment and prognosis of patients with fibrolamellar hepatocellular carcinoma, *J. Am. Coll. Surg.* 215 (6) (2012) 820–830.
- [76] W.G. Melsen, M.M. Rovers, R.H. Groenwold, D.C. Bergmans, C. Camus, T.T. Bauer, et al., Attributable mortality of ventilator-associated pneumonia: a meta-analysis of individual patient data from randomised prevention studies, *Lancet Infect. Dis.* 13 (8) (2013) 665–671.
- [77] S. Morita, I. Okamoto, K. Kobayashi, K. Yamazaki, H. Asahina, A. Inoue, et al., Combined survival analysis of prospective clinical trials of gefitinib for non-small cell lung cancer with EGFR mutations, *Clin. Cancer Res.* 15 (13) (2009) 4493–4498.
- [78] S.G. Muthuri, S. Venkatesan, P.R. Myles, J. Leonardi-Bee, T.S. Al Khuwaitir, A. Al Mamun, et al., Effectiveness of neuraminidase inhibitors in reducing mortality in patients admitted to hospital with influenza A H1N1pdm09 virus infection: a meta-analysis of individual participant data, *Lancet Respir. Med.* 2 (5) (2014) 395–404.
- [79] S.J. Nicholls, G. Brandrup-Wognsen, M. Palmer, P.J. Barter, Meta-analysis of comparative efficacy of increasing dose of atorvastatin versus rosuvastatin versus simvastatin on lowering levels of atherogenic lipids (from VOYAGER), *Am. J. Cardiol.* 105 (1) (2010) 69–76.
- [80] NSCLC Meta-Analyses Collaborative Group, Chemotherapy in addition to supportive care improves survival in advanced non-small-cell lung cancer: a systematic review and meta-analysis of individual patient data from 16 randomized controlled trials, *J. Clin. Oncol.* 26 (28) (2008) 4617–4625.
- [81] NSCLC Meta-analysis Collaborative Group, Adjuvant chemotherapy, with or without postoperative radiotherapy, in operable non-small-cell lung cancer: two meta-analyses of individual patient data, *Lancet* 375 (9722) (2010) 1267–1277.
- [82] NSCLC Meta-analysis Collaborative Group, Preoperative chemotherapy for non-small-cell lung cancer: a systematic review and meta-analysis of individual participant data, *Lancet* 383 (9928) (2014) 1561–1571.
- [83] K. Oba, Efficacy of adjuvant chemotherapy using tegafur-based regimen for curatively resected gastric cancer: update of a meta-analysis, *Int. J. Clin. Oncol.* 14 (2) (2009) 85–89.
- [84] K. Oba, M. Kobayashi, T. Matsui, Y. Koda, J. Sakamoto, Individual patient based meta-analysis of lenitin for unresectable/recurrent gastric cancer, *Anticancer Res.* 29 (7) (2009) 2739–2745.
- [85] N.J. Opdam, F.H. van de Sande, E. Bronkhorst, M.S. Cenci, P. Bottenberg, U. Pallesen, et al., Longevity of posterior composite restorations: a systematic review and meta-analysis, *J. Dent. Res.* 93 (10) (2014) 943–949.
- [86] G.I. Papakostas, M.H. Trivedi, J.E. Alpert, C.A. Seifert, A. Krishen, E.P. Goodale, et al., Efficacy of bupropion and the selective serotonin reuptake inhibitors in the treatment of anxiety symptoms in major depressive disorder: a meta-analysis of individual patient data from 10 double-blind, randomized clinical trials, *J. Psychiatr. Res.* 42 (2) (2008) 134–140.
- [87] G. Patti, C.P. Cannon, S.A. Murphy, S. Mega, V. Pasceri, C. Briguori, et al., Clinical benefit of statin pretreatment in patients undergoing percutaneous coronary intervention: a collaborative patient-level meta-analysis of 13 randomized studies, *Circulation* 123 (15) (2011) 1622–1632.
- [88] R.M. Pearce, J.D. Belsey, J.N. Cole, E.D. Bennett, Effect of dexamethasone infusion on mortality following major surgery: individual patient data meta-regression analysis of published clinical trials, *Crit. Care Med.* 36 (4) (2008) 1323–1329.
- [89] R. Pettengell, M. Schwenkglenks, P. Bacon, S. Lawrinson, U. Duehrsen, Pegfilgrastim primary prophylaxis in patients with non-Hodgkin lymphoma: results from an integrated analysis, *Hematol. Oncol.* 29 (4) (2011) 177–184.
- [90] M.J. Piccart-Gebhart, T. Burzykowski, M. Buyse, G. Sledge, J. Carmichael, H.-J. Luck, et al., Taxanes alone or in combination with anthracyclines as first-line therapy of patients with metastatic breast cancer, *J. Clin. Oncol.* 26 (12) (2008) 1980–1986.
- [91] J.-P. Pignon, H. Tribodet, G.V. Scagliotti, J.-Y. Douillard, F.A. Shepherd, R.J. Stephens, et al., Lung adjuvant cisplatin evaluation: a pooled analysis by the LACE Collaborative Group, *J. Clin. Oncol.* 26 (21) (2008) 3552–3559.
- [92] A. Presson, T.B. Moore, P. Kempert, Efficacy of high-dose chemotherapy and autologous stem cell transplant for recurrent Wilms' tumor: a meta-analysis, *J. Pediatr. Hematol. Oncol.* 32 (6) (2010) 454–461.
- [93] J.M. Raboud, C. Diong, A. Carr, S. Grinspoon, K. Mulligan, J. Sutinen, et al., A meta-analysis of six placebo-controlled trials of thiazolidinedione therapy for HIV lipodystrophy, *HIV Clin. Trials* 11 (1) (2010) 39–50.
- [94] F.A. Raja, N. Counsell, N. Colombo, J. Pfisterer, A. du Bois, M.K. Parmar, et al., Platinum versus platinum-combination chemotherapy in platinum-sensitive recurrent ovarian cancer: a meta-analysis using individual patient data, *Ann. Oncol.* 24 (12) (2013) 3028–3034.
- [95] L. Rejnmark, A. Avenell, T. Masud, F. Anderson, H.E. Meyer, K.M. Sanders, et al., Vitamin D with calcium reduces mortality: patient level pooled analysis of 70,528 patients from eight major vitamin D trials, *J. Clin. Endocrinol. Metab.* 97 (8) (2012) 2670–2681.
- [96] R.N. Rodseth, B.M. Biccari, Y. Le Manach, D.I. Sessler, G.A. Lurati Buse, L. Thabane, et al., The prognostic value of pre-operative and post-operative B-type natriuretic peptides in patients undergoing noncardiac surgery: B-type natriuretic peptide and N-terminal fragment of pro-B-type natriuretic peptide: a systematic review and individual patient data meta-analysis, *J. Am. Coll. Cardiol.* 63 (2) (2014) 170–180.
- [97] U. Ronellenfitsch, M. Schwarzbach, R. Hofheinz, P. Kienle, M. Kieser, T.E. Slinger, et al., Preoperative chemo(radio)therapy versus primary surgery for gastroesophageal adenocarcinoma: systematic review with meta-analysis combining individual patient and aggregate data, *Eur. J. Cancer* 49 (15) (2013) 3149–3158.
- [98] E. Rosenfeld, A. Beyerlein, M. Hadders-Algra, K. Kennedy, A. Singhal, M. Fewtrell, et al., IPD meta-analysis shows no effect of LC-PUFA supplementation on infant growth at 18 months, *Acta Paediatr.* 98 (1) (2009) 91–97.
- [99] A. Rossi, P. Chiodini, J.M. Sun, M.E. O'Brien, C. von Plessen, F. Barata, et al., Six versus fewer planned cycles of first-line platinum-based chemotherapy for non-small-cell lung cancer: a systematic review and meta-analysis of individual patient data, *Lancet Oncol.* 15 (11) (2014) 1254–1262.
- [100] A. Rossi, M. Di Maio, P. Chiodini, R.M. Rudd, H. Okamoto, D.V. Skarlos, et al., Carboplatin- or cisplatin-based chemotherapy in first-line treatment of small-cell lung cancer: the COCIS meta-analysis of individual patient data, *J. Clin. Oncol.* 30 (14) (2012) 1692–1698.
- [101] D. Sargent, A. Sobrero, A. Grothey, M.J. O'Connell, M. Buyse, T. Andre, et al., Evidence for cure by adjuvant therapy in colon cancer: observations based on individual patient data from 20,898 patients on 18 randomized trials, *J. Clin. Oncol.* 27 (6) (2009) 872–877.
- [102] P. Schuetz, M. Briel, M. Christ-Crain, D. Stolz, L. Bouadma, M. Wolff, et al., Procalcitonin to guide initiation and duration of antibiotic treatment in acute respiratory infections: an individual patient data meta-analysis, *Clin. Infect. Dis.* 55 (5) (2012) 651–662.
- [103] S. Shepperd, H. Doll, J. Broad, J. Gladman, S. Illife, P. Langhorne, et al., Early discharge hospital at home, *Cochrane Database Syst. Rev.* 1 (2009) CD000356.

- [104] D.D. Sin, D. Tashkin, X. Zhang, F. Radner, U. Sjobring, A. Thoren, et al., Budesonide and the risk of pneumonia: a meta-analysis of individual patient data, *Lancet* 374 (9691) (2009) 712–719.
- [105] F.M. Struik, Y. Lacasse, R.S. Goldstein, H.A. Kerstjens, P.J. Wijkstra, Nocturnal noninvasive positive pressure ventilation in stable COPD: a systematic review and individual patient data meta-analysis, *Respir. Med.* 108 (2) (2014) 329–337.
- [106] M.E. Thase, S.G. Kornstein, J.-M. Germain, Q. Jiang, C. Guico-Pabia, P.T. Ninan, An integrated analysis of the efficacy of desvenlafaxine compared with placebo in patients with major depressive disorder, *CNS Spectr.* 14 (3) (2009) 144–154.
- [107] S.R. Walsh, J.R. Boyle, A.G. Lynch, U. Sadat, J.P. Carpenter, T.Y. Tang, et al., Suprarenal endograft fixation and medium-term renal function: systematic review and meta-analysis, *J. Vasc. Surg.* 47 (6) (2008) 1364–1370.
- [108] G. Yothers, D.J. Sargent, N. Wolmark, R.M. Goldberg, M.J. O'Connell, J.K. Benedetti, et al., Outcomes among black patients with stage II and III colon cancer receiving chemotherapy: an analysis of ACCENT adjuvant trials, *J. Natl. Cancer Inst.* 103 (20) (2011) 1498–1506.
- [109] J. Young, A. De Sutter, D. Merenstein, G.A. van Essen, L. Kaiser, H. Varonen, et al., Antibiotics for adults with clinically diagnosed acute rhinosinusitis: a meta-analysis of individual patient data, *Lancet* 371 (9616) (2008) 908–914.
- [110] Y. Yuan, W. Xiang, M. Qing, L. Yanhui, L. Jiewen, M. Yunhe, Survival analysis for valproic acid use in adult glioblastoma multiforme: a meta-analysis of individual patient data and a systematic review, *Seizure* 23 (10) (2014) 830–835.
- [111] M.C. Simmonds, J. Tierney, J. Bowden, J.P.T. Higgins, Meta-analysis of time-to-event data: a comparison of two-stage methods, *Res. Synth. Methods* 2 (3) (2011) 139–149.
- [112] I.R. White, P. Royston, A.M. Wood, Multiple imputation using chained equations: issues and guidance for practice, *Stat. Med.* 30 (4) (2011) 377–399.