

Active Compression-Decompression Resuscitation and Impedance Threshold Device for Out-of-Hospital Cardiac Arrest: A Systematic Review and Metaanalysis of Randomized Controlled Trials*

Chih-Hung Wang, MD^{1,2}; Min-Shan Tsai, MD, PhD¹; Wei-Tien Chang, MD, PhD¹;
Chien-Hua Huang, MD, PhD¹; Matthew Huei-Ming Ma, MD, PhD¹; Wen-Jone Chen, MD, PhD^{1,3};
Cheng-Chung Fang, MD¹; Shyr-Chyr Chen, MD, MBA¹; Chien-Chang Lee, MD, MSc^{2,4}

Objective: Active compression-decompression resuscitation and impedance threshold device are proposed to improve survival of patients of cardiac arrest by lowering intrathoracic pressure and increasing cardiac output. The results of clinical studies of active compression-decompression resuscitation or impedance threshold device were controversial. This metaanalysis pooled results of randomized controlled trials to examine whether active compression-decompression resuscitation or impedance threshold device would improve outcomes of out-of-hospital cardiac arrest in comparison with standard cardiopulmonary resuscitation and to explore factors modifying these effects.

Data Sources: Medline and Embase were searched from inception to September 2013.

Study Selection: Randomized controlled trials comparing active compression-decompression resuscitation or impedance threshold device with standard cardiopulmonary resuscitation in out-of-hospital cardiac arrest patients were selected. There were no restrictions for language, population, or publication year.

*See also p. 929.

¹Department of Emergency Medicine, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan.

²Department of Emergency Medicine, National Taiwan University Hospital Yunlin Branch, Douliu, Yunlin, Taiwan.

³Department of Emergency Medicine, Lotung Poh-Ai Hospital, Yilan, Taiwan.

⁴Department of Epidemiology, Harvard School of Public Health, Boston, MA.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website (<http://journals.lww.com/ccmjournal>).

The authors have disclosed that they do not have any potential conflicts of interest.

Address requests for reprints to: Chien-Chang Lee, MD, MSc, Department of Emergency Medicine, National Taiwan University Hospital Yunlin Branch, No. 579, Sec. 2, Yunlin Road, Douliu, Yunlin 640, Taiwan. E-mail: chnchnglee@ntu.edu.tw

Copyright © 2015 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

DOI: 10.1097/CCM.0000000000000820

Data Extraction: Data on study characteristics, including patients, intervention details, and outcome measures, were independently extracted.

Data Synthesis: Fifteen trials, including 16,088 patients, were identified from 331 potentially relevant references. Return of spontaneous circulation was designated as the primary outcome. The pooled result showed no significant improvement in return of spontaneous circulation by active compression-decompression resuscitation or impedance threshold device in comparison with standard cardiopulmonary resuscitation (risk ratio, 1.04; 95% CI, 0.93–1.16; *P*, 46%). There was also no significant difference in survival or neurologic outcome at hospital discharge between active compression-decompression resuscitation or impedance threshold device and standard cardiopulmonary resuscitation. The meta-regression indicated that this minimal effect might be modified by two important prognostic factors, that is, witnessed status and response time. After adjustment of these two factors, impedance threshold device appeared to improve return of spontaneous circulation, which could be further augmented by advanced airway use.

Conclusions: Active compression-decompression resuscitation or impedance threshold device seemed not to improve return of spontaneous circulation in out-of-hospital cardiac arrest patients. The meta-regression indicated two probable prognostic factors causing this minimal effect. Nonetheless, these findings referred to differences between trials and could not necessarily be extrapolated to individual patients. The individual patient-level extrapolation may need to be solved by a future randomized controlled trial. (*Crit Care Med* 2015; 43:889–896)

Key Words: cardiopulmonary resuscitation; critical care; emergency medical service; emergency medicine; heart arrest; metaanalysis

Out-of-hospital cardiac arrest (OHCA) is a clinical condition still characterized by a poor prognosis (1). High-quality cardiopulmonary resuscitation (CPR)

with complete chest recoil is addressed in 2010 American Heart Association guidelines (2). However, standard cardiopulmonary resuscitation (STR), defined as manual chest compressions with rescue breathing, is inherently inefficient (3). The use of active compression-decompression resuscitation (ACDR) or impedance threshold device (ITD) was proposed to improve CPR efficiency.

With the use of a hand-held device with a suction cup, ACDR is performed to lower the intrathoracic pressure by active chest decompression and thereby augment the venous return. Experimental studies on animal models of cardiac arrest have shown that ACDR was associated with a higher rate of return of spontaneous circulation (ROSC) than STR (4). Clinical studies, however, showed inconsistent results. A Cochrane metaanalysis concluded that ACDR might not be superior to STR (5).

ITD, a valve device used to reduce air entry into lungs during chest decompression, is proposed to produce a similar effect as ACDR by maintaining negative intrathoracic pressure during CPR (6). A metaanalysis demonstrated that ITD improved ROSC and early survival (7). However, this analysis included a trial comparing ACDR/ITD with STR (8), of which the observed benefit might not be totally ascribed to ITD alone.

The clinical evidence for the effect of ACDR or ITD in OHCA was conflicting (9, 10). This metaanalysis quantitatively pooled the results of randomized controlled trials (RCTs) to examine whether use of ACDR or ITD would improve outcomes of OHCA in comparison with STR and explore whether other factors would modify the effects of these devices.

MATERIALS AND METHODS

Data Sources and Searches

We performed this metaanalysis in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (11). Two authors (C.-H.W., C.-C.L.) independently searched Medline and Embase from inception to September 2013 according to a prespecified protocol. The search terms included “active compression-decompression,” “impedance threshold device,” “cardiac arrest,” and “cardiopulmonary resuscitation.” There were no restrictions on language, population, or publication year. We did not search abstracts from conferences, proceedings, or clinical trial registries. Instead, we manually reviewed the bibliographies of relevant studies, reviews, and metaanalyses to identify references we may have missed during our primary search. The authors of the included trials were contacted for missing information, if necessary. This study was reviewed by the local institutional review board and is exempt from requiring approval.

Study Selection

Two authors (C.-H.W., M.-S.T.) independently scanned the titles and abstracts of all retrieved articles and selected those that were pertinent to this review. The following prespecified inclusion criteria were used: 1) being an RCT, 2) comparisons were made between the use of ACDR or ITD and STR in OHCA, and 3) the results included ROSC, survival, or

neurologic outcome. If studies shared the same cohort, the study with more patients was selected into analysis.

After retrieving full articles of potentially relevant trials, two reviewers (C.-H.W., M.-S.T.) independently assessed each study's eligibility on the basis of the inclusion criteria and settled differing opinions by consensus or consultation with a third investigator (W.-J.C.).

Data Extraction and Quality Assessment

Four authors independently extracted data by using a prespecified protocol (W.-T.C., C.-H.H., M.H.-M.M., C.-C.F.). Each trial was reviewed independently by two of these four authors to ensure the correctness of the extracted data, which included 1) study design, 2) study population characteristics, 3) details of interventions, and 4) types of outcomes measured.

ROSC was prespecified as the primary outcome in the analysis. Other prespecified secondary outcomes included survival for 24 hours, survival to hospital discharge, and satisfactory neurologic outcome at hospital discharge.

The Cochrane risk of bias tool was adopted to assess the risk of bias for each RCT (12). Each RCT was scored as “high risk,” “low risk,” or “unclear” with respect to random sequence generation, allocation concealment, blinding process, incomplete outcome data, and selective reporting (12). Discrepancies in assessment were resolved through discussions between the extraction authors or consultation with the supervising investigator (S.-C.C.).

Data Synthesis and Analysis

Metaanalysis and meta-regression were performed using the *rmeta* and *metafor* packages in the R 2.15.3 software (R Foundation for Statistical Computing, Vienna, Austria). Two-sided *p* value of less than 0.05 was considered statistically significant. The estimated risk ratio (RR) and the 95% CIs of each individual study were recorded for computing the pooled (i.e., weighted averaged) estimate of effect size from all collected studies. The response variable to be analyzed in metaanalysis and meta-regression was the transformed “log(RR)” for a more symmetric response distribution.

Metaanalysis was conducted first. Heterogeneity in the estimated RRs between individual studies was examined using 1) the *I*² statistic, in which *I*² more than 40% indicated a substantial level of heterogeneity, and 2) the test of heterogeneity, in which *p* value of less than 0.05 indicated heterogeneity. If heterogeneity was not evident, fixed-effects summary estimates (Mantel-Haenszel method) were reported in forest plots; otherwise, random-effects summary estimates (DerSimonian-Laird method [13]) were reported.

If significant statistical heterogeneity was found between individual studies in metaanalysis, subgroup analysis was performed by stratifying collected studies according to the interventions (ACDR or ITD) used in experimental groups. Next, to account for the heterogeneity, meta-regression was performed with potentially relevant covariates (listed in **Table 1**) as moderators. The stepwise variable selection procedure (with iterations between the forward and backward steps) was applied to obtain the final regression model. All the univariate significant/

TABLE 1. Characteristics of the Included Randomized Controlled Trials

Study	Year	n	Country	Mean Age/ Male %	Witnessed Cardiac Arrest %	Bystander Cardio- pulmonary Resuscitation %	Response Time	Initial Arrest Rhythm (Shockable Rhythm/ Pulseless Electrical Activity/ Asystole %)	ITD Attach- ment Site (Endotrache- al Tube/ Advanced Airways %)
ACDR vs STR									
Lurie et al (17)	1994	130	United States	67.1/71.5	68.5	36.9	3.4	61.5/25.4/13.1	NA
Schwab et al (18) (Fresno)	1995	253	United States	64.5/61.3	61.3	33.6	4.9	32.8/17.8/42.3	NA
Schwab et al (18) (San Francisco)	1995	607	United States	65.5/67.1	52.6	18.1	3.95	22.5/36.6/40.9	NA
Luiz et al (19)	1996	56	Germany	68.9/73.2	73.2	30.4	6	26.8/17.8/55.4	NA
Mauer et al (20)	1996	220	Germany	68.8/NA	68.6	15.5	7.2	45/34.1/20.9	NA
Stiell et al (21)	1996	1,011	Canada	68/63.6	59.2	19.6	NA	32.5/21.9/41.3	NA
Goralski et al (22)	1998	145	France	61.6/73.1	81.5	22.4	16.5	20.7/3.4/75.9	NA
Mauer et al (23)	1998	120	Germany	68.3/NA	69.2	8.3	7.65	45.2/34.1/20.8	NA
Nolan et al (24)	1998	576	United Kingdom	67.2/68.4	70.3	32.5	6.5	60.2/14.6/25.2	NA
Plaisance et al (16)	1999	750	France	58.5/68	73.9	7.6	NA	12.5/6/82	NA
Skogvoll and Wik (25)	1999	302	Norway	71/73	81.5	57.4	9	57/NA/NA	NA
STR/ITD vs STR									
Aufderheide et al (26)	2005	230	United States	65.5/61.3	58.7	22.2	4.7	25.7/22.6/50.9	NA/NA
Pirrallo et al (27)	2005	22	United States	61.2/59.1	45.5	22.7	8.7	18.2/31.8/45.5	100/100
Aufderheide et al (9)	2011	8,718	United States/ Canada	66.9/64.6	48.7	38.5	5.8	24.5/23.9/44.2	76.2/86.9
ACDR/ITD vs STR									
Wolcke et al (8)	2003	210	Germany	67/61.9	74.8	24.3	5.95	40/29.5/30.5	100/100
Frascone et al (15)	2013	2,738	United States	63.8/62.4	54.6	40.8	6.6	25.1/23.5/50	NA/86.7

ITD = impedance threshold device, ACDR = active compression-decompression resuscitation, STR = standard cardiopulmonary resuscitation, NA = not applicable.

nonsignificant covariates and some of their interaction terms (including ACDR \times ITD and ITD \times response time) were put on the variable list to be selected. The significance levels for entry and for stay were set to 0.15 (or larger) for being conservative. With the aid of substantive knowledge, the final regression model was identified manually by dropping the covariates with *p* value of more than 0.05 one at a time until all regression coefficients were statistically significant. Generalized additive models were fitted to detect nonlinear effects of continuous covariates and identify appropriate cutoff point(s) for dichotomizing a continuous covariate, if necessary, during the stepwise variable selection procedure.

In meta-regression, if the test of residual heterogeneity was not statistically significant, the result of a fixed-effects meta-regression model was reported; otherwise, that of mixed-effects meta-regression model with the Knapp and Hartung adjustment was reported instead. The potential publication bias was examined by funnel plot.

RESULTS

Search Results and Description

The review process is shown in **Figure 1**. Two trials (10, 14) were excluded because more patients were enrolled from the




Figure 1. Flow diagram of the literature search. The literature search process was shown. Two studies were excluded after full-text review because more patients were enrolled from similar cohorts in later published trials.

similar cohorts in later published trials (15, 16). We identified 15 RCTs (8, 9, 15–27) including 16,088 patients for metaanalysis (Table 1). According to interventions in experimental groups, the included RCTs were categorized into three

subgroups: ACDR versus STR, STR/ITD versus STR, and ACDR/ITD versus STR. The trial by Schwab et al (18) provided data of two cities, which were reported separately. The trial by Stiell et al (21) enrolled patients of OHCA and in-hospital cardiac arrest; only the data of OHCA were presented.

Patients with noncardiac causes of arrest were excluded in six trials (8, 20, 23, 25–27). Patients were excluded from trials if the delay between cardiac arrest and basic CPR exceeded 15 minutes in three trials (8, 9, 21) and 30 minutes in another three trials (16, 22, 24). In most trials, the response time from dispatch to the arrival of first emergency medical service (EMS) team (either basic life support or advanced life support team) was within 10 minutes. Two trials provided time from collapse to EMS arrival instead, which were 5.1 minutes (21) and 9.2 minutes (16), respectively. In trials involving ITD use, ITD was used with endotracheal tube (ETT) in two trials (8, 27), whereas in the other two trials (9, 15), ITD could be attached between the ventilation bag and a facemask or an advanced airway (e.g., supraglottic airways or ETT).

As shown in **Supplemental Table 1** (Supplemental Digital Content 1, <http://links.lww.com/CCM/B152>), the blinding process could not be performed in trials involving ACDR. Several trials used alternating periods (day, week, or month) to allocate patients (clustered randomization) and showed a high risk of bias in allocation concealment (8, 16–18, 24). All trials were managed by intention-to-treat analysis with complete follow-up.

The outcomes are presented in **Supplemental Table 2** (Supplemental Digital Content 2, <http://links.lww.com/CCM/B153>). The timing of ROSC recording was not defined in most trials, and the results were pooled together as ROSC before hospital admission. Goralski et al (22) reported survival to ICU discharge, which was classified as survival to hospital discharge. Modified Rankin scale of up to 3 (28) and Glasgow–Pittsburgh cerebral performance category score of up to 2 (29)




Figure 2. Forest plot of the summary effect estimates of return of spontaneous circulation (ROSC) before hospital admission. The comparison was made between use of active compression-decompression resuscitation or impedance threshold device (intervention) and standard cardiopulmonary resuscitation (control) in patients with out-of-hospital cardiac arrest.

were categorized as satisfactory neurologic outcomes (30). Lurie et al (17) reported the number of patients who were awake, alert, and oriented to person, place, and time, which was also counted as satisfactory neurologic outcome.

Quantitative Data Synthesis

There was no significant difference in ROSC between use of ACDR or ITD and STR (RR, 1.04; 95% CI, 0.93–1.16; P , 46%) (Fig. 2) (Table 2). There were also no differences in survival and neurologic outcomes in either main or subgroup analysis (Table 2).

In the meta-regression, when all covariates except ITD attachment site in Table 3 were analyzed as independent variables with log(RR) of ROSC as the response variable, witnessed cardiac arrest ($p < 0.001$) and ITD use ($p = 0.002$) were noted to be positively associated with log(RR) of ROSC, whereas response time ($p = 0.004$) was inversely correlated with log(RR) of ROSC (Table 4). The general additive models plot demonstrated that log(RR) of ROSC would be higher

when the response time was less than 6.5 minutes, which was used as the cutoff point (Supplemental Fig. 1, Supplemental Digital Content 3, <http://links.lww.com/CCM/B154>). When interaction terms of ACDR \times ITD and ITD \times response time (< 6.5 min) were put into meta-regression model, these two interaction terms were not statistically significant.

When ITD attachment site was considered, the meta-regression was restricted to only four studies involving ITD use (8, 9, 15, 27). Multivariate analysis was not performed because of the limited number of trials. The univariate analysis on ITD attachment site showed that advanced airway use was positively associated with log(RR) of ROSC ($p = 0.018$) in these four studies (Table 3). The funnel plot did not show obvious publication bias (Supplemental Fig. 2, Supplemental Digital Content 4, <http://links.lww.com/CCM/B155>).

DISCUSSION

Previous metaanalyses examined the effect of ACDR or ITD for OHCA separately (5, 7). The interaction among ACDR, ITD,

TABLE 2. Pooled Outcomes of the Included Randomized Controlled Trials

Outcomes	No. of Studies	Events/No. of Intervention	Events/No. of Control	Risk Ratio (95% CI)	P
Use of ACDR or ITD vs STR					
ROSC before hospital admission	14	2,368/7,533	2,318/7,544	1.04 (0.93–1.16)	46%
Survival for 24 hr	4	217/875	147/881	1.49 (1.24–1.79)	0
Survival to hospital discharge	12	663/7,537	622/7,549	1.07 (0.96–1.18)	0
Satisfactory neurologic outcome at hospital discharge	10	472/7,362	428/7,349	1.24 (0.93–1.65)	19%
ACDR vs STR					
ROSC before hospital admission	9	489/1,530	495/1,629	1.00 (0.86–1.16)	48%
Survival for 24 hr	2	93/447	63/448	1.12 (0.44–2.87)	78%
Survival to hospital discharge	9	122/1,658	119/1,762	1.11 (0.87–1.41)	0
Satisfactory neurologic outcome at hospital discharge	7	94/1,483	82/1,562	1.24 (0.93–1.65)	0
STR/ITD vs STR					
ROSC before hospital admission	3	1,231/4,497	1246/4,473	0.98 (0.92–1.05)	0
Survival for 24 hr	1	19/114	14/116	1.38 (0.73–2.62)	NA
Survival to hospital discharge	1	357/4,373	355/4,345	1.00 (0.87–1.15)	NA
Satisfactory neurologic outcome at hospital discharge	1	254/4,373	260/4,345	0.97 (0.82–1.15)	NA
ACDR/ITD vs STR					
ROSC before hospital admission	2	648/1,506	577/1,442	1.21 (0.86–1.69)	79%
Survival for 24 hr	1	38/103	24/107	1.64 (1.07–2.54)	NA
Survival to hospital discharge	2	184/1,506	148/1,442	1.19 (0.97–1.46)	0
Satisfactory neurologic outcome at hospital discharge	2	124/1,506	86/1,442	1.39 (1.06–1.81)	0

ACDR = active compression-decompression resuscitation, ITD = impedance threshold device, STR = standard cardiopulmonary resuscitation, ROSC = return of spontaneous circulation, NA = not applicable.

TABLE 3. Results of Univariate Meta-Regression Analysis on Log Relative Risk of Return of Spontaneous Circulation

Variable	Odds Ratio (95% CI)	p
Mean age	0.979 (0.960–0.997)	0.023
Male (%)	1.000 (0.982–1.018)	0.980
Studies of North America	0.942 (0.840–1.056)	0.304
Witnessed cardiac arrest (%)	1.004 (0.999–1.009)	0.086
Bystander cardiopulmonary resuscitation (%)	0.998 (0.994–1.003)	0.471
Average response time (min)	0.971 (0.936–1.006)	0.105
Initial shockable rhythm (%)	1.000 (0.995–1.004)	0.911
Use of active compression-decompression resuscitation	1.072 (0.977–1.177)	0.143
Use of ITD	0.993 (0.888–1.111)	0.905
ITD attachment site (% of advanced airways use)	1.028 (1.005–1.051)	0.018

ITD = impedance threshold device.

and other factors was less examined. The current metaanalysis demonstrated that use of ACDR or ITD, in comparison with STR, seemed not to improve survival and neurologic outcomes of OHCA. Nonetheless, this minimal effect might be confounded by two important prognostic factors, that is, witnessed status and response time. The meta-regression uncovered that if these factors were controlled, ITD could still be effective. Use of an advanced airway also seemed to augment ITD effect.

Laboratory research has demonstrated that blood flow is significantly increased by ACDR compared with STR (31, 32). The minimal effect of ACDR in OHCA observed in the subgroup analysis was consistent with the Cochrane metaanalysis (5). There were several explanations for this discrepancy

between animal and clinical studies, such as the higher physical demand of ACDR in real practice (33) and the nonadherence to the chest of the ACDR device (17). The only study showing a survival benefit of ACDR was that by Plaisance et al (16), in which more staff were dispatched to perform ACDR, and with more experience in its use, as compared with other studies.

ITD improved hemodynamics and neurologic outcomes in animal studies (34–36). In comparison with ACDR, ITD is relatively easy to use in clinical settings. A previous metaanalysis (7) indicated that ITD was effective in improving short-term survival in OHCA patients, but a study (8) comparing ACDR/ITD with STR was also included in the analysis. In current subgroup analysis, ITD seemed not to improve either short-term or long-term outcome. Nonetheless, the meta-regression uncovered that this minimal effect might be caused by two important prognostic factors for OHCA, that is, witnessed status and response time.

The meta-regression showed that higher proportion of witnessed cardiac arrest and shorter response time were associated with higher log(RR) of ROSC, consistent with findings of other studies (37). Basically, these two factors similarly implied that time factor seemed to influence the ITD effect significantly. The use of ITD might be most effective within the circulatory phase (4–10 min following collapse) (38) when chest compression was addressed. Although the interaction term of ITD \times response time (< 6.5 min) was not statistically significant, the limited number of included RCTs might lead to this type II error. Some secondary studies (39, 40) of the Resuscitation Outcomes Consortium Prehospital Resuscitation Impedance Valve and Early Versus Delayed Analysis trial also pointed out that certain important CPR metrics, such as perishock pause and chest compression rates, had significantly influenced the survival of patients in this large trial. These results suggested that some subgroup of patient might still benefit from the use of ITD. More importantly, it has been noted that the response time or witnessed status is a summary of patient characteristics rather than a specific attribute of the trial. The finding is an aggregated association that cannot necessarily be extrapolated to individual patients. The individual patient-level extrapolation may need to be solved by an RCT.

Animal studies have shown the superiority of the drop in intrathoracic pressure and the increase in cerebral perfusion when ACDR and ITD were used in combination in comparison with STR, ACDR alone, or ITD alone (41, 42). Current subgroup analysis did not reveal the probable synergistic effect of ACDR/ITD. In the meta-regression, the interaction term of ACDR \times ITD was not statistically significant either. However, the limited number of RCTs studying ACDR/ITD combination might lead to this result.

The meta-regression indicated that advanced airway use might be synergistic with ITD. Most animal studies showing a survival benefit of ITD actually used ITD and ETT in combination rather than ITD with a facemask (34–36). The negative intrathoracic pressure might be better maintained by the occluded ventilation circuit with advanced airways. Plaisance et al (43) had demonstrated that the airway pressures with

TABLE 4. Results of Multivariate Meta-Regression Analysis on Log Relative Risk of Return of Spontaneous Circulation

Variable	Odds Ratio (95% CI)	p
Witnessed cardiac arrest (%)	1.017 (1.008–1.027)	< 0.001
Average response time (min)	0.936 (0.895–0.978)	0.004
Use of impedance threshold device	1.366 (1.120–1.665)	0.002

an ITD on either a facemask or ETT were similar in OHCA. Nonetheless, in their study (43), there was a trend showing that ITD with ETT could induce more negative pressure than ITD with a facemask, either in mean (-5.8 vs -3.4 mm Hg) or maximal pressure (-7.3 vs -4.6 mm Hg) (43).

Historically, effective use of facemask ventilation has been challenging even among skilled clinicians (44, 45). The single-handed “E-C” technique often tilts the facemask to the left, allowing air to leak under the right side of the mask (46). Plaisance et al (43) reduced this leak by using a strap around the head of the patient and by ventilating through a two-rescuer technique. Nonetheless, in most emergency services, the limited number of dispatched staff did not allow such a practice. In the subgroup analysis of the RCT by Aufderheide et al (9), there was also a nonsignificant trend in favor of ITD with ETT than with a supraglottic airway or a facemask in patients receiving STR. Nonetheless, there were only four RCTs included in this meta-regression model examining the effect of advanced airway use without other confounders controlled. Further studies should be conducted to examine this association.

In summary, despite that the pooled result indicated ACDR or ITD might not be effective in improving outcomes of OHCA patients, the meta-regression uncovered that time factor appeared to influence the ITD effect significantly. It might be difficult to shorten response time for every OHCA patient. Accordingly, this hypothesis might be examined in patients of in-hospital cardiac arrest, in whom the response time could probably be shorter and intubation could possibly be performed for every event although the etiology and confounding variables might be different from OHCA. ACDR has been examined in patients of in-hospital cardiac arrest; however, there are no similar studies for ITD (5).

Limitations

There was tremendous clinical and methodological heterogeneity among the included trials. Especially, the differences in EMS practice might make difficulties in shortening response time and limit the generalizability of this result. Prehospital intubation was not performed in most EMS cases. In some studies, prehospital intubation was viewed as a potentially harmful procedure for OHCA patients (47).

CONCLUSIONS

ACDR or ITD seemed not to improve ROSC in OHCA. This result might be confounded by two important prognostic factors, that is, witnessed status and response time. However, these hypotheses were based on aggregated association and might be better examined in future studies.

ACKNOWLEDGMENTS

We acknowledge statistical review provided by the Taiwan Clinical Trial Bioinformatics and Statistical Center, Training Center, and Pharmacogenomics Laboratory (which is

founded by National Research Program for Biopharmaceuticals at the National Science Council of Taiwan; NSC 102-2325-B-002-088).

REFERENCES

- Nichol G, Thomas E, Callaway CW, et al; Resuscitation Outcomes Consortium Investigators: Regional variation in out-of-hospital cardiac arrest incidence and outcome. *JAMA* 2008; 300:1423–1431
- Field JM, Hazinski MF, Sayre MR, et al: Part 1: Executive summary: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010; 122:S640–S656
- Andreka P, Frenneaux MP: Haemodynamics of cardiac arrest and resuscitation. *Curr Opin Crit Care* 2006; 12:198–203
- Chang MW, Coffeen P, Lurie KG, et al: Active compression-decompression CPR improves vital organ perfusion in a dog model of ventricular fibrillation. *Chest* 1994; 106:1250–1259
- Lafuente-Lafuente C, Melero-Bascones M: Active chest compression-decompression for cardiopulmonary resuscitation. *Cochrane Database Syst Rev* 2002; 3:CD002751
- Lurie K, Plaisance P, Sukhum P, et al: Mechanical advances in cardiopulmonary resuscitation. *Curr Opin Crit Care* 2001; 7:170–175
- Cabrini L, Beccaria P, Landoni G, et al: Impact of impedance threshold devices on cardiopulmonary resuscitation: A systematic review and meta-analysis of randomized controlled studies. *Crit Care Med* 2008; 36:1625–1632
- Wolcke BB, Mauer DK, Schoefmann MF, et al: Comparison of standard cardiopulmonary resuscitation versus the combination of active compression-decompression cardiopulmonary resuscitation and an inspiratory impedance threshold device for out-of-hospital cardiac arrest. *Circulation* 2003; 108:2201–2205
- Aufderheide TP, Nichol G, Rea TD, et al; Resuscitation Outcomes Consortium (ROC) Investigators: A trial of an impedance threshold device in out-of-hospital cardiac arrest. *N Engl J Med* 2011; 365:798–806
- Aufderheide TP, Frascone RJ, Wayne MA, et al: Standard cardiopulmonary resuscitation versus active compression-decompression cardiopulmonary resuscitation with augmentation of negative intrathoracic pressure for out-of-hospital cardiac arrest: A randomised trial. *Lancet* 2011; 377:301–311
- Moher D, Liberati A, Tetzlaff J, et al; PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Ann Intern Med* 2009; 151:264–269, W64
- Higgins JPT, Altman DG, Sterne JAC: Chapter 8: Assessing risk of bias in included studies. In: *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). Higgins JPT, Green S (Eds). The Cochrane Collaboration, 2011. Available at: <http://www.cochrane-handbook.org>. Accessed April 1, 2013
- DerSimonian R, Laird N: Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7:177–188
- Plaisance P, Adnet F, Vicaud E, et al: Benefit of active compression-decompression cardiopulmonary resuscitation as a prehospital advanced cardiac life support. A randomized multicenter study. *Circulation* 1997; 95:955–961
- Frascone RJ, Wayne MA, Swor RA, et al: Treatment of non-traumatic out-of-hospital cardiac arrest with active compression-decompression cardiopulmonary resuscitation plus an impedance threshold device. *Resuscitation* 2013; 84:1214–1222
- Plaisance P, Lurie KG, Vicaud E, et al: A comparison of standard cardiopulmonary resuscitation and active compression-decompression resuscitation for out-of-hospital cardiac arrest. French Active Compression-Decompression Cardiopulmonary Resuscitation Study Group. *N Engl J Med* 1999; 341:569–575
- Lurie KG, Shultz JJ, Callahan ML, et al: Evaluation of active compression-decompression CPR in victims of out-of-hospital cardiac arrest. *JAMA* 1994; 271:1405–1411
- Schwab TM, Callahan ML, Madsen CD, et al: A randomized clinical trial of active compression-decompression CPR vs standard

- CPR in out-of-hospital cardiac arrest in two cities. *JAMA* 1995; 273:1261–1268
19. Luiz T, Ellinger K, Denz C: Active compression-decompression cardiopulmonary resuscitation does not improve survival in patients with prehospital cardiac arrest in a physician-manned emergency medical system. *J Cardiothorac Vasc Anesth* 1996; 10:178–186
 20. Mauer D, Schneider T, Dick W, et al: Active compression-decompression resuscitation: A prospective, randomized study in a two-tiered EMS system with physicians in the field. *Resuscitation* 1996; 33:125–134
 21. Stiell IG, Hébert PC, Wells GA, et al: The Ontario trial of active compression-decompression cardiopulmonary resuscitation for in-hospital and prehospital cardiac arrest. *JAMA* 1996; 275:1417–1423
 22. Goralski M, Villéger JL, Cami G, et al: Evaluation of active compression-decompression cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *Réan Urg* 1998; 7:543–550
 23. Mauer D, Schneider T, Elich D, et al: Carbon dioxide levels during pre-hospital active compression-decompression versus standard cardiopulmonary resuscitation. *Resuscitation* 1998; 39:67–74
 24. Nolan J, Smith G, Evans R, et al: The United Kingdom pre-hospital study of active compression-decompression resuscitation. *Resuscitation* 1998; 37:119–125
 25. Skogvoll E, Wik L: Active compression-decompression cardiopulmonary resuscitation: A population-based, prospective randomised clinical trial in out-of-hospital cardiac arrest. *Resuscitation* 1999; 42:163–172
 26. Aufderheide TP, Pirralo RG, Provo TA, et al: Clinical evaluation of an inspiratory impedance threshold device during standard cardiopulmonary resuscitation in patients with out-of-hospital cardiac arrest. *Crit Care Med* 2005; 33:734–740
 27. Pirralo RG, Aufderheide TP, Provo TA, et al: Effect of an inspiratory impedance threshold device on hemodynamics during conventional manual cardiopulmonary resuscitation. *Resuscitation* 2005; 66:13–20
 28. Lai SM, Duncan PW: Stroke recovery profile and the Modified Rankin assessment. *Neuroepidemiology* 2001; 20:26–30
 29. Idris AH, Becker LB, Ornato JP, et al: Utstein-style guidelines for uniform reporting of laboratory CPR research. A statement for healthcare professionals from a Task Force of the American Heart Association, the American College of Emergency Physicians, the American College of Cardiology, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, the Institute of Critical Care Medicine, the Safar Center for Resuscitation Research, and the Society for Academic Emergency Medicine. *Resuscitation* 1996; 33:69–84
 30. Rittenberger JC, Raina K, Holm MB, et al: Association between Cerebral Performance Category, Modified Rankin Scale, and discharge disposition after cardiac arrest. *Resuscitation* 2011; 82:1036–1040
 31. Lindner KH, Pfenninger EG, Lurie KG, et al: Effects of active compression-decompression resuscitation on myocardial and cerebral blood flow in pigs. *Circulation* 1993; 88:1254–1263
 32. Prengel AW, Lindner KH, Pfenninger EG, et al: Effects of ventilation on hemodynamics and myocardial blood flow during active compression-decompression resuscitation in pigs. *Anesthesiology* 1996; 84:135–142
 33. Shultz JJ, Mianulli MJ, Gisch TM, et al: Comparison of exertion required to perform standard and active compression-decompression cardiopulmonary resuscitation. *Resuscitation* 1995; 29:23–31
 34. Pantazopoulos IN, Xanthos TT, Vlachos I, et al: Use of the impedance threshold device improves survival rate and neurological outcome in a swine model of asphyxial cardiac arrest. *Crit Care Med* 2012; 40:861–868
 35. Lurie KG, Zielinski T, McKnite S, et al: Use of an inspiratory impedance valve improves neurologically intact survival in a porcine model of ventricular fibrillation. *Circulation* 2002; 105:124–129
 36. Yannopoulos D, Sigurdsson G, McKnite S, et al: Reducing ventilation frequency combined with an inspiratory impedance device improves CPR efficiency in swine model of cardiac arrest. *Resuscitation* 2004; 61:75–82
 37. Abrams HC, Moyer PH, Dyer KS: A model of survival from out-of-hospital cardiac arrest using the Boston EMS arrest registry. *Resuscitation* 2011; 82:999–1003
 38. Weisfeldt ML, Becker LB: Resuscitation after cardiac arrest: A 3-phase time-sensitive model. *JAMA* 2002; 288:3035–3038
 39. Cheskes S, Schmicker RH, Verbeek PR, et al: Resuscitation Outcomes Consortium (ROC) Investigators: The impact of peri-shock pause on survival from out-of-hospital shockable cardiac arrest during the Resuscitation Outcomes Consortium PRIMED trial. *Resuscitation* 2014; 85:336–342
 40. Idris AH, Guffey D, Pepe PE, et al: Abstract 16132: Chest compression rates between 100 and 120 per minute are independently associated with increased adult survival following out-of-hospital cardiac arrest. *Circulation* 2012; 126:A16132
 41. Lanhelle A, Strømme T, Sunde K, et al: Inspiratory impedance threshold valve during CPR. *Resuscitation* 2002; 52:39–48
 42. Metzger AK, Herman M, McKnite S, et al: Improved cerebral perfusion pressures and 24-hr neurological survival in a porcine model of cardiac arrest with active compression-decompression cardiopulmonary resuscitation and augmentation of negative intrathoracic pressure. *Crit Care Med* 2012; 40:1851–1856
 43. Plaisance P, Soleil C, Lurie KG, et al: Use of an inspiratory impedance threshold device on a facemask and endotracheal tube to reduce intrathoracic pressures during the decompression phase of active compression-decompression cardiopulmonary resuscitation. *Crit Care Med* 2005; 33:990–994
 44. Kurola J, Harve H, Kettunen T, et al: Airway management in cardiac arrest—comparison of the laryngeal tube, tracheal intubation and bag-valve mask ventilation in emergency medical training. *Resuscitation* 2004; 61:149–153
 45. Elling R, Politis J: An evaluation of emergency medical technicians' ability to use manual ventilation devices. *Ann Emerg Med* 1983; 12:765–768
 46. Matic AA: The adult ergonomic face mask concept: Historical and theoretical perspectives. *J Clin Anesth* 2009; 21:300–304
 47. Hasegawa K, Hiraide A, Chang Y, et al: Association of prehospital advanced airway management with neurologic outcome and survival in patients with out-of-hospital cardiac arrest. *JAMA* 2013; 309:257–266

Copyright of Critical Care Medicine is the property of Lippincott Williams & Wilkins and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.