

# MATH 265 HW1

Hanzhang Yin

Aug/26/2024

## Question 1

*Proof.* Define a function  $f$  as:

$$f : \mathbb{N} \rightarrow \{1, 4, 7, 10, 13, \dots\} : n \mapsto 3n - 2$$

Now we need to show  $f$  is *one to one* and *onto*.

- Prove  $f$  is *one to one*

*Proof.* To prove  $f$  is *one to one*, we must show that if  $f(n_1) = f(n_2)$ , then  $n_1 = n_2$ .

Suppose  $f(n_1) = f(n_2)$ . Then:

$$\begin{aligned} 3n_1 - 2 &= 3n_2 - 2 \\ \Rightarrow 3n_1 &= 3n_2 \Rightarrow n_1 = n_2 \end{aligned}$$

Since  $n_1 = n_2$ ,  $f$  is injective. □

- Prove  $f$  is *onto*

*Proof.* To prove  $f$  is *onto*, we need to show for every  $m \in \{1, 4, 7, 10, 13, \dots\}$ , there exists an  $n \in \mathbb{N}$  s.t.  $f(n) = m$ .

Take any  $m \in \{1, 4, 7, 10, 13, \dots\}$ . We want to find  $n$  such that  $3n - 2 = m$ . Solving for  $n$ :

$$3n - 2 = m \Rightarrow 3n = m + 2 \Rightarrow n = \frac{m + 2}{3}$$

Since  $m$  is given of the form  $3k + 1$  for some integer  $k$ . Therefore,  $n \in \mathbb{N}$ , and  $f$  is onto. □

Since  $f$  is *one to one* and *onto*, it is a bijection. Hence,  $f(n) = 3n - 2$  is a explicit bijection from the set  $\mathbb{N}$  to  $\{1, 4, 7, 10, 13, \dots\}$ . □

## Question 2

*Proof.* Define a function  $f$  as:

$$f : (0, 1) \rightarrow \mathbb{R}_{>0} : n \mapsto \frac{n}{1-n}$$

Now we need to show  $f$  is *one to one* and *onto*.

- Prove  $f$  is *one to one*

*Proof.* To prove  $f$  is *one to one*, we must show that if  $f(n_1) = f(n_2)$ , then  $n_1 = n_2$ .

Suppose  $f(n_1) = f(n_2)$ . Then:

$$\begin{aligned}\frac{n_1}{1-n_1} &= \frac{n_2}{1-n_2} \\ \Rightarrow n_1(1-n_2) &= n_2(1-n_1) \\ \Rightarrow n_1 - n_1n_2 &= n_2 - n_2n_1 \\ \Rightarrow n_1 &= n_2\end{aligned}$$

Thus,  $f$  is one to one. □

- Prove  $f$  is *onto*

*Proof.* To prove  $f$  is *onto*, we need to show for every  $m \in \mathbb{R}_{>0}$ , there exists an  $n \in (0, 1)$  s.t.  $f(n) = m$ .

Take any  $m > 0$ . We want to find  $n$  such that  $\frac{n}{1-n} = m$ . Solving for  $n$ :

$$n = m(1-n) \Rightarrow n + nm = m \Rightarrow n(1+m) = m \Rightarrow n = \frac{m}{1+m}$$

Note that  $m > 0$ , this implies  $0 < \frac{m}{1+m} < 1$ , hence  $n \in (0, 1)$ .

Thus, for every  $m > 0$ , there is an  $n \in (0, 1)$  s.t.  $f(n) = m$ . (i.e.  $f$  is onto) □

Since  $f$  is *one to one* and *onto*, it is a bijection. Hence,  $f(n) = \frac{n}{1-n}$  is a explicit bijection from the set  $(0, 1)$  to  $\mathbb{R}_{>0}$ . □

### Question 3

*Proof.* Let  $x \in [0, 1)$  s.t.  $g(x) = w, w \in \mathbb{R}_{>0}$ , modified the function  $f$  from Q2 to  $g$  as:

$$g : [0, 1) \rightarrow \mathbb{R}_{>0} : g(x) = \begin{cases} \frac{\frac{1}{2}}{1 - \frac{1}{2}} = 1, & \text{if } x = 0 \\ \frac{\frac{2n+1}{2^{n+1}}}{1 - \frac{1}{2^{n+1}}} = \frac{1}{2^{n+1}-1}, & \text{if } x = \frac{1}{2^n}, n \in \mathbb{N}^+ \\ \frac{x}{1-x}, & \text{if } x \neq \frac{1}{2^n}, n \in \mathbb{N} \end{cases}$$

Now we need to show  $g$  is *one to one* and *onto*.

- Prove  $g$  is *one to one*

*Proof.* To prove  $g$  is *one to one*, we must show that if  $g(x_1) = g(x_2)$ , then  $x_1 = x_2$ .

**Case 1:**  $x_1 = 0$  and  $x_2 \neq 0$ :

- If  $x_1 = 0$ , then  $g(x_1) = 1$
- If  $x_2 \neq 0$ ,  $g(x_2)$  will either be in form  $\frac{2n+1}{1-2^{n+1}}$  or  $\frac{x_2}{1-x_2}$ , both are not equal to 1. Thus,  $g(x_1) \neq g(x_2)$ .

**Case 2:**  $x_1 = \frac{1}{2^n}, x_2 = \frac{1}{2^m}$  s.t.  $n \neq m$

- For  $x_1 = \frac{1}{2^n}$ , we have  $g(x_1) = \frac{2n+1}{1-\frac{1}{2^{n+1}}}$ .
- For  $x_2 = \frac{1}{2^m}$ , we have  $g(x_2) = \frac{2m+1}{1-\frac{1}{2^{m+1}}}$ .

Since  $n \neq m$ ,  $g(x_1) \neq g(x_2)$ .

**Case 3:**  $x_1 \neq \frac{1}{2^n}$  and  $x_2 \neq \frac{1}{2^m}$  with  $x_1 \neq x_2$

If  $x_1, x_2 \notin \{\frac{1}{2^n}\}$ , then  $g(x_1) = \frac{x_1}{1-x_1}$  and  $g(x_2) = \frac{x_2}{1-x_2}$ . Since  $x_1 \neq x_2$ ,  $\frac{x_1}{1-x_1} \neq \frac{x_2}{1-x_2}$ . Hence,  $g(x_1) \neq g(x_2)$ .

Combining all cases,  $g(x)$  is injective.  $\square$

- Prove  $g$  is *onto*

*Proof.* To prove  $g$  is *onto*, we need to show for every  $w \in \mathbb{R}_{>0}$ , there exists an  $x \in [0, 1)$  s.t.  $g(x) = w$ .

- For any  $w = 1$ , we have  $g(0) = 1$ .
- For any  $w = \frac{2n+1}{1-\frac{1}{2^{n+1}}}$ , there exists and  $x = \frac{1}{2^n}$  such that  $g(x) = w$ .
- For any other  $w > 0$ , there exists an  $x \in (0, 1) \setminus \{\frac{1}{2^n}\}$  s.t.  $g(x) = \frac{x}{1-x} = w$ , we can get:

$$x = \frac{w}{1+w}$$

Since  $w > 0, x \in (0, 1)$ .

Overall, for any  $w > 0$ , there is an  $x \in [0, 1]$  s.t.  $g(x) = w$ . Hence,  $g(x)$  is surjective.  $\square$

Since  $g$  is *one to one* and *onto*, it is a bijection.  $\square$

## Question 4

**(a) Show that  $g \circ f$  is injective if both of  $f$  and  $g$  are injective.**

*Proof.* Given that  $f : A \rightarrow B$  and  $g : B \rightarrow C$  are injective. We can write  $g \circ f$  as  $g \circ f : A \rightarrow C$ , defined as  $(g \circ f)(x) = g(f(x))$ .

Assume:  $(g \circ f)(x) = (g \circ f)(y)$ , for some  $x, y \in A$ ,

1. By the definition of composite function, we can get  $g(f(x)) = g(f(y))$
2. Given,  $g$  is injective,  $f(x) = f(y)$
3. Also given,  $f$  is injective,  $x = y$

Hence,  $g \circ f$  is injective.  $\square$

**(b) Show that  $g \circ f$  is surjective if both of  $f$  and  $g$  are surjective.**

*Proof.* Given that  $f : A \rightarrow B$  and  $g : B \rightarrow C$  are surjective. We can write  $g \circ f$  as  $g \circ f : A \rightarrow C$ , defined as  $(g \circ f)(x) = g(f(x))$ .

Let an arbitrary value  $z \in C$ . We need to find some  $x \in A$  such that  $(g \circ f)(x) = z$

1. Given  $g$  is surjective, hence there exists a  $y \in B$  such that  $g(y) = z$
2. Given  $f$  is surjective, hence there exists a  $x \in A$  such that  $f(x) = y$
3. Hence,  $(g \circ f)(x) = g(f(x)) = g(y) = z$

Hence,  $g \circ f$  is surjective.  $\square$