

MATH 240 HW1

Hanzhang Yin

Aug/26/2024

Question 1

Proof. Define a function f as:

$$f : \mathbb{N} \rightarrow \{1, 4, 7, 10, 13, \dots\} : n \mapsto 3n - 2$$

Now we need to show f is *one to one* and *onto*.

- Prove f is *one to one*

Proof. To prove f is *one to one*, we must show that if $f(n_1) = f(n_2)$, then $n_1 = n_2$.

Suppose $f(n_1) = f(n_2)$. Then:

$$\begin{aligned} 3n_1 - 2 &= 3n_2 - 2 \\ \Rightarrow 3n_1 &= 3n_2 \Rightarrow n_1 = n_2 \end{aligned}$$

Since $n_1 = n_2$, f is injective. □

- Prove f is *onto*

Proof. To prove f is *onto*, we need to show for every $m \in \{1, 4, 7, 10, 13, \dots\}$, there exists an $n \in \mathbb{N}$ s.t. $f(n) = m$.

Take any $m \in \{1, 4, 7, 10, 13, \dots\}$. We want to find n such that $3n - 2 = m$. Solving for n :

$$3n - 2 = m \Rightarrow 3n = m + 2 \Rightarrow n = \frac{m + 2}{3}$$

Since m is of the form $3k - 2$ for some integer k , and $n = k$ is an integer. Therefore, $n \in \mathbb{N}$, and f is onto. □

Since f is *one to one* and *onto*, it is a bijection. Hence, $f(n) = 3n - 2$ is a explicit bijection from the set \mathbb{N} to $\{1, 4, 7, 10, 13, \dots\}$. □

Question 2

Proof. Define a function f as:

$$f : (0, 1) \rightarrow \mathbb{R}_{>0} : n \mapsto \frac{n}{1-n}$$

Now we need to show f is *one to one* and *onto*.

- Prove f is *one to one*

Proof. To prove f is *one to one*, we must show that if $f(n_1) = f(n_2)$, then $n_1 = n_2$.

Suppose $f(n_1) = f(n_2)$. Then:

$$\begin{aligned}\frac{n_1}{1-n_1} &= \frac{n_2}{1-n_2} \\ \Rightarrow x_1(1-x_2) &= x_2(1-x_1) \\ \Rightarrow x_1 - x_1x_2 &= x_2 - x_2x_1 \\ \Rightarrow x_1 &= x_2\end{aligned}$$

Thus, f is one to one. □

- Prove f is *onto*

Proof. To prove f is *onto*, we need to show for every $m \in \mathbb{R}_{>0}$, there exists an $n \in (0, 1)$ s.t. $f(n) = m$.

Take any $m > 0$. We want to find n such that $\frac{n}{1-n} = m$. Solving for n :

$$n = m(1-n) \Rightarrow n + nm = m \Rightarrow n(1+m) = m \Rightarrow n = \frac{m}{1+m}$$

Note that $m > 0$, this implies $0 < \frac{m}{1+m} < 1$, hence $n \in (0, 1)$.

Thus, for every $m > 0$, there is an $n \in (0, 1)$ s.t. $f(n) = m$. (i.e. f is onto) □

Since f is *one to one* and *onto*, it is a bijection. Hence, $f(n) = \frac{n}{1-n}$ is a explicit bijection from the set $(0, 1)$ to $\mathbb{R}_{>0}$. □

Question 3

Proof. Modified the function f from Q2 to g as:

$$g : [0, 1) \rightarrow \mathbb{R}_{>0} : g(x) = \begin{cases} \frac{x}{1-x}, & \text{if } x \in (0, 1) \\ a, & \text{if } x = 0 \end{cases}$$

Where $a \in \mathbb{R}_+$ and is not in the range of $f(x)$ for $x \in (0, 1)$.

Now we need to show g is *one to one* and *onto*.

- Prove g is *one to one*

Proof. To prove g is *one to one*, we must show that if $g(x_1) = g(x_2)$, then $x_1 = x_2$.

- If $x_1, x_2 \in (0, 1)$, then from Q2, $\frac{x_1}{1-x_1} = \frac{x_2}{1-x_2}$ implies $x_1 = x_2$
- If $x_1 = 0$ and $x_2 \in (0, 1)$, then $g(0) = a, g(x_2) = \frac{x_2}{1-x_2}$, but $a \neq \frac{x_2}{1-x_2}, \forall x_2 \in (0, 1)$. Thus, $g(x_1) \neq g(x_2)$ and it is one to one.

□

- Prove g is *onto*

Proof. To prove g is *onto*, we need to show for every $y \in \mathbb{R}_{>0}$, there exists an $x \in (0, 1)$ s.t. $g(x) = y$.

- For $y \neq a$, there exists an $x \in (0, 1)$ s.t. $\frac{x}{1-x} = y$.
- For $y = a$, we have $g(0) = a$.

Hence, $g(x)$ covers \mathbb{R}_+ , which is onto.

□

Since g is *one to one* and *onto*, it is a bijection.

□

Question 4

(a) Show that $g \circ f$ is injective if both of f and g are injective.

Proof. Given that $f : A \rightarrow B$ and $g : B \rightarrow C$ are injective. We can write $g \circ f$ as $g \circ f : A \rightarrow C$, defined as $(g \circ f)(x) = g(f(x))$.

Assume: $(g \circ f)(x) = (g \circ f)(y)$, for some $x, y \in A$,

1. By the definition of composite function, we can get $g(f(x)) = g(f(y))$
2. Given, g is injective, $f(x) = f(y)$
3. Also given, f is injective, $x = y$

Hence, $g \circ f$ is injective.

□

(b) Show that $g \circ f$ is surjective if both of f and g are surjective.

Proof. Given that $f : A \rightarrow B$ and $g : B \rightarrow C$ are surjective. We can write $g \circ f$ as $g \circ f : A \rightarrow C$, defined as $(g \circ f)(x) = g(f(x))$.

Let an arbitrary value $z \in C$. We need to find some $x \in A$ such that $(g \circ f)(x) = z$

1. Given g is surjective, hence there exists a $y \in B$ such that $g(y) = z$
2. Given f is surjective, hence there exists a $x \in A$ such that $f(x) = y$
3. Hence, $(g \circ f)(x) = g(f(x)) = g(y) = z$

Hence, $g \circ f$ is surjective. □