On Existence of Digital Measure Constant

F. G. Ersezer, *Student Member, IEEE*

***Abstract*—Unidentified continuous functions are a generalized classification for all signals. The author assumes that any unidentified continuous function (any signal) is either digital (non-Archimedean continuum) or analog (Archimedean continuum), and gives a method to indicate a distinction between both continuum models. The distinction is measured by the digital measure constant Hyp_1, the area difference between the two models. In the non-Archimedean model with infinitesimal step dx, we show that Hyp_1 is a nonzero infinitesimal for every non-constant function, that it vanishes at large scale, and that Hyp_1/dx equals one half of the total variation of the function. The factor one half is independent of scale and of the function. The method can therefore be applied to all unidentified continuous functions in order to identify their continuum models.**

***Index Terms*—Time-frequency analysis, non-standard analysis, digital topology**

I. Introduction

THE article "Approaches to analysis with infinitesimals following Robinson, Nelson and others" [18] has the section "The CD+II mindset". The terms of that section will be regarded, since they are foundations of set-theoretical identifications that are based on an "intuitively equal" algebraic partition (arithmetization) of the real number line (all real numbers) into a countably infinite amount of interval groups. (The work of Sophus Lie is a general reference for the historical viewpoint of continuous symmetric groups [2]–[4].)

"The Cantor–Dedekind postulate (CD) identifying the line in physical space with the real number line" (Cantor's main work is [5]; his arguments on reals and infinitesimals are on pages 26–55 of [17]).

"The Intended Interpretation hypothesis (II), entailing an identification of a standard $\mathbb{N}$ in its set-theoretic context, on the one hand, with ordinary intuitive counting numbers, on the other."

These main arguments will be the building blocks for the separation of unidentified continuous functions (generalized signals). The mathematical foundations are generally regarded whenever there is a need for deductive (Fourier or harmonic) functional analysis (see [20]). The author will regard non-standard analysis, since the main arguments for a digital continuum can be founded via inductive (instantaneous) non-standard notions of analysis. Rather than concerning applications about the statistical large-scale construction of signals on the real numbers to generate a generalized theory, the author establishes a deterministic model of small-scale construction for signals on $\mathbb{R}^*$ as a generalized theory and applies it to larger scales.




Fig. 1. Cartesian Coordinate System

The Cartesian Coordinate System is mainly a product of the algebraic partition of the real number line into "intuitively equal", countable and infinitely many interval parts. The branches of mathematics, mainly set theory, measure theory and analysis, define different terms regarding this application of partition. The author adds another term, the "Coarsity of Compression Model Hypothesis" (CCMH). This hypothesis is the assumption that all (a priori) proportionately partitioned parts can always be partitioned proportionately again, as a generalization. In order to explain the CCMH, the author will regard types of continuum and the general view of scales as perspectives. Finally, the author will define the digital measure constant as a continuum relation.

While the Cartesian Coordinate System is the human mind's intuition to partition physical space into patterns in order to recognize physical space itself, the term "intuitively equal" is used rather than "equal" because the real number set (the real number line) is uncountable in terms of counting numbers. Via non-standard analysis one can construct the same a priori idea of partition on the unit interval, and then apply this idea of partition to unidentified continuous functions. Work of this type has been done before in Physics Letters A by R. F. Werner and M. P. H. Wolff [21].

A. Types of Continuum for Unidentified Continuity

We begin with the idea of certain graphs (images) which look identical when looked at from far away. Since a "function" is the

Revised version, 31 October 2026. This version keeps the original argument and states its results in precise form: the value of Hyp_1, the unit relation behind Hyp_0, and the logical energy unit u.

representation of an image on a graph, the distinctions between small-scale models of *"functions that look identical from far away"* will be separated into two kinds of continuum. This analysis was made in the article *"Tension between Intuitive Infinitesimals and Formal Mathematical Analysis"* [19] by Mikhail G. Katz and David Tall. The author mainly changed Luzin's saw (sawtooth) into ceil/floor series as a generalization.




Fig. 2. Analog Continuum Model as x




Fig. 3. Digital Continuum Model as $\lceil x \rceil$

In Figures 2–3, the two graphs (images) of functions look identical when looked at from a far distance. For the generalization, the term "unidentified" will be used for functions f (see below) that look like generalizations of an infinitely differentiable function (n indicates infinite) when looked at from far away (large scale), but are "unknown" in terms of continuum (small scale):

$$\frac{\partial^n f}{\partial x^n} \cong 0\,; \qquad f(x) \cong \sum_{i\in\mathbb{N}} b_i \cdot x^i\,; \quad b_i \in \mathbb{R}$$

Here and below, $A \cong B$ means that $A - B$ is infinitesimal, that is, A and B are equal when looked at from large scale.

Any area below the curve of the two possible continuum models of an "unidentified" function will also look like the same constant number when looked at from far away. But in reality there will be an area difference between the defined continuum models.

The reason why ceil/floor series are used for the digital (non-Archimedean) continuum model, rather than the functional representations described in *"Tension between Intuitive Infinitesimals and Formal Mathematical Analysis"* at pages 14–15, is that the *"Natural Number Set"* representation as *"intuitively equal"* interval partitions can be shown with ceil/floor series.

II. Generalized Functional Model Interpretations

The abstraction is for the functional integration of natural numbers (as groups) with real numbers. Any continuum model of an "unidentified" function f at small scale is therefore shown with these functional representations. For a step $dx > 0$, write the ceil and floor at that step as

$$\lceil x \rceil_{dx} = dx \cdot \left\lceil \frac{x}{dx} \right\rceil, \qquad \lfloor x \rfloor_{dx} = dx \cdot \left\lfloor \frac{x}{dx} \right\rfloor$$

The author will regard the (Riemann summation [1]) upper summation model as the difference between the ceil and the analytical function, and the lower summation model as the difference between the analytical and the floor function. The constant Hyp_1 is the representation of the area difference between the models. The coefficients a_i are natural numbers here, so f is increasing for $x \geq 0$ and the ceil and floor models are exactly the upper and lower summation models:

$$\int_0^H \left(\sum_{i\in\mathbb{N}} a_i \lceil x \rceil_{dx}^i - \sum_{i\in\mathbb{N}} a_i x^i \right) dx \;\cong\; Hyp_1$$

$$\int_0^H \left(\sum_{i\in\mathbb{N}} a_i x^i - \sum_{i\in\mathbb{N}} a_i \lfloor x \rfloor_{dx}^i \right) dx \;\cong\; Hyp_1$$

Both equations hold together up to the relation $\cong$: their two sides differ only by an infinitesimal of higher order than Hyp_1 itself (Proposition 1).

Note: The integration is taken on the domain $[0, H]$ because the generalized functional model is applied from a small-scale perspective. Any definite integration of length, area or volume that results in a constant number from the large-scale perspective is assumed to be "potentially infinite" in small-scale models. Here H is such a potential infinity: an infinite hyperreal bound, so that the integral is a well-defined element of $\mathbb{R}^*$. Hence the representation of the digital continuum is possible regardless of the basis domain. The term "actual infinite" (∞) will be used as the "singular infinite", and "potential infinite" (H) as a perspective.

III. Model Interpretations on Small Scale

The abstraction is to complement the smaller-scale perspective. The symbol $(dx)_{\|\mu\|}$ will indicate the digital (non-Archimedean) continuum at small-scale perspective. The digital continuum $(dx)_{\|\mu\|}$ has an interrelation with ceil/floor series as a functional

representation. This is mainly indicated via the hyperreal set, shown with the symbol $\mathbb{R}^*$. (Here the author must cite Hilbert [8]–[10], Hausdorff [6], [7], Lebesgue [11], Borel [12], [13] and Robinson [15], because of various foundations and relations in analysis, set theory and measures.)

$$0 < a < +\infty\,; \qquad a = n \cdot (dx)_{\|\mu\|}$$

$$a \in \mathbb{R}^*\,; \quad n \in \mathbb{N}^+\,; \qquad \frac{\pm 1}{(dx)_{\|\mu\|}} \triangleq \pm\infty$$

Here the non-integer coefficient variants of $(dx)_{\|\mu\|}$ unit/group lengths are non-definitive for the $(dx)_{\|\mu\|}$ units/groups. So the CCMH is basically an "Intended Interpretation" model on the unit interval.




Fig. 4. A proportionate partitioning

IV. Digital Measure Constant

On earlier pages the statement *"Any area below the curve of the two possible continuum models of an unidentified function will also look like the same constant number when looked at from far away. But in reality there will be an area difference between the defined continuum models"* was given. The author will regard the type of "unidentified" function f with the term

$$\Delta x \cdot \int_{\Omega_1} f$$

for functions that are variations of analytical models, which have an Archimedean continuum at small scale. The author will also regard the type of "unidentified" function f from a closer view with the term

$$\int_{\Omega_1} f\,dx$$

for functions that are variations of synthetical models, which have a digital (non-Archimedean) continuum at small scale. Precisely, $\Delta x \cdot \int_{\Omega_1} f$ is the integral of the analytical model, and $\int_{\Omega_1} f\,dx$ is the step sum of the synthetical model, $\sum_k f(\xi_k)\,dx$, taken over cells of length dx with $f(\xi_k)$ the upper (or lower) value of f on each cell.

Since any finite area below the curve of an "unidentified" function in the large-scale view looks non-finite when looked at from the small-scale view, the area difference between the models can be shown as nearly equal to zero in the large-scale view.

If on $\Omega_1 := [a,b]$ the large-scale integration of f is known to be a finite constant S,




Fig. 5. Area under the curve of unidentified function

then the equations below must hold for the large-scale view:

$$\Delta x \cdot \int_{\Omega_1} f \cong S\,, \qquad \int_{\Omega_1} f\,dx \cong S\,, \qquad S \in \mathbb{R}$$

$$\left|\Delta x \cdot \int_{\Omega_1} f - \int_{\Omega_1} f\,dx\right| \cong 0$$

Note: If the domain or range of the function is extended to infinity anyhow, it must be extended to infinity regardless of all perspectives. Here the functional representation of "proportion" is entirely due to the perspective; therefore a potential point resides *at some infinity, not "the infinity" itself* [14]. This is due to Carathéodory's extension theorem (lemma):

$$\mu^*(S) = \mu^*(S \cap M) + \mu^*(S \cap M^c)$$

However, for the case where on the defined domain $\Omega_1 := [a,b]$ the integration of f is known to be non-finite also at large scale,

$$\Delta x \cdot \int_{\Omega_1} f = \infty \quad \text{and} \quad \int_{\Omega_1} f\,dx = \infty$$

Hence for this condition, the difference between the models must result in the digital measure constant, since the extension of an infinite range is singular for all perspectives:

$$\left|\Delta x \cdot \int_{\Omega_1} f - \int_{\Omega_1} f\,dx\right| \cong Hyp_1$$

Hyp_1 is a constant because it determines the measurability of definite integrals. It cannot be equal to zero, since that would mean the two continuum models are equal. And it must be a constant, since if the integration of the two possible continuum models did not hold as a constant, the integration of definite (finite) integrals (in the large-scale view) would give different results for the two models because of the expansion of the compared maps (graphs, images), which would contradict large-scale definite integration (Gromov [16]). The value of this constant is given precisely as follows.

Proposition 1: Let f be continuously differentiable on $\Omega_1 = [a,b]$, let dx be the step, and let $V_a^b(f) = \int_a^b |f'(x)|\,dx$ be the total variation of f. Then, for both the upper and the lower summation model,

$$Hyp_1 = \left|\Delta x \cdot \int_{\Omega_1} f - \int_{\Omega_1} f\,dx\right| = \frac{1}{2}\,dx \cdot V_a^b(f) + o(dx)$$

For monotone f, $V_a^b(f) = |f(b) - f(a)|$.

Proof: On a cell of length dx, the upper value of f exceeds $f(x)$ by approximately $|f'|$ times the distance from x to the point where the upper value is attained. That distance averages $dx/2$ over the cell, so the cell contributes $\frac{1}{2}|f'|\,dx^2$. Summing over the cells gives $\frac{1}{2}\,dx \int_a^b |f'|\,dx$. The lower model is symmetric. For monotone f the leading term is the first term of the Euler–Maclaurin formula. ■

Proposition 1 gives the claimed properties of Hyp_1 exactly:

1) *Non-zero:* $Hyp_1 = 0$ only when $V_a^b(f) = 0$, that is, when f is constant and the two continuum models coincide. For every other function, Hyp_1 is a nonzero infinitesimal.
2) *Zero at large scale:* $Hyp_1 \cong 0$, since it is a multiple of the infinitesimal dx.
3) *Scale invariant:* $Hyp_1/dx = \frac{1}{2}V_a^b(f)$ does not depend on the step dx. The factor $\frac{1}{2}$ is universal: it is the same at every scale and for every function.
4) *Ceil and floor agree:* the two equations of Section II give the same leading term $\frac{1}{2}\,dx \cdot V_a^b(f)$, and differ only at order dx^2.

Start by expanding the generalized equation into

$$\left|\Delta x \cdot \int_0^{\infty} f - \int_0^{\infty} f \cdot dx\right| = Hyp_1$$

There is a special case when f is dx. This is because dx acts as a *constant* on $\mathbb{R}$, because $\mathbb{R}$ is *Archimedean*, and acts as an *infinitesimal* on $\mathbb{R}^*$, because it is *non-Archimedean*:

$$f(x) \cong dx$$

The generalized equation therefore has to abide by two situations. The first is when f acts as the *constant number* $dx > 0$, with the actual infinity as upper bound:

$$\left|\Delta x \cdot \int_0^{\infty} dx - \int_0^{\infty} dx \cdot dx\right| = Hyp_1$$

$$\int_0^{\infty} dx \cdot dx = dx \cdot \int_0^{\infty} dx = \infty$$

The equations above hold since the large-scale integration of this integral is infinite. The second is when f acts as the *infinitesimal number* dx. The upper bound is then the potential infinity of the small-scale perspective, $H = 1/dx$:

$$\frac{\pm 1}{(dx)_{\|\mu\|}} \triangleq \pm\infty$$

$$\int_0^H dx = \int_0^{1/dx} dx = \frac{1}{dx} = 1 + 1 + 1 + \ldots$$

where the sum has $1/dx$ terms. Therefore

$$\int_0^H dx \cdot dx = dx \cdot \int_0^{1/dx} dx = 1$$

The two situations use different upper bounds, the actual infinity ∞ and the potential infinity $H = 1/dx$, which is why they give different values. This special case does not measure the area difference of Proposition 1: for the constant function dx the two continuum models coincide, so that difference is zero. Instead it compares the two perspectives. In the analytical perspective one unit of length is Δx; in the small-scale perspective the same unit is $dx \cdot (1/dx)$ steps. Their ratio defines the second constant:

$$\frac{\Delta x}{dx} = Hyp_0$$

that is,

$$\Delta x = Hyp_0 \cdot dx$$

Meaning: any unit distance Δx on $\mathbb{R}$ contains Hyp_0 steps of the digital continuum $\mathbb{R}^*$. Hyp_0 is the conversion factor between the two units, like the number of centimetres in a metre. With $\Delta x = 1$, $Hyp_0 = 1/dx$ is an infinite hypernatural number. In order to obtain the values of the digital measure constants Hyp_0 and Hyp_1, the author indicates that the generalized functional model interpretations of Section II lead to the result: the area difference between ceil/floor and analytic functions with natural coefficients is the Hyp_1 constant, whose value is given by Proposition 1.

V. Digital Measure

Any function that represents a generalized signal and has unidentified continuity shown on the real numbers $\mathbb{R}$ (far-away view) can also be shown on the hyperreal number set $\mathbb{R}^*$. The author indicates the existence of another digital measure constant, Hyp_0, via generalization, in order to show that a point at real coordinate x lies at digital coordinate (step count) x^* with

$$x^* = Hyp_0 \cdot x\,; \qquad x \in \mathbb{R}\,; \quad x^* \in \mathbb{R}^*$$

The digital measure of an interval $[a,b]$ on $\mathbb{R}$ is its number of steps:

$$\mu_D[a,b] = Hyp_0 \cdot (b-a)\,; \qquad b > a$$

Equivalently, an interval $[a^*,b^*]$ given in digital coordinates has real length

$$\frac{1}{Hyp_0} \cdot (b^* - a^*)$$

VI. Applications

If there is a signal f (shown via a function) that is known to have unidentified continuity,

$$\frac{\partial^n f}{\partial x^n} \cong 0\,; \qquad f(x) \cong \sum_{i \in \mathbb{N}} b_i \cdot x^i\,; \quad b_i \in \mathbb{R}$$

then via this signal one can obtain the digital continuum model or the analog continuum model of this signal. So any unidentified continuous function f can be transferred from one continuum model to the other by

$$f(x) = f\left(\frac{x^*}{Hyp_0}\right) ; \qquad x \in \mathbb{R} ; \quad x^* \in \mathbb{R}^*$$

An example is the energy of the photon (wave–particle), shown via functions ($E(x)$, $f(x)$ and $\lambda(x)$ for the functional representations of energy, frequency and wavelength) that have unidentified continuity:

$$E(x) = h \cdot f(x) ; \qquad f(x) = \frac{c}{\lambda(x)} ; \qquad \lambda(x) = x$$

Then the digital continuum model of wavelength (as an interval) is

$$\lambda = \frac{\lambda^*}{Hyp_0} ; \qquad \lambda^* \in \mathbb{R}^* ; \quad \lambda \in \mathbb{R}$$

with the bounded range

$$\lambda^* \in \mathbb{R}^* ; \qquad 0 \leq \lambda^* \leq \frac{1}{dx}$$

If every wavelength is an integer multiple of the interval $[0, dx]$, then $\lambda^* = n$ with $n \in \mathbb{N}^+$ and $n \leq N$, where $N = 1/dx$. The energies are then

$$E_n = h \cdot \frac{c}{\lambda} = \frac{h\,c\,Hyp_0}{n} ; \qquad n = 1, 2, \ldots, N$$

These energy levels are not integer multiples of each other; they form the sequence $E_1, E_1/2, E_1/3, \ldots$ Nevertheless, a single *logical* unit u exists such that every level is an integer multiple of it. Let $L = \mathrm{lcm}(1, 2, \ldots, N)$, which exists as a hypernatural number by transfer. Then

$$u = \frac{h\,c\,Hyp_0}{L} ; \qquad E_n = \frac{L}{n} \cdot u , \quad \frac{L}{n} \in \mathbb{N}^+$$

For example, for a finite range with $N = 10$, $L = 2520$ and the levels are $2520u, 1260u, 840u, \ldots$ The bounded range of λ^* is essential: without an upper bound on n, no single unit divides every level. The author therefore indicates that if every wavelength in a bounded range is an integer multiple of the interval $[0, dx]$, then every energy level is an integer multiple of the logical constant u. This holds within the digital continuum model and is not a statement about measured photon spectra.

VII. Conclusion

Within the non-Archimedean continuum model, the digital measure constant exists: Hyp_1 is a nonzero infinitesimal for every non-constant function, it vanishes at large scale, and $Hyp_1/dx = \frac{1}{2}V_a^b(f)$ is independent of the scale, with the universal factor $\frac{1}{2}$. The constant $Hyp_0 = \Delta x/dx$ converts between the real and the digital units of length.

Data Availability Statement

Some of the data is available to the public, while some of the data has prerequisites, and some of the data is located at an outdated website. The author has given information about the contents he has read below. All referenced pages are at the equivalent of the internet domain's reference pages (logical), not the original book's or reference's pages (physical).

References and Footnotes

[1] Riemann, Bernhard, *Gesammelte mathematische Werke und wissenschaftlicher Nachlass*, pages: (18–58), (78–151), (152–159), (228–265), (269–283), (298–330), (346–359), (372–419), (448), (464–470). https://archive.org/details/bernardrgesamm00riemrich/page/n17/mode/2up?view=theater

[2] Lie, Sophus, *Theorie der Transformationsgruppen*, Reihe 1, Seite: (III–VIII), (1–8), (11–136), (171–183), (184–211), (289–310), (554–597). https://gdz.sub.uni-goettingen.de/id/PPN585383391

[3] Lie, Sophus, *Theorie der Transformationsgruppen*, Reihe 2, Seite: (35–44), (77–114), (171–176), (178–206), (292–293), (298–327). https://gdz.sub.uni-goettingen.de/id/PPN588388580

[4] Lie, Sophus, *Theorie der Transformationsgruppen*, Reihe 3, Seite: (IX–XXV), (1–109), (122–170), (180–232), (266–282), (296–398), 427, (437–471), (485–497), (524–635), (665–753). https://gdz.sub.uni-goettingen.de/id/PPN588388718

[5] Cantor, Georg, *Gesammelte Abhandlungen* (ed. Ernst Zermelo), Seite 115 (§2). https://gdz.sub.uni-goettingen.de/id/PPN237853094

[6] Hausdorff, Felix, *Grundzüge der Mengenlehre*: (73), (101–139), (158–161), (185–189), (209–219), (229–239), (247–249), (253–335), (369–398), (403–417), (430–448). https://gallica.bnf.fr/ark:/12148/bpt6k9736980b

[7] Hausdorff, Felix, *Gesammelte Abhandlungen, Band 3: Deskriptive Mengenlehre und Topologie*, U. Felgner, H. Herrlich, M. Hušek, V. Kanovei, P. Koepke, G. Preuß, W. Purkert, E. Scholz (eds.), Springer: (27–62), (91–341), (374–430), (468–472), (483–486), 494–497, (543–548), 559, (725–745), (757–776), (785–802), (805–832), (847–899), (909–987).

[8] Hilbert, David, *Gesammelte Abhandlungen*, Band I, Seite: (5–23), (26–42), (53–68), (69–92), (112–149), (176–242), (275–279), (291–302), (390–393), (412–427), (507–509). https://gdz.sub.uni-goettingen.de/id/PPN237821559

[9] Hilbert, David, *Gesammelte Abhandlungen*, Band II, Seite: (1–24), (34–37), (102–147), (162–263), (287–319), (341–344), (376–414). https://gdz.sub.uni-goettingen.de/id/PPN237833719

[10] Hilbert, David, *Gesammelte Abhandlungen*, Band III, Seite: (1–25), (33–37), (43–80), (94–145), (290–377). https://gdz.sub.uni-goettingen.de/id/PPN237834022

[11] Lebesgue, Henri, *Leçons sur l'intégration et la recherche des fonctions primitives*, pages: (1–14), (17–35), (37–140), (156–184), (202–226), (262–271), (314–340). https://gallica.bnf.fr/ark:/12148/bpt6k377935x/f359.item

[12] Borel, Émile, *Leçons sur la théorie des fonctions*, pages: (20–81). https:

//archive.org/details/leonssurlathori00boregoog/page/n19/mode/2up

[13] Borel, Émile, *Leçons sur les séries divergentes*, pages: (83–113), (147–163). https://archive.org/details/leonssurlessrie01boregoog/page/n114/mode/2up

[14] Carathéodory, Constantin, *Mass und Integral und ihre Algebraisierung*, Birkhäuser, Seite: (10–172), (192–200), (247–281), (285–287), (294–310).

[15] Robinson, Abraham, *Non-Standard Analysis* (first edition), pages: (24–64), (65–118), (132–149), (189–192), (193–245).

[16] Gromov, Michael, "Groups of polynomial growth and expanding maps (with an appendix by Jacques Tits)," *Publications Mathématiques de l'IHÉS*, Tome 53 (1981), pp. 53–78. http://www.numdam.org/item/PMIHES_1981__53__53_0

[17] Ehrlich, Philip, "The rise of non-Archimedean mathematics and the roots of a misconception I: the emergence of non-Archimedean systems of magnitudes," pages: (16), (26–55). https://link.springer.com/article/10.1007/s00407-005-0102-4

[18] Katz, Mikhail G., Hrbacek, Karel, Kanovei, Vladimir, Fletcher, Peter, Lobry, Claude, Sanders, Sam, "Approaches to analysis with infinitesimals following Robinson, Nelson and others," page: (4). https://arxiv.org/abs/1703.00425

[19] Katz, Mikhail G., Tall, David, "Tension between intuitive infinitesimals and formal mathematical analysis," pages: (14–15). https://arxiv.org/abs/1110.5747

[20] Soto-Bajo, Moisés, Fraguela Collar, Andrés, Herrera-Vega, Javier, "On the concept of frequency in signal processing: a discussion [Perspectives]." https://ieeexplore.ieee.org/document/10243508

[21] Werner, R. F., Wolff, M. P. H., "Classical mechanics as quantum mechanics with infinitesimal $\hbar$," *Physics Letters A*. https://www.sciencedirect.com/science/article/abs/pii/0375960195003443