

London Air Quality Pre and Post ULEZ

...

By Ioana and Mina

Changes to levels of PM₁₀ and Nitrogen Dioxide in Central London before and after the introduction of the Ultra-Low Emissions Zone (ULEZ)

What is NO_2 and PM_{10} ?

- Nitrogen Dioxide (NO_2) is produced by burning of **fuel**, especially **diesel-run engines**
 - Causes irritation in respiratory system
 - Particles smaller than about 10 micrometers or PM_{10} (**dust, rubber** and **metal** from engine wear)
 - Can settle in the airways and lungs, worsening heart and lung diseases
-

The Research



Our hypotheses are:

1. The introduction of ULEZ decreased the amount of NO_x in Zone 1 of Central London by an effect size of at least 0.3. Null hypothesis - ULEZ did not reduced the amount of NO_x in Zone 1 by any significant effect size
2. The introduction of ULEZ was at least 1.25 as effective than the introduction of LEZ in reducing NO₂ levels in Central London. Null hypothesis - The effect of ULEZ on NO₂ levels was not significantly more effective than LEZ in reducing NO₂ levels in Central London.
3. The introduction of ULEZ decreased the amount of PM₁₀ in Zone 1 of Central London by an effect size of at least 0.4. Null hypothesis - ULEZ did not reduce the amount of PM₁₀ in Zone 1 by any significant amount.
4. The introduction of ULEZ was at least 1.5 as effective as the introduction of LEZ in reducing PM₁₀ levels in Central London. Null hypothesis - The effect of ULEZ on PM₁₀ was not significantly more effective than LEZ in reducing PM₁₀ levels in Central London.

Methodology and data

Took King's College London Air quality data from the following sites for the following pollutants:

NO₂

CT3	City of London, Sir John Cass School
KC1	Kensington and Chelsea, North Kensington
HK6	Hackney, Old Street
CT6	City of London, Walbrook Wharf
WM6	Westminster, Oxford Street

PM₁₀


CT3	City of London, Sir John Cass School
CT8	City of London, Upper Thames Street
MY7	Westminster Marylebone Road FDMS
HK6	Hackney, Old Street
WM6	Westminster, Oxford Street



Methodology and data

- Data by King's College London (London Air Quality Network, LAQN)
- Sampled data for two months of years 2006, 2007, 2018, 2019 to get normal distributions
- Welch T-test
- Final outcome: Cohen's d effect size
- Alpha threshold set to 0.05






NO2 Values


**Pre LEZ
(2007)**

**And
Post LEZ
(2008)**

A




B




A. NO2 Levels
Pre- and Post LEZ

B. NO2 Levels
Pre- and Post ULEZ

C




D



C. PM10 Level
Pre- and Post LEZ


D. PM10 Levels
Pre- and Post ULEZ




NO₂
Samples pre- and
post-ULEZ
Cohen's d = -3.657



PM₁₀ samples pre-
and post-ULEZ
Cohen's d = -7.025



NO₂ samples
before and after
ULEZ and LEZ
Ratio of Cohen's
d's : 0.848



PM₁₀ samples
before and
after ULEZ and
LEZ
Ratio of
Cohen's d's :
3.66

Going forward:


Limitation of our results:

- Smaller population of data for post-ULEZ
- Unevenly distributed data for the Central London sensors

Next steps:


- Repeating sampling and analysis for other pollutants ($\text{PM}_{2.5}$ and CO)
- Choosing Inner & Outer London sites for testing impact of LEZ

London's air pollution before ULEZ



Reference: LAEI, 2016

And in 2025, after ULEZ and the Mayor's other actions



Reference: London Environment Strategy, 2018

Conclusion

Our research shows that ULEZ was effective in reducing NO_2 and PM_{10} levels by a significant margin.

Interestingly, ULEZ was significantly more effective in reducing PM_{10} than LEZ was, whereas LEZ had a moderately larger impact on NO_2 than ULEZ.

Appendix

NO2 Level (ULEZ)

Based on the p value of 0.0 and our alpha of 0.05 we reject the null hypothesis.

Due to these results, we can state that there is a difference between our samples with an effect size, Cohen's d, of -3.469 and power of 1.0.

PM10 Level (ULEZ)

Based on the p value of 0.0 and our alpha of 0.05 we reject the null hypothesis.

Due to these results, we can state that there is a difference between our samples with an effect size, Cohen's d, of -6.773 and power of 1.0.

NO2

LEZ vs ULEZ

Based on the p value of 0.0 and our alpha of 0.05 we reject the null hypothesis.

Due to these results, we can state that there is a difference between our samples with an effect size, Cohen's d, of -3.657 and power of 1.0.

The effect of ULEZ on the levels of NO2 was **0.848 times** than the effect of LEZ.'

PM10

LEZ vs ULEZ

Based on the p value of 0.0 and our alpha of 0.05 we reject the null hypothesis.

Due to these results, we can state that there is a difference between our samples with an effect size, Cohen's d, of -7.025 and power of 1.0.

The effect of ULEZ on the levels of PM10 was **3.66 times greater** than the effect of LEZ.