Principles of Quantum Mechanics

Notes made by Finley Cooper

23rd October 2026

Contents

1 Introduction **3**

2 Dirac Formalism **3**
2.1 Concepts and Definitions 3
2.2 The Harmonic Oscillator 6
2.2.1 Construction of the eigenstates 6
2.2.2 Importance of the oscillator 9
2.3 Bases and Measurements 10
2.3.1 Bases of Eigenstates 10
2.3.2 Functions of Operators 10
2.3.3 Measurement Axioms 11
2.3.4 Simultaneous measurement and complete commuting sets 12
2.4 Bases, Representations and Wavefunctions 13
2.4.1 Discrete and Continuous Bases 13
2.4.2 Position and Momentum Bases 14
2.4.3 Position and Momentum Eigenstates 17

3 Evolution in Time and Pictures **18**
3.1 The Schrödinger Equation and the Schrödinger Picture 18
3.2 Heisenberg Picture and Heisenberg Equations 21
3.2.1 Unitary operators 21
3.2.2 Changing Pictures 21

1 Introduction

Recall from IB Quantum Mechanics we have the following:

(i) Wave-particle duality. Waves behave like particles, like light quanta, and vice versa (particles behave like waves).

(ii) We have a wavefunction $\psi(x)$ for a particle which we interpret as a probability density $|\psi(x)|^2$ (i.e. probability is intrinsic to the theory).

(iii) Observables become operators on wavefunctions:

- $\hat{x} \to x$;
- $\hat{p} \to -i\hbar\frac{\partial}{\partial x}$

The lack of commutation, since $[\hat{x}, \hat{p}] = i\hbar$, leads to the uncertainty relation.

(iv) Schrödinger's equation with Hamiltonian H specifies dynamics, and eigenvalues of H are the allowed energy levels of the system.

(v) All of this explains the hydrogen atom structure and the resulting line spectra.

The aims of this course are the following:

(i) Reformulate quantum mechanics in a more powerful, flexible and useful form: the Dirac formulation, which goes beyond wavefunctions. The previous description appears as a particular realisation of our generalisation of quantum states and also leads to alternatives such as

$$\tilde{\psi}(p) \quad \text{with} \quad \hat{x} \to i\hbar\frac{\partial}{\partial p} \quad \text{and} \quad \hat{p} \to p.$$

(ii) This more general approach allows much simpler analysis of known problems (for example, the Simple Harmonic Oscillator), but also lots of new things such as *spin* of particles and symmetries (translations, rotations) and the related conservation laws.

(iii) We will also look at identical particles (bosons and fermions).

(iv) We will develop a framework for quantising more general systems such as EM fields.

(v) We will also set up perturbation theory, which is a practical tool for computation, but we will not heavily consider the detailed applications.

2 Dirac Formalism

2.1 Concepts and Definitions

A quantum system is described at each instant by a *state* $|\psi\rangle$ which belongs to a complex vector space V; any non-zero element of this space V is a possible state. Thus

$$|\psi\rangle, |\phi\rangle \in V \implies \alpha|\psi\rangle + \beta|\phi\rangle \in V \quad \text{for any } \alpha, \beta \in \mathbb{C}.$$

Physically, this is a superposition principle leading to wave-like behaviour, but these states are *not wavefunctions*. There are also *dual* or conjugate states $\langle\psi|$ which belong to the dual space

$V^\dagger$. By definition, states and duals can be combined or paired to give a complex number. The map

$$(\langle\psi|\,, |\phi\rangle) \mapsto \langle\psi|\phi\rangle$$

defines a map from $V^\dagger \times V \to \mathbb{C}$ with

$$\langle\phi|\,(\alpha_1\,|\psi_1\rangle + \alpha_2\,|\psi_2\rangle) = \alpha_1\,\langle\phi|\psi_1\rangle + \alpha_2\,\langle\phi|\psi_2\rangle$$

and

$$(\beta_1\,\langle\phi_1| + \beta_2\,\langle\phi_2|)\,|\psi\rangle = \beta_1\,\langle\phi_1|\psi\rangle + \beta_2\,\langle\phi_2|\psi\rangle\,.$$

The spaces V and $V^\dagger$ come with an inner product which can be described as a one-to-one correspondence between the spaces

$$\begin{aligned} V &\to V^\dagger \\ |\psi\rangle &\to \langle\psi| = (|\psi\rangle)^\dagger\,. \end{aligned}$$

We also have that if we take a linear combination of states to the dual space, the coefficients have to have their complex conjugates taken when mapping from the dual space to V or vice versa. Hence the inner product is

$$|\phi\rangle\,, |\psi\rangle \mapsto \langle\phi|\psi\rangle = (|\phi\rangle)^\dagger\,|\psi\rangle$$

and it is assumed to obey

$$\langle\phi|\psi\rangle = (\langle\psi|\phi\rangle)^*$$

since the inner product is Hermitian, and the inner product defines a positive-definite norm on V:

$$\|\,|\psi\rangle\,\|^2 = \langle\psi|\psi\rangle\,.$$

Remark. Knowing $\langle\phi|\psi\rangle$ for all $\langle\phi|$ determines $|\psi\rangle$ and vice versa.

We can show this by

$$\langle\phi|\psi_1\rangle = \langle\phi|\psi_2\rangle \implies \langle\phi|\chi\rangle = 0,$$

where $|\chi\rangle = |\psi_1\rangle - |\psi_2\rangle$. If this holds for all $\langle\phi|$, we can set $\langle\phi| = \langle\chi|$, so by positive definiteness we have that the norm of $|\chi\rangle$ is zero, and therefore $|\chi\rangle = 0$ too.

We also have that the physical content of any state is unchanged by

$$|\psi\rangle \to \alpha\,|\psi\rangle$$

for any non-zero $\alpha \in \mathbb{C}$. So we will usually normalise to make the state have norm 1, but we can still change it by a phase factor.

Remark. Relative phases can be important in expressions where we're adding a linear combination of elements of V. We also assume the space V is complete so appropriate sequences and series converge in V.

A complete inner product space is called a *Hilbert space*.

Definition. (Operator) An *operator* Q is a linear map on states

$$|\psi\rangle \to Q\,|\psi\rangle\,.$$

The same operator can be thought of as acting to the left on dual states, so

$$\langle\phi| \to \langle\phi|\,Q$$

where the right-hand side is defined by

$$(\langle\phi|\,Q)\,|\psi\rangle = \langle\phi|\,(Q\,|\psi\rangle) = \langle\phi|\,Q\,|\psi\rangle\,.$$

Remark. States and operators $|\psi\rangle, \langle\phi|, Q$ can be dealt with by rulers that mirror those for complex vectors $\mathbf{u}(\text{col}), \mathbf{v}^\dagger(\text{row})$ and a matrix A. So we can compare $\langle\psi|\,Q\,|\phi\rangle$ with $\mathbf{v}^\dagger A\mathbf{u}$ like we do with inner products.

Definition. (Eigenstate) For any operator Q with say that $|\psi\rangle$ non-zero is an *eigenstate* with *eigenvalue* λ if

$$Q\,|\psi\rangle = \lambda\,|\psi\rangle$$

or equivalently

$$\langle\psi|\,Q^\dagger = \lambda^*\,\langle\psi|\,.$$

Remark. For a general Q, λ may be complex.

Definition. (Hermitian Conjugate) For any operator Q, the *Hermitian conjugate* or *adjoint map* $Q^\dagger$ is an operator defined by

$$\langle\psi|\,Q^\dagger = (Q\,|\psi\rangle)^\dagger$$

for any $|\psi\rangle$.

Equivalently

$$\begin{aligned}\langle\phi|\,Q^\dagger\,|\psi\rangle &= (Q\,\langle\phi|)^\dagger\,|\psi\rangle \\ &= \langle\psi|\,Q\,|\phi\rangle^\dagger\end{aligned}$$

for all $|\psi\rangle, |\phi\rangle$.

Definition. (Hermitian) An operator Q is *Hermitian* or *self-adjoint* if

$$Q^\dagger = Q$$

and such operators are called *observables* because they correspond to physical quantities that can be measured, for example position, momentum, energy.

To define a quantum system, we must specify the *Hamiltonian* whose eigenvalues are the allowed energy levels of the system,

$$H|\chi\rangle = E|\chi\rangle.$$

We have the following key results for any Hermitian operator Q.

Proposition. For a Hermitian operator Q,
(i) All eigenvalues are real;
(ii) Eigenstates with distinct eigenvalues are orthogonal;
(iii) Any state can be written as a linear combination of its eigenstates (i.e its matrix representation is diagonalisable).

For a proof of (iii) we have a proof in IB Linear Algebra. We'll cover a proof of (i) and (ii) here.

Proof. Suppose that $Q|\psi\rangle = \lambda|\psi\rangle$. So taking Hermitian conjugates on both sides we have that

$$\begin{aligned}\langle\psi|Q^\dagger &= \lambda^*\langle\psi| \\ &= \langle\psi|Q.\end{aligned}$$

Hence $\langle\psi|Q|\psi\rangle = \lambda\langle\psi|\psi\rangle = \lambda^*\langle\psi|\psi\rangle$. Since $|\psi\rangle$ is non-zero, it's norm is non-zero hence $\lambda = \lambda^*$.

For the second part let $|n\rangle$ be an eigenstate of Q with eigenvalues $\lambda = q_n$ real. Then we have that

$$\begin{aligned}Q|n\rangle &= q_n|n\rangle \\ Q|m\rangle &= q_m|m\rangle \\ \langle m|Q &= q_m\langle m|\end{aligned}$$

hence $\langle m|Q|n\rangle = q_n\langle m|n\rangle = q_m\langle m|n\rangle$. But we have that $q_m \neq q_n$ so $\langle m|n\rangle = 0$. □

2.2 The Harmonic Oscillator

2.2.1 Construction of the eigenstates

A one-dimensional harmonic oscillator of mass m and frequency ω is a system with position $\hat{x}$, momentum $\hat{p}$ and Hamilitonian

$$H = \frac{1}{2m}\hat{p}^2 + \frac{1}{2}m\omega^2\hat{x}^2.$$

We seek the energy eigenstates and eigenvalues of H. We do this by introducting *annihilation, creation, and number operators.*

Define

$$a = \left(\frac{m\omega}{2\hbar}\right)^{\frac{1}{2}} \left(\hat{x} + \frac{i\hat{p}}{m\omega}\right)$$
$$a^\dagger = \left(\frac{m\omega}{2\hbar}\right)^{\frac{1}{2}} \left(\hat{x} - \frac{i\hat{p}}{m\omega}\right).$$

Equivalently

$$\hat{x} = \left(\frac{\hbar}{2m\omega}\right)^{\frac{1}{2}} \left(a + a^\dagger\right)$$
$$\hat{p} = \left(\frac{\hbar m\omega}{2}\right)^{\frac{1}{2}} i\left(a^\dagger - a\right).$$

It's easy to check that

$$[\hat{x}, \hat{p}] = i\hbar \iff [a, a^\dagger] = 1.$$

In fact,

$$\begin{aligned} aa^\dagger &= \frac{m\omega}{2\hbar}\hat{x}^2 + \frac{1}{2m\omega\hbar}\hat{p}^2 - \frac{i}{2\hbar}(\hat{x}\hat{p} - \hat{p}\hat{x}) \\ &= \frac{1}{\hbar\omega}H + \frac{1}{2} \end{aligned}$$

but

$$a^\dagger a = \frac{1}{\hbar\omega}H - \frac{1}{2}.$$

This confirms the commutation relation, but we also find that

$$H = \hbar\omega\left(N + \frac{1}{2}\right)$$

with $N = a^\dagger a$, the *number operator* which is clearly Hermitian, so finding eigenvalues and eigenstates of N is equivalent to finding these for H.

Let $|\lambda\rangle$ be any eigenstate of N with eigenvalue λ, so

$$N|\lambda\rangle = \lambda|\lambda\rangle \iff H|\lambda\rangle = E|\lambda\rangle$$

where $E = \hbar\omega\left(\lambda + \frac{1}{2}\right)$. Then assuming that $|\lambda\rangle$ is normalised, write

$$\lambda = \langle\lambda| N |\lambda\rangle = \lambda a^\dagger a|\lambda\rangle = (a|\lambda\rangle)^\dagger (a|\lambda\rangle) = ||a|\lambda\rangle||^2$$

we could have just seen this directly, since N is clearly positive semi-definite from its construction as the outer product of an operator. So we have that $\lambda \geq 0$ and $\lambda = 0 \iff a|\lambda\rangle = 0$.

Now we consider

$$\begin{aligned} N(a^\dagger|\lambda\rangle) &= ([N, a^\dagger] + a^\dagger N)|\lambda\rangle \\ &= (a^\dagger + a^\dagger\lambda)|\lambda\rangle \\ &= (\lambda + 1)(a^\dagger|\lambda\rangle). \end{aligned}$$

Similarly we have that

$$\begin{aligned} N(a\,|\lambda\rangle) &= = ([N,a] + aN)\,|\lambda\rangle \\ &= (-a + a\lambda)\,|\lambda\rangle \\ &= (\lambda - 1)a\,|\lambda\rangle\,. \end{aligned}$$

Here we have used the results for commutators $[N, a^\dagger] = [a^\dagger, a^\dagger] = a^\dagger[a, a^\dagger] = a^\dagger$ and similarily $[N, a] = -a$.

Now the news states with be eigenstates provided their norms are non-zero. We can compute

$$\begin{aligned} ||a\,|\lambda\rangle\,||^2 &= \lambda \\ ||a^\dagger\,|\lambda\rangle\,||^2 &= \langle\lambda|\,aa^\dagger\,|\lambda\rangle = \lambda a^\dagger a + 1\,|\lambda\rangle \\ &= \lambda + 1. \end{aligned}$$

Because of these properties $a^\dagger$ and a are called *creation* (or raising) and *annihilation* (or lowering) operators.

We can't lower below zero, since the eigenvalues are positive so the eigenvalues must be a subset of the positive integers to avoid a contradiction. We can lower and raise all we want, so we can generate any other state with eigenvalues being all of the non-negative integers.

More formally, since we know that $\lambda \geq 0$, so for a given eigenstate $|\lambda\rangle$, if λ is *not* an integer, then

$$a\,|\lambda\rangle\,, a^2\,|\lambda\rangle\,, \ldots, a^n\,|\lambda\rangle$$

are states with eigenvalues

$$\lambda - 1, \lambda - 2, \ldots, \lambda - n$$

but for n sufficiently large $\lambda - n < 0$ which is a contradiction.

By contrast if $\lambda = k \in \mathbb{N}$ then the contradiction is avoided because

$$a\,|k\rangle\,, a^2\,|k\rangle\,, \ldots, a^k\,|k\rangle$$

with eigenvalues

$$k - 1, k - 2, \ldots, 0$$

so we can't lower anymore because our eigenstate has collapsed to zero, i.e. if we force integer eigenvalues the resultant eigenstate is *nilpotent*.

We can use the raising operator $a^\dagger$ to generate the eigenstates with eigenvalues $k + 1, k + 2, \ldots$ forever.

Hence the eigenvalues of N are therefore precisely the non-negative integers and the oscillator has energy levels

$$E_n = \hbar\omega\left(n + \frac{1}{2}\right).$$

From our calculation of norms above, we can choose normalised eigenstates $|n\rangle$ related by

$$\begin{aligned} a^\dagger\,|n\rangle &= \sqrt{n+1}\,|n+1\rangle \\ a\,|n\rangle &= \sqrt{n}\,|n-1\rangle\,. \end{aligned}$$

Starting from any state, $|n\rangle$, we can generate all others. In particular, starting from the *ground state* $|0\rangle$, characterised by

$$a\,|0\rangle = 0$$

we can construct normalised eigenstates

$$|n\rangle = \frac{1}{\sqrt{n!}}(a^\dagger)^n\,|0\rangle\,.$$

Remark. Beware that $|0\rangle$ is *not* zero in V. It is an eigenstate and therefore non-zero by definition.

2.2.2 Importance of the oscillator

(i) A smooth potential $V(x)$ with equilibrium point $x = x_0$ (so that $V'(x_0) = 0$) can be expanded by

$$V(x_0 + x) = V(x_0) + \frac{1}{2}V''(x_0)x^2 + O(x^3)$$

so, if $x = x_0$ is *stable*, then we can set $V''(x_0) = m\omega^2 > 0$ and for displacements x not too large, we can expect the system to be well approximated by a oscillator.

(ii) This can also be applied in the case for a diatomic molecules where we have quantisation of vibrational energies which is important in understanding the internal energy and heat capacities which result in macroscopic consequences.

(iii) More complicated systems can be analysed in terms of normal modes. Coherent motion in which all degrees of freedom oscillate with the same common frequency ω. The general solution for a classical oscillator is

$$x(t) = Ae^{-i\omega t} + A^* e^{i\omega t}$$

where A is a complex constant.

Normal modes for a system with variables x_N are solutions of the form

$$x_N(t) = Au_N e^{-i\omega t} + A^* u_N^* e^{i\omega t}$$

with u_N a complex vector specific to each normal frequency ω.

(iv) For example consider a crystal with a number of atoms $\mathcal{N} \sim 10^{23}$ where the force are approximately elastic and N labels diretions of motion for each atom, so we have $3 \times \mathcal{N}$ values.

(v) Or we consider an EM field, where normal modes are oscillations of electric and mangetic at each point in space are given by

$$\mathbf{E}(\mathbf{x}, t) = A\mathbf{u}(\mathbf{x})e^{-i\omega t} + A^*\mathbf{u}(\mathbf{x})^* e^{i\omega t}.$$

This is exact.

(vi) Normal modes are independent oscillator: treat these as above. For each mode we have

$$a(\omega) \quad \text{and} \quad a(\omega)^\dagger$$

where are ground state $|0\rangle$ with $a(\omega)\,|0\rangle = 0$. Non-ground states created with the raising operators add energy and momentum and behave like particles. In this way we can see that a destroys particles and $a^\dagger$ creates particles.

2.3 Bases and Measurements

2.3.1 Bases of Eigenstates

We know that for any observable Q there is an orthonormal basis of eigenstates $\{|n\rangle\}$ for V,

$$Q\,|n\rangle = q_n\,|n\rangle\,.$$

We have orthonormality $\langle m|n\rangle = \delta_{mn}$ and for any state $|\psi\rangle \in V$ we have that

$$|\psi\rangle = \sum_n \alpha_n\,|n\rangle$$

where $\alpha_n = \langle n|\psi\rangle$. Note that $||\,|\psi\rangle\,||^2 = 1$ if and only if

$$\langle\psi|\psi\rangle = \left(\sum_m \alpha_m^*\,\langle m|\right)\left(\sum_n \alpha_n\,|n\rangle\right) = \sum_n |\alpha_n|^2 = 1.$$

Note that we can write

$$1 = \sum_n |n\rangle\,\langle n|\,.$$

since for any state $|\psi\rangle$ we have that

$$\begin{aligned}|\psi\rangle &= \left(\sum_n |n\rangle\,\langle n|\right)|\psi\rangle \\ &= \sum_n |n\rangle\,(\langle n|\psi\rangle)\end{aligned}$$

In above there may be several independent staets with the same eigenvalue λ. The eigenspace for a given eigenvalue is

$$V_\lambda = \{|\psi\rangle : Q\,|\psi\rangle = \lambda\,|\psi\rangle\}.$$

Note that V_λ has basis $\{|n\rangle : q_n = \lambda\}$. The *degeneracy* of λ is the number of sattes in this basis (i.e the dimension of V_λ). If V_λ has dimension 1, then we say that λ is non-degenerate.

2.3.2 Functions of Operators

It is straight forward to define $f(Q)$ for some operator Q if f is a polynmoial or power series (assuming convergence). For example we can consider $f(Q) = Q^3 + Q$ or $f(Q) = e^Q$ in this way. Important examples are $e^{-iHt/\hbar}$ which is the time-evolution operator and $e^{-ia\hat{p}/\hbar}$ which is the translation operator through some fixed displacement a in the x direction.

More generally, for Q an observable, with basis of eigenstates, we can define

$$f(Q)\,|n\rangle = f(q_n)\,|n\rangle$$

and then defined on any state, provided that $f(q_n)$ makes sense. For example for $f(Q) = \frac{1}{Q}$, we have that

$$\frac{1}{Q}\,|n\rangle = \frac{1}{q_n}\,|n\rangle$$

for $q_n \neq 0$.

In addition we have that

$$Q = \sum_n q_n \left|n\right\rangle \left\langle n\right|$$
$$f(Q) = \sum_n f(q_n) \left|n\right\rangle \left\langle n\right|.$$

Recall our eigenspace V_λ and consider a projection operation P_λ onto this subspace, $V \to V_\lambda$ can be defined by its action on basis states

$$P_\lambda \left|n\right\rangle = \begin{cases} \left|n\right\rangle & q_n = \lambda \\ 0 & \text{otherwise} \end{cases}$$

which can be written as

$$P_\lambda = \sum_{n:\ q_n=\lambda} \left|n\right\rangle \left\langle n\right|.$$

2.3.3 Measurement Axioms

Consider a measurement of Q (observable) when the system is in a normalised state $\left|\psi\right\rangle$. Then

(i) The result in an eigenvalue λ of Q;

(ii) This result is obtained with probability

$$p(\lambda) = ||P_\lambda \left|\psi\right\rangle||^2 = \sum_{n:\ q_n=\lambda} |\alpha_n|^2.$$

(iii) Immediately after the measurement the state is

$$cP_\lambda \left|\psi\right\rangle = c \sum_{n:\ q_n=\lambda} \alpha_n \left|n\right\rangle$$

where c is a constant for normalisation.

It is often the case with λ being a non-generate eigenstate. In that case we just get that $p(\lambda) = |\left\langle n|\psi\right\rangle|^2 = |\alpha_n|^2$ where α_n is often called the amplitude of the state.

We have that

$$\sum_\lambda p(\lambda) = 1.$$

Example. Consider a system with space V with orthonormal basis states

$$\left|1\right\rangle, \left|2\right\rangle, \left|3\right\rangle$$

which are eigenstaets of Q with eigenvalues $q_1 = q_2 = 0$ and $q_3 = 1$. Consider

$$\left|\psi\right\rangle = \frac{1}{\sqrt{6}} \left(2\left|1\right\rangle + \left|2\right\rangle + \left|3\right\rangle\right)$$

So we get that

$$p(1) = \left|\frac{1}{\sqrt{6}}\right|^2 = \frac{1}{6}$$
$$p(0) = \left|\frac{2}{\sqrt{6}}\right|^2 + \left|\frac{1}{\sqrt{6}}\right|^2 = \frac{5}{6}$$

and the state afterwards is $|3\rangle$ if 1 was measured and is

$$\frac{1}{\sqrt{5}}(2\,|1\rangle + |2\rangle)$$

if we measured 0.

Definition. (Expectation value) The *expectation value* or *mean* of Q in a state $|\psi\rangle$ is

$$\langle Q\rangle_\psi = \langle\psi|\, Q\,|\psi\rangle = \sum_\lambda \lambda p(\lambda) = \sum_n q_n |\alpha_n|^2.$$

Definition. (Uncertainty) The *uncertainty* or *variance* of an operator Q in a state $|\psi\rangle$ is

$$(\Delta Q)^2_\psi = \langle (Q - \langle Q\rangle_\psi)^2\rangle_\psi = \langle Q^2\rangle_\psi - \langle Q\rangle^2_\psi.$$

Remark. In thinking about many repeated measurements, we must prepare the system in the same state $|\psi\rangle$ each time.

Remark. If $|\psi\rangle$ is an eigenstate of Q with eigenvalue λ then

$$\langle Q\rangle_\psi = \lambda \quad \text{and} \quad (\Delta Q)_\psi = 0.$$

Quantum mechanical behaviour arises from the fact that certain obserables do not commute. In general, in any state $|\psi\rangle$

$$(\Delta A)_\psi (\Delta B)_\psi \geq \frac{1}{2}\,|\langle [A,B]\rangle_\psi|$$

so $[A,B] \neq 0$ implies that we cannot expect to measure A and B at the same time.

A proof of the general uncertainty relation can be seen from expanding $||(A + i\lambda B)\,|\psi\rangle\,||^2 \geq 0$.

The prototypical case is with position and momentum $\hat{x}, \hat{p}$ which obey

$$[\hat{x}, \hat{p}] = i\hbar$$

in one dimension and $\Delta x \Delta p \geq \frac{\hbar}{2}$.

In three dimensions we have

$$[\hat{x}_i, \hat{p}_j] = i\hbar\delta_{ij}$$

and the uncertainty applies to position and momentum in a particular direction.

2.3.4 Simultaneous measurement and complete commuting sets

If an observable Q does not distinguish states completely because some eigenvalues are degenerate, then we can look for an additional obserable Q' such that Q, Q' are simultaneously measureable (i.e simultaneously diagonalisable) and preserves the eigenspaces of the other operator. This means we have joint eigenstates $|\lambda, \lambda'\rangle$ with

$$Q\,|\lambda,\lambda'\rangle = \lambda\,|\lambda,\lambda'\rangle$$
$$Q'\,|\lambda,\lambda'\rangle = \lambda'\,|\lambda,\lambda'\rangle$$

so $[Q, Q'] = 0$. Conversely if $[Q, Q'] = 0$, it is easy to check that $Q' : V_\lambda \to V_\lambda$ and hence we can find a basis of eigenstates of Q' in V_λ which gives us joint eigenstates for Q and Q'. Since we assume that Q is Hermitian so V is a direct sum of eigenstates, we can do this for all eigenstates, giving us a complete basis of joint eigenstates, which we call $|\lambda, \lambda'\rangle$.

Definition. (Complete commuting set) A set of observables $\{Q, Q', Q'', \dots\}$ is a *complete commuting set* if their pairwise commute. This implies that there is a basis of joint eigenstates, and the set of eigenvalues $\lambda, \lambda', \lambda'', \dots$ label states uniquely (up to normalisation)

Remark. Equivalently if A is any observable such that $[A, Q] = [A, Q'] = \cdots = 0$ then $A = f(Q, Q', \dots)$.

Example. Consider the previous example with orthonormal satets $|1\rangle, |2\rangle, |3\rangle$ and eigenstates of Q with $\lambda = 0$ for $|1\rangle$ and $|2\rangle$ and $\lambda = 1$ for $|3\rangle$. Now consider a second observabke Q' with $Q'|1\rangle = |2\rangle, Q'|2\rangle = |1\rangle, Q'|3\rangle = 0$. Now by writing a general vector in the orthonormal basis, we have that $[Q, Q'] = 0$. Now let

$$|\pm\rangle = \frac{1}{\sqrt{2}}(|1\rangle \pm |2\rangle)$$

so then

$$Q'|\pm\rangle = \pm|\pm\rangle$$

Q, Q' is a complete commuting set with basis of joint eigenstates $|+\rangle, |-\rangle, |3\rangle$.

2.4 Bases, Representations and Wavefunctions

2.4.1 Discrete and Continuous Bases

Consider orthonormal basis $\{|n\rangle\}$ with n discrete label. We can regard all states and operators as row or column vectors and matrices respectively

$$|\psi\rangle = \sum_n \alpha_n |n\rangle \iff \alpha_n \qquad = \langle n|\psi\rangle$$
$$|\phi\rangle = \sum_n \beta_n |n\rangle \iff \beta_n \qquad = \langle n|\phi\rangle$$

and inner product

$$\langle\phi|\psi\rangle = \sum_n \beta_n^* \alpha_n.$$

For an operator A,

$$A|n\rangle = \sum_m |m\rangle A_{mn}$$

where A_{mn} is a complex matrix with entries $\langle m| A |n\rangle$.

It's simple to check that

$$|\phi\rangle = A|\psi\rangle \iff \beta_m = \sum_n A_{mn}\alpha_n$$

and also if B is another operator with matrix $B_{mn} = \langle m| B |n\rangle$ then

$$(AB)_{mn} = \sum_p A_{mp} B_{pn}$$

i.e matrix multiplication.

The choice of basis provides a representation of the quantum states and operators. We can also allow the index n to be continuous. This requires modifications such as the following:

$$\begin{aligned} \sum_n &\longrightarrow \int \mathrm{d}n \\ \langle n|m\rangle = \delta_{nm} &\longrightarrow \delta(n-m) \\ |\psi\rangle = \sum_n \alpha \, |n\rangle &\longrightarrow \int \mathrm{d}n \; \alpha_n \, |n\rangle \end{aligned}$$

and the completeness relation goes to

$$I = \sum_n |n\rangle \langle n| \longrightarrow \int \mathrm{d}n \, |n\rangle \langle n| \, .$$

Note that if basis a bove arise as eigenstates of an obserable then

$$|\alpha_n|^2 = |\langle n|\psi\rangle|^2 \longrightarrow \text{probability density for continuous range } n.$$

2.4.2 Position and Momentum Bases

Considera particle in one dimension with position $\hat{x}$ and momentum $|p\rangle$ obeying the relation $[\hat{x}, \hat{p}] = i\hbar$. Let $|x\rangle$ be the position eigenstates

$$\hat{x} \, |x\rangle = x \, |x\rangle$$

which are continuous eigenvalues with

$$\langle x|x'\rangle = \delta(x - x'), \quad \int \mathrm{d}x \, |x\rangle \langle x| = I.$$

Any state can be written as

$$|\psi\rangle = \int \mathrm{d}x \; \psi(x) \, |x\rangle$$

with $\psi(x) = \langle x|\psi\rangle$ and this is the usual position space wavefunction. From our remarks in the previous section $|\psi(x)|^2$ is a probability density for measuring position.

Our inner product of two wavefunctions from Part IB Quantum mechanics is recovered through

$$\begin{aligned} \langle \phi|\psi\rangle &= \left(\int \mathrm{d}x \; \phi(x)^* \, \langle x| \right) \left(\int \mathrm{d}x' \; \psi(x') \, |x\rangle' \right) \\ &= \int \mathrm{d}x \int \mathrm{d}x' \; \phi(x)^* \psi(x') \, \langle x|x'\rangle = \int \mathrm{d}x \; \phi(x)^* \psi(x) \end{aligned}$$

and

$$|| \, |\psi\rangle \, ||^2 = \langle\psi|\psi\rangle = \int \mathrm{d}x |\psi(x)|^2 = 1$$

if normalised.

Similarly, define momentum eigenstate $|p\rangle$ with

$$\hat{p} \, |p\rangle = p \, |p\rangle$$

and

$$\langle p|p'\rangle = \delta(p - p'), \quad \int \mathrm{d}p \, |p\rangle \, \langle p| = I.$$

Now we can expand

$$|\psi\rangle = \int \mathrm{d}p \; \tilde{\psi}(p) \, |p\rangle$$

with $\tilde{\psi}(p) = \langle p|\psi\rangle$ momentum space wavefunction and $|\tilde{\psi}(p)|^2$ probability distribution for measurements of momentum.

To proceed, we need the following result that

$$\langle x|p\rangle = \frac{1}{(2\pi\hbar)^{1/2}} e^{ipx/\hbar}$$
$$\langle p|x\rangle = \frac{1}{(2\pi\hbar)^{1/2}} e^{-ipx/\hbar}$$

which we will assume holds. Consider the action of $\hat{x}$ and $\hat{p}$ on position space wavefunctions. So suppose we have $\hat{x} \, |\psi\rangle$ has wavefunction

$$\langle x| \, \hat{x} \, |\psi\rangle = x \, \langle x|\psi\rangle = x\psi(x).$$

Similarly if we have $\hat{p} \, |\psi\rangle$, the wavefunction is

$$\begin{aligned} \langle x| \, \hat{p} \, |\psi\rangle &= \int \mathrm{d}p \, \langle x| \, \hat{p} \, |p\rangle \, \langle p|\psi\rangle \\ &= \int \mathrm{d}p \; p \, \langle x|p\rangle \, \langle p|\psi\rangle \end{aligned}$$

and note that $p \, \langle x|p\rangle = -i\hbar \frac{\partial}{\partial x} \, \langle x|p\rangle$, so hence

$$\begin{aligned} \langle x| \, \hat{p} \, |\psi\rangle &= -i\hbar \frac{\partial}{\partial x} \int \mathrm{d}p \;\; \langle x|p\rangle \, \langle p|\psi\rangle \\ &= -i\hbar \frac{\partial}{\partial x} \, \langle x|\psi\rangle \\ &= -i\hbar\psi'(x). \end{aligned}$$

which gives us the action of $\hat{x}$ and $\hat{p}$ on position space wavefunctions. Similarly on momentum space wavefunctions we calculate $\langle p| \, \hat{x} \, |\psi\rangle$ and $\langle p| \, \hat{p} \, |\psi\rangle$ and the result is that

$$\hat{x} \longrightarrow i\hbar \frac{\partial}{\partial p}$$
$$\hat{p} \longrightarrow p.$$

The relationship between $\psi(x)$ and $\tilde{\psi}(p)$ also follows from our assumed result, as

$$\begin{aligned}\tilde{\psi}(p) = \langle p|\psi\rangle &= \int \mathrm{d}x \, \langle p|x\rangle \, \langle x|\psi\rangle \\ &= \frac{1}{(2\pi\hbar)^{1/2}} \int \mathrm{d}x \; e^{-ipx/\hbar}\psi(x)\end{aligned}$$

and

$$\begin{aligned}\psi(x) = \langle x|\psi\rangle &= \frac{1}{(2\pi\hbar)^{1/2}} \int \mathrm{d}p \, \langle x|p\rangle \, \langle p|\psi\rangle \\ &= \frac{1}{(2\pi\hbar)^{1/2}} \int \mathrm{d}p \; e^{ixp/\hbar}\tilde{\psi}(p).\end{aligned}$$

Now given $H(\hat{x}, \hat{p})$ we can consider the time independent Schrödinger equation

$$H\,|\psi\rangle = E\,|\psi\rangle\,.$$

If $H = \frac{\hat{p}^2}{2m} + V(\hat{x})$ then on $\psi(x)$ we get that

$$H \to -\frac{\hbar^2}{2m}\frac{\partial^2}{\partial x^2} + V(x)$$

and on $\tilde{\psi}(p)$ we get that

$$H \to \frac{p^2}{2m} + V\left(i\hbar\frac{\partial}{\partial p}\right).$$

It may be easy to interpret the potential term in momentum space for something like $V(x) = \lambda x^n$ which gives

$$V\left(i\hbar\frac{\partial}{\partial p}\right) = \lambda(i\hbar)^n\left(\frac{\partial}{\partial p}\right)^n$$

for a positive integer n. More generally this isn't obvious what the potential term in momentum space means. We will argue from first principles as follows.

Consider

$$\begin{aligned}\langle p|\,V(\hat{x})\,|\psi\rangle &= \int \mathrm{d}x \, \langle p|\,V(\hat{x})\,|x\rangle \, \langle x|\psi\rangle \\ &= \int \mathrm{d}x \int \mathrm{d}p' \, \langle p|\,V(\hat{x})\,|x\rangle \, \langle x|p'\rangle \, \langle p'|\psi\rangle \\ &= \int \mathrm{d}p' \left(\int \mathrm{d}x \; V(x) \, \langle p|x\rangle \, \langle x|p'\rangle \, \langle p'|\psi\rangle\right) \\ &= \int \mathrm{d}p' \int \mathrm{d}x \; V(x)\frac{e^{-i(p-p')x/\hbar}}{2\pi\hbar}\tilde{\psi}(p) \\ &= \frac{1}{(2\pi\hbar)^{1/2}} \int \mathrm{d}p' \; \tilde{V}(p-p')\tilde{\psi}(p')\end{aligned}$$

Which is a reproduction of the convolution theorem. So $H\,|\psi\rangle = E\,|\psi\rangle$ becomes

$$-\frac{\hbar^2}{2m}\psi''(x) + V(x)\psi(x) = E\psi(x)$$

in position form. Or

$$\frac{p^2}{2m}\tilde{\psi}(p) + \frac{1}{(2\pi\hbar)^{1/2}}\int \mathrm{d}p'\tilde{V}(p-p')\tilde{\psi}(p') = E\tilde{\psi}(p)$$

in momentum space.

To generalise from one to three dimensions we use the commutation relation that

$$[\hat{x}_i, \hat{p}_j] = i\hbar\delta_{ij}.$$

One complete commuting set is

$$\hat{\mathbf{x}} = (\hat{x}_1, \hat{x}_2, \hat{x}_3)$$

with joint eigenstates

$$\hat{x}_i \left|\mathbf{x}\right\rangle = x_i \left|\mathbf{x}\right\rangle$$

which are orthonormal, so $\langle \mathbf{x}|\mathbf{x}'\rangle = \delta^{(3)}(\mathbf{x}-\mathbf{x}')$.

An alternative set

$$\hat{\mathbf{p}} = (\hat{p}_1, \hat{p}_2, \hat{p}_3)$$

with joint eigenstates

$$\hat{p}_i \left|\mathbf{p}\right\rangle = p_i \left|\mathbf{p}\right\rangle$$

which are orthonormal in the same sense as the position operator. For any state $|\psi\rangle$, we can expand

$$\left|\psi\right\rangle = \int \mathrm{d}^3\mathbf{x}\ \psi(\mathbf{x})\left|\mathbf{x}\right\rangle = \int \mathrm{d}^3\mathbf{p}\ \tilde{\psi}(\mathbf{p})\left|\mathbf{p}\right\rangle$$

where $\psi(\mathbf{x}) = \langle \mathbf{x}|\psi\rangle$ and $\tilde{\psi}(\mathbf{p}) = \langle \mathbf{p}|\psi\rangle$. We also get that

$$\langle \mathbf{x}|\mathbf{p}\rangle = \frac{1}{(2\pi\hbar)^{3/2}} e^{i\mathbf{p}\cdot\mathbf{x}/\hbar}$$

and the wavefunctions are related by

$$\tilde{\psi}(\mathbf{p}) = \frac{1}{(2\pi\hbar)^{3/2}}\int \mathrm{d}^3\mathbf{x}\ e^{i\mathbf{p}\cdot\mathbf{x}/\hbar}\psi(\mathbf{x})$$

and the corresponding Fourier inverse.

2.4.3 Position and Momentum Eigenstates

Now we return to one dimension and seek to justify the result for $\langle x|p\rangle$ using only the commutator relation. Given $|x\rangle$ with $\hat{x}\left|x\right\rangle = x\left|x\right\rangle$, consider

$$T(a) = e^{-ia\hat{p}/\hbar} = \sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{-ia}{\hbar}\right)^n \hat{p}^n$$

and note that

$$[\hat{x}, T(a)] = \left[\hat{x}, \frac{-ia\hat{p}}{\hbar}\right] T(a) = aT(a).$$

Hence

$$\begin{aligned}\hat{x}(T(x)\,|x\rangle) &= ([\hat{x}, T(a)] + T(a)\hat{x})\,|x\rangle \\ &= (x+a)(T(x)\,|x\rangle)\end{aligned}$$

so we can think of $T(a)\,|x\rangle$ as a translated eigenstate.

Similarly consider $S(b) = e^{ib\hat{x}/\hbar}$ and note that $[\hat{p}, S(b)] = bS(b)$. Hence

$$\hat{p}\,|p\rangle = p\,|p\rangle \implies \hat{p}(S(b)\,|p\rangle) = (p+b)(S(b)\,|p\rangle)$$

so $S(b)$ gives a translation in momentum space.

Now take fixed values x_0 and p_0 and define all eigenstates by

$$|x_0 + a\rangle = T(a)\,|x_0\rangle\,, \quad |p_0 + b\rangle = S(b)\,|p_0\rangle\,.$$

Now observe that

$$\begin{aligned}\langle x_0 + a | p_0 + b\rangle &= \langle x_0|\, e^{ia\hat{p}/\hbar}\,|p_0 + b\rangle \\ &= \langle x_0|\, e^{ia(p_0+b)/\hbar} e^{ib\hat{x}/\hbar}\,|p_0\rangle \\ &= e^{(ab+ap_0+bx_0)/\hbar}\,\langle x_0|p_0\rangle\,.\end{aligned}$$

Now choose $\langle x_0|p_0\rangle = \frac{1}{\sqrt{2\pi\hbar}}$ for correct and consistent normalisation. Then set $x_0 = p_0 = 0$, so we get

$$\langle a|b\rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{iab/\hbar}.$$

3 Evolution in Time and Pictures

3.1 The Schrödinger Equation and the Schrödinger Picture

The evolution in time of state $|\psi(t)\rangle$ is governed (in absence of external influences of measurement) by the Schödinger equation

$$i\hbar \frac{\partial}{\partial t}\,|\psi(t)\rangle = H\,|\psi(t)\rangle$$

where $H = H^\dagger$ is the Hamiltonian.

Equivalently

$$-i\hbar \frac{\partial}{\partial t}\,\langle\psi(t)| = \langle\psi(t)|\,H.$$

This implies that

$$\frac{\partial}{\partial t}\,(\langle\psi(t)|\psi(t)\rangle) = 0$$

so that the normalisation of $|\psi(t)\rangle$ is preserved in time.

The solution of the Schrödinger equation can be written as

$$|\psi(t)\rangle = U(t)\,|\psi(0)\rangle$$

where $U(t) = e^{-iHt/\hbar}$ is the time-evolution operator, assuming that H is independent of t. We can check immediately from $i\hbar\frac{\partial U(t)}{\partial t} = HU(t)$.

Recall that time-evolution problems can be conveniently solved by working with basis of energy eigenstates $\{|n\rangle\}$ with

$$H|n\rangle = E_n|n\rangle .$$

with

$$e^{-iE_nt/\hbar}|n\rangle$$

being a solution of the time-dependent Schrödinger equation.

Generally if

$$|\psi(0)\rangle = \sum_n \alpha_n |n\rangle$$

where

$$H|n\rangle = E_n|n\rangle$$

then

$$|\psi(t)\rangle = \sum_n \alpha_n e^{-iE_nt/\hbar}|n\rangle .$$

Example. Consider a system with two energy eigenstates $|1\rangle$ and $|2\rangle$. Consider measuring Q defined by

$$Q|1\rangle = |2\rangle , \quad Q|2\rangle = |1\rangle$$

So $[Q, H] \neq 0$. The eigenstates of Q are

$$|\pm\rangle = \frac{1}{\sqrt{2}}(|1\rangle \pm |2\rangle)$$

with eigenvalues $q_\pm = \pm 1$. Take the initial state

$$|\psi(0)\rangle = |+\rangle .$$

This implies that

$$|\psi(t)\rangle = \frac{1}{\sqrt{2}}\left(e^{-iE_1t/\hbar}|1\rangle + e^{-iE_2t/\hbar}|2\rangle\right)$$

so the probability of measuring Q at time t and getting ± 1 is

$$\begin{aligned} |\langle\pm|\psi(t)\rangle|^2 &= \left|\frac{1}{2}(\langle 1| \pm |2\rangle)(e^{-iE_1t/\hbar}|1\rangle + e^{-iE_2t/\hbar}|2\rangle)\right|^2 \\ &= \left|\frac{1}{2}(e^{-iE_1t/\hbar} \pm e^{-iE_2t/\hbar})\right|^2 \\ &= \begin{cases} \cos^2(E_1 - E_2)t/2\hbar \\ \sin^2(E_1 - E_2)t/2\hbar \end{cases} . \end{aligned}$$

In the basis

$$|1\rangle \longrightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$
$$|2\rangle \longrightarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

then

$$H \longrightarrow \begin{pmatrix} E_1 & 0 \\ 0 & E_2 \end{pmatrix}$$

and

$$U = e^{-iHt/\hbar} \longrightarrow \begin{pmatrix} e^{-iE_1 t/\hbar} & 0 \\ 0 & e^{-iE_2 t/\hbar}. \end{pmatrix}$$

But instead we can work in the alternate basis where

$$|+\rangle \longrightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$
$$|-\rangle \longrightarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

So we have that

$$\begin{aligned} H\,|\pm\rangle &= \frac{1}{\sqrt{2}}(E_1\,|1\rangle + E_2\,|2\rangle) \\ \langle\pm|\,H\,|\pm\rangle &= \frac{1}{2}(E_1 + E_2) = \varepsilon \\ \langle\mp|\,H\,|\mp\rangle &= \frac{1}{2}(E_1 - E_2) = \hbar\lambda \end{aligned}$$

where we define ε and λ to be constants. Then we have that

$$H \longrightarrow \begin{pmatrix} \varepsilon & \hbar\lambda \\ \hbar\lambda & \varepsilon \end{pmatrix} = \varepsilon I + \hbar\lambda M$$

where I is the identity matrix and M is the matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ so now we have that

$$\begin{aligned} U(t) = e^{-iHt/\hbar} &= e^{-it(\varepsilon I + \lambda\hbar M)/\hbar} \\ &= e^{-it\varepsilon/\hbar} e^{-it\lambda M}. \end{aligned}$$

Note that $M^2 = I$ and hence we have that

$$\begin{aligned} \exp(-it\lambda M) &= \sum_{r=0}^{\infty} \frac{1}{(2r)!}(-i\lambda r)^{2r} I + \sum_{r=0}^{\infty} \frac{1}{(2r+1)!}(-i\lambda t)^{2r+1} M \\ &= (\cos\lambda t)I - i(\sin\lambda t)M. \end{aligned}$$

So we have that

$$|\psi(0)\rangle = |+\rangle \longrightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

so

$$|\psi(t)\rangle = e^{-iHt/\hbar}\,|+\rangle \longrightarrow e^{-it\varepsilon/\hbar} \begin{pmatrix} \cos\lambda t \\ -i\sin\lambda t \end{pmatrix}$$

which is therefore

$$e^{-it\varepsilon/\hbar}(\cos\lambda t\,|+\rangle - i\sin\lambda t\,|-\rangle).$$

Now compare with previous result for probability of measuring W and finding ± 1.

3.2 Heisenberg Picture and Heisenberg Equations

3.2.1 Unitary operators

Physical predictions in quantum mechanics are given by probability amplitudes, eigenvalues, expectation values, etc, hence we have expressions like

$$\langle\psi|\phi\rangle\,, \quad \langle\phi|\,A\,|\psi\rangle$$

and similar things.

Definition. (Unitary) An operator U is called *unitary* if

$$UU^\dagger = U^\dagger U = I$$

or equivalently

$$U^\dagger = U^{-1}.$$

Given any such U consider a map on states and operators taking

$$\begin{aligned} |\psi\rangle &\longrightarrow U\,|\psi\rangle = |\psi'\rangle \\ \langle\psi| &\longrightarrow \langle\psi|\,U^\dagger = \langle\psi'| \\ A &\longrightarrow UAU^\dagger = A'. \end{aligned}$$

Under this, all physical predictions remain unchanged sinc we have that

$$\langle\phi|\psi\rangle \longrightarrow \langle\phi'|\psi'\rangle = \langle\phi|\,U^\dagger U\,|\psi\rangle = \langle\phi|\psi\rangle$$

and

$$\langle\phi|\,A\,|\psi\rangle \longrightarrow \langle\phi'|\,A'\,|\psi'\rangle = \langle\phi|\,U^\dagger U A U^\dagger U\,|\psi\rangle = \langle\phi|\,A\,|\psi\rangle\,.$$

Furthermore we have that

$$C = AB \longrightarrow C' = A'B'$$

and

$$C = [A, B] \longrightarrow C' = [A', B'].$$

We also have that if Q is Hermitian, then Q' is also Hermitian. We also have the implication that

$$Q\,|\psi\rangle = \lambda\,|\psi\rangle \implies Q'\,|\psi'\rangle = \lambda\,|\psi'\rangle$$

with the same eigenvalue λ.

3.2.2 Changing Pictures

In the Schrödinger picture we have that

$$\begin{aligned} &i\hbar\frac{\partial}{\partial t}\,|\psi(t)\rangle = H\,|\psi(t)\rangle \\ \implies &|\psi(t)\rangle = U(t)\,|\psi(0)\rangle \end{aligned}$$

where $U(t) = \exp(-iHt/\hbar)$ is unitary. In fact we have that $U(t)^\dagger = U(-t) = U(t)^{-1}$ and that $U(t_1)U(t_2) = U(t_1 + t_2)$. The states depend on t but observables do not in the Schrödinger

picture. To pass to the Heisenberg picture we shift t-dependence from states to operators as follows. In the Schrödinger picture we have states and operators

$$|\psi(t)\rangle_{\mathrm{S}} \quad A_{\mathrm{S}}.$$

But in the Heisenberg picture we have

$$\begin{aligned} |\psi\rangle_{\mathrm{H}} &= U^{\dagger} |\psi(t)\rangle_{\mathrm{S}} = e^{itH/\hbar} |\psi(t)\rangle_{\mathrm{S}} = |\psi(0)\rangle_{\mathrm{S}} \\ A_{\mathrm{H}}(t) &= U^{\dagger} A_{\mathrm{S}} U = e^{itH/\hbar} A_{\mathrm{S}} e^{-itH/\hbar} \end{aligned}$$

and because the transformation $U(t)$ is unitary we have that

$$\langle \phi(t)| A_{\mathrm{S}} |\psi(t)\rangle_{\mathrm{S}} = \langle \phi| A_{\mathrm{H}}(t) |\psi\rangle_{\mathrm{H}}$$

Note that $H_{\mathrm{H}} = H_{\mathrm{S}} = H$.

To specify dynamics in the Heisenberg picture consider

$$\begin{aligned} \frac{d}{dt} A_{\mathrm{H}}(t) &= \frac{d}{dt} \left(e^{iHt/\hbar} A_{\mathrm{S}} e^{-iHt/\hbar} \right) \\ &= \frac{iH}{\hbar} e^{iHt/\hbar} A_{\mathrm{S}} e^{-iHt/\hbar} - e^{iHt/\hbar} A_{\mathrm{S}} e^{-iHt/\hbar} \frac{iH}{\hbar} \\ &= \frac{i}{\hbar} [H, A_{\mathrm{H}}(t)] \end{aligned}$$

or

$$i\hbar \frac{d}{dt} A_{\mathrm{H}}(t) = [A_{\mathrm{H}}(t), H]$$

which is the Heisenberg equations of motion.

Example. Consider a particle in one dimension $\hat{x}(t), \hat{p}(t)$ in the Heisenberg picture. Consider $H = \frac{\hat{p}^2}{2m} + V(\hat{x})$ so the Heisenberg equations of motion become

$$\frac{d}{dt} \hat{x}(t) = \frac{1}{i\hbar} [\hat{x}(t), H] = \frac{1}{m} \hat{p}(t)$$

and

$$\frac{d}{dt} \hat{p}(t) = \frac{1}{i\hbar} [\hat{p}(t), H] = -V'(\hat{(}t))$$

using $[\hat{x}(t), \hat{p}(t)] = i\hbar$.