

Multivariate Analysis (slides 8)

- Today we consider Linear Discriminant Analysis (LDA) and Quadratic Discriminant Analysis (QDA).
- These are used if it is assumed that there exists a set of k groups within the data and that there is a subset of the data that is labelled, *i.e.*, whose group membership is known.
- Discriminant analysis refers to a set of ‘supervised’ statistical techniques where the class information is used to help reveal the structure of the data.
- This structure then allows the ‘classification’ of future observations.

Discriminant Analysis

- We want to be able to use knowledge of labelled data (*i.e.*, those whose group membership is known) in order to classify the group membership of unlabelled data.
- We previously considered the k -nearest neighbours technique for this problem.
- We shall now consider the alternative approaches of:
 - LDA (linear discriminant analysis)
 - QDA (quadratic discriminant analysis)

LDA & QDA

- Unlike k -Nearest Neighbours (and all the other techniques so far covered), both LDA and QDA assume the use of a distribution over the data.
- Once we introduce distributions (and parameters of those distributions), we can start to quantify uncertainty over the structure of the data.
- As far as classification is concerned, this means that we can start to talk about the probability of group assignment.
- The distinction between a point that is assigned a probability of 0.51 to one group and 0.49 to another, against a point that is assigned a probability of 0.99 to one group and 0.01 to another, can be quite important.

Multivariate Normal Distribution

- Let $\mathbf{x}^T = (x_1, x_2, \dots, x_m)$, where x_1, x_2, \dots, x_m are random variables.
- The Multi-Variate Normal (MVN) distribution has two parameters:
 - Mean μ , an m -dimensional vector.
 - Covariance matrix Σ , with dimension $m \times m$.
- A vector \mathbf{x} is said to follow a MVN distribution, denoted $\mathbf{x} \sim MVN(\mu, \Sigma)$, if it has the following probability density function:

$$f(\mathbf{x}|\mu, \Sigma) = \frac{1}{(2\pi)^{\frac{m}{2}} |\Sigma|^{\frac{1}{2}}} \exp \left[-\frac{1}{2}(\mathbf{x} - \mu)^T \Sigma^{-1}(\mathbf{x} - \mu) \right]$$

- Here $|\Sigma|$ is used to denote the determinant of Σ .

Multivariate Normal Distribution


- The MVN distribution is very useful when modelling multivariate data.
- Notice:

$$\{\mathbf{x} : f(\mathbf{x}|\mu, \Sigma) > C\} = \left\{ \mathbf{x} : (\mathbf{x} - \mu)^T \Sigma^{-1} (\mathbf{x} - \mu) < -2 \log \left[C (2\pi)^{\frac{m}{2}} |\Sigma|^{\frac{1}{2}} \right] \right\}$$

- This corresponds to an m -dimensional ellipsoid centered at point μ .
- If it is assumed that the data within a group k follows a MVN distribution with mean μ_k and covariance Σ_k , then the scatter of the data should be roughly elliptical.
- The mean fixes the location of the scatter and the covariance affects the shape of the ellipsoid.


Normal Contours

- For example, the contour plot of a MVN $\left[\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & 0.8 \\ 0.8 & 3 \end{pmatrix} \right]$ is:





Normal Contours: Data

- Sampling from this distribution and overlaying the results on the contour plot gives:



Shape of Scatter

- If we assume that the data within each group follows a MVN distribution with mean μ_k and covariance Σ_k , then we also assume that the scatter is roughly elliptical.
- The mean sets the location of this scatter and the covariance sets the shape of the ellipse.




Mahalanobis Distance

- The Mahalanobis distance from a point \mathbf{x} to a mean μ is D , where

$$D^2 = (\mathbf{x} - \mu)^T \Sigma^{-1} (\mathbf{x} - \mu).$$

- Two points have the same Mahalanobis distance if they are on the same ellipsoid centered on μ (as defined earlier).



Which Is Closest?

- Suppose we wish to find the mean μ_k that a point \mathbf{x} is closest to as measured by Mahalanobis distance.
- That is, we want to find the k that minimizes the expression:

$$(\mathbf{x} - \mu_k)^T \Sigma_k^{-1} (\mathbf{x} - \mu_k)$$

- The point \mathbf{x} is closer to μ_k than it is to μ_l (under Mahalanobis distance) when:

$$(\mathbf{x} - \mu_k)^T \Sigma_k^{-1} (\mathbf{x} - \mu_k) < (\mathbf{x} - \mu_l)^T \Sigma_l^{-1} (\mathbf{x} - \mu_l).$$

- Note that this is a quadratic expression for \mathbf{x} .

When Covariance is Equal

- If $\Sigma_k = \Sigma$ for all k , then the previous expression becomes:

$$(\mathbf{x} - \mu_k)^T \Sigma^{-1} (\mathbf{x} - \mu_k) < (\mathbf{x} - \mu_l)^T \Sigma^{-1} (\mathbf{x} - \mu_l).$$

- This can be simplified as:

$$-2\mathbf{x}^T \Sigma^{-1} \mu_k + \mu_k^T \Sigma^{-1} \mu_k < -2\mathbf{x}^T \Sigma^{-1} \mu_l + \mu_l^T \Sigma^{-1} \mu_l$$

$$\Leftrightarrow -2\mu_k^T \Sigma^{-1} \mathbf{x} + \mu_k^T \Sigma^{-1} \mu_k < -2\mu_l^T \Sigma^{-1} \mathbf{x} + \mu_l^T \Sigma^{-1} \mu_l$$

- This is now a linear expression for \mathbf{x}
- *Note the names of ‘linear’ discriminant analysis and ‘quadratic’ discriminant analysis.*

Estimating Equal Covariance

- In LDA we need to pool the covariance matrices of individual classes.
- Remember that the sample covariance matrix Q for a set of n observations of dimension m is the matrix whose elements are

$$q_{ij} = \frac{1}{n-1} \sum_{k=1}^n (x_{ki} - \bar{x}_i)(x_{kj} - \bar{x}_j)$$

for $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, m$.

- Then the pooled covariance matrix is defined as:

$$Q_p = \frac{1}{n-g} \sum_{l=1}^g (n_l - 1) Q_l$$

Where g is the number of classes, Q_l is the estimated sample covariance matrix for class l , n_l is the number of data points in class l , whilst n is the total number of data points.

Estimating Equal Covariance

- This formula arises from summing the squares and cross products over data points in all classes:

$$W_{ij} = \sum_{l=1}^g \sum_{k=1}^{n_l} (x_{ki} - \bar{x}_{li})(x_{kj} - \bar{x}_{lj})$$

for $i = 1, \dots, m$ and $j = 1, \dots, m$.

- Hence:

$$W = \sum_{l=1}^g (n_l - 1) Q_l$$

- Given n data points falling in g groups, we have $n - g$ degrees of freedom because we need to estimate the g group means.
- This results in the previous formula for the pooled covariance matrix:

$$Q_p = \frac{W}{n - g}$$

Modelling Assumptions

- Both LDA and QDA are *parametric* statistical methods.
- In order to classify a new observation \mathbf{x} into one of the known K groups, we need to know $\mathbb{P}(\mathbf{x} \in k | \mathbf{x})$ for $k = 1, \dots, K$.
- That is to say, we need to know the posterior probability of belonging to each of the possible groups given the data.
- Then classify the new observation as belonging to the class which has largest posterior probability.
- Bayes' Theorem states that the posterior probability of observation \mathbf{x} belonging to group k is:

$$\mathbb{P}(\mathbf{x} \in k | \mathbf{x}) = \frac{\pi_k f(\mathbf{x}_i | \mathbf{x} \in k)}{\sum_{l=1}^K \pi_l f(\mathbf{x}_i | \mathbf{x} \in l)}$$

Modelling Assumptions

- Discriminant analysis assumes that observations from group k follow a MVN distribution with mean μ_k and covariance Σ_k .

- That is

$$f(\mathbf{x}|\mathbf{x} \in k) = f(\mathbf{x}|\mu_k, \Sigma_k) = \frac{1}{(2\pi)^{\frac{m}{2}} |\Sigma_k|^{\frac{1}{2}}} \exp \left[-\frac{1}{2} (\mathbf{x} - \mu_k)^T \Sigma_k^{-1} (\mathbf{x} - \mu_k) \right]$$

- Discriminant analysis (as presented here) also assumes values for $\pi_k = \mathbb{P}(\mathbf{x} \in k)$, which is the proportion of population objects belonging to class k (this can be known or estimated).
- Note that $\sum_{k=1}^K \pi_k = 1$.
- Typically, $\pi_k = 1/K$ is used.
- π_k are sometimes referred to as prior probabilities.
- Using all this we can compute $\mathbb{P}(\mathbf{x} \in k|\mathbf{x})$ and assign data points to groups so as to maximise this probability.

Some Calculations

- The probability of \mathbf{x} belonging to group k conditional on \mathbf{x} being known satisfies:

$$\mathbb{P}(\mathbf{x} \in k|\mathbf{x}) \propto \pi_k f(\mathbf{x}|\mu_k, \Sigma_k).$$

- Hence,

$$\mathbb{P}(\mathbf{x} \in k|\mathbf{x}) > \mathbb{P}(\mathbf{x} \in l|\mathbf{x}) \Leftrightarrow \pi_k f(\mathbf{x}|\mu_k, \Sigma_k) > \pi_l f(\mathbf{x}|\mu_l, \Sigma_l)$$

- Taking logarithms and substituting in the probability density function for a MVN distribution we find after simplification:

$$\begin{aligned} \log \pi_k - \frac{1}{2} \log |\Sigma_k| - \frac{1}{2}(\mathbf{x} - \mu_k)^T \Sigma_k^{-1}(\mathbf{x} - \mu_k) \\ > \log \pi_l - \frac{1}{2} \log |\Sigma_l| - \frac{1}{2}(\mathbf{x} - \mu_l)^T \Sigma_l^{-1}(\mathbf{x} - \mu_l) \end{aligned}$$

Linear Discriminant Analysis

- If equal covariances are assumed then $\mathbb{P}(\mathbf{x} \in k | \mathbf{x}) > \mathbb{P}(\mathbf{x} \in l | \mathbf{x})$ if and only if:

$$\log \pi_k + \mathbf{x}^T \Sigma^{-1} \mu_k - \frac{1}{2} \mu_k^T \Sigma^{-1} \mu_k > \log \pi_l + \mathbf{x}^T \Sigma^{-1} \mu_l - \frac{1}{2} \mu_l^T \Sigma^{-1} \mu_l$$

- Hence the name linear discriminant analysis.
- If $\pi_k = 1/K$ for all k , then this reduces further.

$$\left(\mathbf{x} - \frac{1}{2}(\mu_k + \mu_l) \right)^T \Sigma^{-1} (\mu_k - \mu_l) > 0$$

Quadratic Discriminant Analysis

- No simplification arises in the unequal covariance case, hence $\mathbb{P}(\mathbf{x} \in k|\mathbf{x}) > \mathbb{P}(\mathbf{x} \in l|\mathbf{x})$ if and only if:

$$\begin{aligned} \log \pi_k - \frac{1}{2} \log |\Sigma_k| - \frac{1}{2}(\mathbf{x} - \mu_k)^T \Sigma_k^{-1}(\mathbf{x} - \mu_k) \\ > \log \pi_l - \frac{1}{2} \log |\Sigma_l| - \frac{1}{2}(\mathbf{x} - \mu_l)^T \Sigma_l^{-1}(\mathbf{x} - \mu_l) \end{aligned}$$

- Hence the name quadratic discriminant analysis.
- If $\pi_k = 1/K$ for all k , then some simplification arises.

Summary

- In LDA the decision boundary between class k and class l is given by:

$$\log \frac{P(k|\mathbf{x})}{P(l|\mathbf{x})} = \log \frac{\pi_k}{\pi_l} + \log \frac{f(\mathbf{x}|k)}{f(\mathbf{x}|l)} = 0$$

- Unlike k -nearest neighbour, both LDA and QDA are model based classifiers where $P(\text{data}|\text{group})$ is assumed to follow a MVN distribution:
 - The model based assumption allows for the generation of the probability for class membership.
 - The MVN assumption means that groups are assumed to follow an elliptical shape.
- Whilst LDA assumes groups have the same covariance matrix, QDA permits different covariance structures between groups.