

A
Same-line control: M = K562 essential − K562 genome-wide (6,207 genes); filled: two-sided p ≤ 0.05
single-line features
EXPLORATORY (post hoc)
K562 − Jurkat
RPE1 − HepG2 †
RPE1 − K562 †
K562 − HepG2
Jurkat − RPE1 †
Jurkat − HepG2
p = 0.0448
p = 0.0194
p = 0.0017
p = 0.169
p = 0.122
p = 0.398
−0.0020 −0.0015 −0.0010 −0.0005 0.0000 0.0005
θ̂ against same-line M
K562 CPM
HepG2 CPM
Jurkat CPM
RPE1 CPM
0.000 0.025 0.050 0.075
partial r with same-line M
B
Signal track: Tn5 insertion CPM (primary) vs MACS2 fold change (FC); filled: two-sided p ≤ 0.05
K562 − Jurkat: CPM (primary)
K562 − Jurkat: FC, all lines
RPE1 − HepG2 †: CPM (primary)
RPE1 − HepG2 †: FC, all lines
RPE1 − HepG2 †: only RPE1 = 4DN FC
RPE1 − K562 †: CPM (primary)
RPE1 − K562 †: only RPE1 = 4DN FC
Jurkat − RPE1 †: CPM (primary)
Jurkat − RPE1 †: only RPE1 = 4DN FC
p = 0.838
p = 0.85
p = 0.0001
p = 0.267
p = 0.161
p = 0.0001
p = 0.668
p = 0.0002
p = 0.238
−0.03 −0.02 −0.01 0.00 0.01
θ̂ (95% CI)
C
EXPLORATORY (post hoc): RPE1 diagnostic, selected variants
RPE1 − K562 (mixed)
Jurkat − RPE1 (mixed)
RPE1 − HepG2 (matched)
primary (reproduced)
RPE1 CN = DepMap clone consensus
RPE1 thinned to Jurkat depth (3 seeds)
exclude chr10 and chrX
exclude both p53 gene sets
promoter feature
distal feature
RPE1 feature alone
partner feature alone
RPE1 = 4DN FC (others CPM)
RPE1 = local FC (others CPM)
partner = FC, RPE1 CPM
RPE1 CPM part shared with FC
RPE1 CPM part not shared with FC
top tercile of RPE1 expression
bottom tercile of RPE1 expression
−0.06 −0.04 −0.02 0.00 0.02
−0.04 −0.02 0.00 0.02
θ̂ (95% CI)
two-sided p ≤ 0.05
p > 0.05
primary θ̂
CN / depth / chromosomes / location / expression
signal-track variants
one side only