
EXPLORATORY (post hoc): RPE1 diagnostic, all variants (filled: two-sided p ≤ 0.05; dotted: primary θ̂)
insertion TSS enrichment RPE1 10.6, other lines 13.7–28.3
primary
copy number
accessibility source
location
chromosomes
p53 / expression
side
two-sided p > 0.05
RPE1 − K562 (mixed)
Jurkat − RPE1 (mixed)
RPE1 − HepG2 (matched)
primary (frozen main test; reproduced)
CN: RPE1 = DepMap clone consensus (features + cn_diff)
CN: DepMap, features only
CN: DepMap, cn_diff covariate only
CN: no CN correction for RPE1
RPE1 CPM from rep1 only
RPE1 CPM from rep2 only
RPE1 = 4DN-processed FC (others CPM)
RPE1 = local-pipeline FC (others CPM)
RPE1 thinned to Jurkat depth (mean of 3 seeds)
RPE1 thinned to 58M insertions (mean of 3 seeds)
all lines FC (frozen FC sensitivity)
all lines FC, RPE1 = 4DN FC + peaks
partner = FC, RPE1 kept CPM
partner and RPE1 = FC (others CPM)
RPE1 CPM part shared with RPE1 FC
RPE1 CPM part not shared with RPE1 FC
promoter feature
distal feature
exclude chr10
exclude chrX
exclude chr10 and chrX
exclude Fischer p53 core targets
exclude HALLMARK_P53_PATHWAY
exclude both p53 sets
bottom tercile of RPE1 expression
middle tercile of RPE1 expression
top tercile of RPE1 expression
RPE1 feature alone (oriented, resid. on C)
partner feature alone (oriented, resid. on C)
−0.06
−0.04
−0.02
0.00
0.02
θ̂ (95% CI)
LOCO: 0/23 drops > 0; jackknife SE 0.0046
−0.06
−0.04
−0.02
0.00
0.02
θ̂ (95% CI)
LOCO: 0/23 drops > 0; jackknife SE 0.0053
−0.04
−0.02
0.00
0.02
θ̂ (95% CI)
LOCO: 0/23 drops > 0; jackknife SE 0.0032