

# Chapter 12: Linear Maps

March 27, 2023

---

## Contents

<b>1</b>	<b>General approach</b>	<b>1</b>
1.1	Definition . . . . .	1
1.2	Notation . . . . .	1
1.3	Specific Linear Maps . . . . .	1
1.3.1	Definition . . . . .	1
1.4	Necessary Condition . . . . .	1
1.4.1	Proof . . . . .	1
<b>2</b>	<b>Kernel and Images</b>	<b>2</b>
2.1	Definition . . . . .	2
2.2	Graphics representation . . . . .	2
2.3	Example . . . . .	2
2.4	Proposition . . . . .	3
<b>3</b>	<b>Projects and Symmetries</b>	<b>3</b>
3.1	Reminder: . . . . .	3
3.1.1	Definition of supplementary subspaces . . . . .	3
3.2	Proposition . . . . .	3
3.3	Definition . . . . .	4
<b>4</b>	<b>Rank Nullity Theorem (RNT)</b>	<b>4</b>
4.1	Definition of Rank . . . . .	4
4.2	Proposition . . . . .	4
4.3	Theorem (Rank Nullity Theorem) . . . . .	4
4.4	Corollary . . . . .	5
4.5	Involvement . . . . .	5
4.6	Proof of corollary . . . . .	5

# 1 General approach

## 1.1 Definition

Let  $E, F$  two  $\mathbb{K} - VS$ , and  $f$  a mapping from  $E$  to  $F$ . We say that  $f$  is a linear (or  $f$  is a linear map) if:

$$\forall(\alpha, X, Y) \in \mathbb{K} \times E \times E, f(\alpha \cdot X + Y) = \alpha \cdot f(X) + f(Y)$$

$$\Longleftrightarrow$$

$$\forall(\alpha, \beta, X, Y) \in \mathbb{K} \times \mathbb{K} \times E \times E, f(\alpha \cdot X + \beta \cdot Y) = \alpha \cdot f(X) + \beta \cdot f(Y)$$

## 1.2 Notation

We denote  $L(E, F)$  the set of all linear maps from  $E$  to  $F$ .

## 1.3 Specific Linear Maps

### 1.3.1 Definition

1. Let  $f \in \mathcal{L}(E, F)$ : we say  $f$  is an endomorphism if  $E = F$  we then denote  $\mathcal{L}(E)$  the set of all endomorphism of  $E$ .
2. Let  $f \in \mathcal{L}(E, F)$ : we say  $f$  is an isomorphism if  $f$  is bijective.
3. Let  $f \in \mathcal{L}(E, F)$ : we say  $f$  is an automorphism if  $f$  is an endomorphism and an isomorphism. ( $E = F$  and bijective)

## 1.4 Necessary Condition

$$f \in \mathcal{L}(E, F) \implies f(0_E) = 0_F$$

### 1.4.1 Proof

<p>Let <math>X \in E</math></p> $f(0_E) = f(0_E \times X)$ $f(0_E) = 0_F \times f(X)$ $f(0_E) = 0_F$	<p>Let <math>X \in E</math></p> $f(0_E) = f(X - X)$ $f(0_E) = f(X) - f(X)$ $f(0_E) = 0_F$
--	---

## 2 Kernel and Images

### 2.1 Definition

Let  $E$  and  $F$  two  $\mathbb{K} - VS$  and  $f \in \mathcal{L}(E, F)$ . Then:

1. We call kernel of  $f$  and denote  $Ker(f)$  the subset of  $E$  defined as follows:


$$Ker(f) = \{X \in E, f(X) = 0_F\} = f^{-1}(\{0_F\})$$

*Note:  $f^{-1}()$  is NOT the inverse of  $f$  because  $f$  is not necessarily bijective.*

2. We call image of  $f$  and denote  $Im(f)$  the subset of  $F$  defined as follows:

$$Im(f) = \{f(X), X \in E\} = \{Y \in F, \exists X \in E, f(X) = Y\}$$

### 2.2 Graphics representation



### 2.3 Example

$$f: \mathbb{R}^2 \longrightarrow \mathbb{R}^3 \quad \text{① } f \in \mathcal{L}(\mathbb{R}^2, \mathbb{R}^3)?$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \longmapsto \begin{pmatrix} x \\ 0 \\ y \end{pmatrix} \quad \text{② } Kerf = ?$$

$$\quad \quad \quad \text{③ } Imf = ?$$

① Necessary condition:  $f(0_E) = 0_F : f(0_{\mathbb{R}^2}) = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} ?$

$$\forall (\alpha, X, Y) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2, X = \begin{pmatrix} x \\ y \end{pmatrix} \text{ and } Y = \begin{pmatrix} x' \\ y' \end{pmatrix}, x, y, x', y' \in \mathbb{R}$$

$$f(\alpha \cdot X + Y) = \begin{pmatrix} \alpha \cdot x + x' \\ \alpha \cdot y + y' \end{pmatrix} = \begin{pmatrix} \alpha \cdot x + x' \\ 0 \\ \alpha \cdot y + y' \end{pmatrix} = \alpha \cdot \begin{pmatrix} x \\ 0 \\ y \end{pmatrix} + \begin{pmatrix} x' \\ 0 \\ y' \end{pmatrix} = \alpha \cdot f(X) + f(Y)$$

so ①  $f \in \mathcal{L}(\mathbb{R}^2, \mathbb{R}^3) \checkmark$

②  $Ker(f) = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^2, \begin{pmatrix} x \\ 0 \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \right\} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$

$$\textcircled{3} \quad \text{Im}(f) = \left\{ \begin{pmatrix} x \\ 0 \\ y \end{pmatrix}, (x, y) \in \mathbb{R}^2 \right\} = \left\{ x \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + y \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, (x, y) \in \mathbb{R}^2 \right\}$$

$$\text{Im}(f) = \text{span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$$

## 2.4 Proposition

1. Let  $f \in \mathcal{L}(E, F)$  and  $g \in \mathcal{L}(F, G)$ . Then:

$$g \circ f \in \mathcal{L}(E, G)$$

2. If  $f$  is bijective then  $f^{-1}$  is bijective and  $f^{-1} \in \mathcal{L}(F, E)$

3.  $\mathcal{L}(E, F)$  is a  $\mathbb{K} - VS$ :

$$\begin{aligned} \mathcal{L}(E, F): \quad E &\longrightarrow F \\ X &\mapsto B_F \in \mathcal{L}(F, E) \end{aligned} \quad \forall (\alpha, f, g) \in \mathbb{K} \times \mathcal{L}^2(E, F)$$

$$\alpha \cdot f + g \in \mathcal{L}(E, F)$$

4. Let  $f \in \mathcal{L}(A, B)$ :

- $f$  is injective  $\iff \text{Ker}(f) = \{0_A\}$
- $f$  is surjective  $\iff \text{Im}(f) = B$

## 3 Projects and Symmetries

### 3.1 Reminder:

#### 3.1.1 Definition of supplementary subspaces

Let  $E$  a  $\mathbb{K} - VS$ . Let  $F$  and  $G$  two supplementary  $\mathbb{K} - VSS$  of  $E$ :

$$E = F \oplus G \iff \forall X \in E, \exists! (X_F, X_G) \in F \times G, X = X_F + X_G$$

### 3.2 Proposition

Let us consider:

$$p: \quad \begin{array}{ccc} E & \longrightarrow & F \subset E \\ & & \underbrace{\hspace{1.5cm}} \\ X & \xrightarrow{p} & X_F \xrightarrow{p} X_F \end{array}$$

1.  $p \in \mathcal{L}(E)$
2.  $p \circ p (= p^2) = p$
3.
  - $\text{Ker}(p) = G$
  - $\text{Im}(p) = F$

### 3.3 Definition

We call projector from  $\mathbb{K}$  - VS  $E$  over  $\mathbb{K}$  - VS  $F$ , any endomorphism  $p$  of  $E$  such that  $p \circ p = p$ . (We often say that  $p$  is a projector over  $F$  parallel of/alongside  $G$ .)


$$\text{Im}(p) \oplus \text{Ker}(p) = E$$

Let us consider:

$$S : \quad E \longrightarrow E$$

$$X \longmapsto (2P - \text{Id}_E)(X) = 2p(x) - X$$

1.  $S \in \mathcal{L}(E)$
2.  $\forall X \in E, S(X) = X_E - X_G$
3.  $S \circ S = \text{Id}_E$



## 4 Rank Nullity Theorem (RNT)

### 4.1 Definition of Rank

Let  $E$  and  $F$  two finite dimensional vector spaces. We call rank of any mapping  $f$  from  $\mathcal{L}(E, F)$  and denote  $\text{rank}(f)$  the dimension of  $\text{Im}(f)$ :

$$\text{rank}(f) = \dim(\text{Im}(f))$$

### 4.2 Proposition

Let  $E$  a finite dimensional  $\mathbb{K}$  - VS such that  $B = (U_1, U_2, \dots, U_n)$  a basis of  $E$ . Let  $F$  a  $\mathbb{K}$  - VS. Then let  $f \in \mathcal{L}(E, F)$ . We have:

$$\text{Im}(f) = \text{sp}(\{f(U_1), f(U_2), \dots, f(U_n)\})$$

We then deduce the following theorem:

### 4.3 Theorem (Rank Nullity Theorem)

Let  $E$  a finite dimensional VS and  $f \in \mathcal{L}(E, F)$ . Then:

$$\dim(\mathbf{E}) = \dim(\text{Ker}(\mathbf{f})) + \overbrace{\text{rank}(\mathbf{f})}^{\dim(\text{Im}(\mathbf{f}))}$$

#### 4.4 Corollary

Let  $f \in \mathcal{L}(E, F)$  where  $E$  and  $F$  two finite dimensional vector spaces.

$$\text{If } \dim(E) = \dim(F) (\iff \dim(\text{Ker}(f)) = 0 \text{ or } \dim(\text{Im}(f)) = 0)$$

$$\implies$$

$$f \text{ injective} \iff f \text{ surjective} \iff f \text{ bijective}$$

#### 4.5 Involvement

$$f \text{ injective} \iff \text{Ker}(f) = \{0_E\}$$

$$[f \text{ injective} \implies \dim(E) \leq \dim(F)] \iff [\dim(E) > \dim(F) \implies f \text{ not injective}]$$

$$f \text{ surjective} \iff \text{Im}(f) = F$$

$$[f \text{ surjective} \implies \dim(E) \geq \dim(F)] \iff [\dim(E) < \dim(F) \implies f \text{ not surjective}]$$

#### 4.6 Proof of corollary

Hypothesis:  $\dim(E) = \dim(F)$

$$f \text{ injective} \iff \text{Ker}(f) = \{0_E\}$$

$$\iff \dim(\text{Ker}(f)) = 0$$

$$\iff \dim(E) = \dim(\text{Ker}(f)) + \dim(\text{Im}(f))$$

$$\iff \dim(E) = \dim(\text{Im}(f))$$

$$\iff \dim(F) = \dim(\text{Im}(f))$$

$$f \text{ surjective} \iff \text{Im}(f) = F$$