

## MATU9D2 : PRACTICAL STATISTICS : FORMULA SHEET

### 1. One Sample Summary Statistics

$$\bar{x} = \frac{\sum x_i}{n}$$
$$s^2 = \frac{\sum x_i^2 - \frac{(\sum x_i)^2}{n}}{n-1}$$

### 2. Two Sample Summary Statistics

$$\bar{x}_1 = \frac{\sum x_{1i}}{n_1} \quad s_1^2 = \frac{\sum x_{1i}^2 - \frac{(\sum x_{1i})^2}{n_1}}{n_1 - 1}$$
$$\bar{x}_2 = \frac{\sum x_{2i}}{n_2} \quad s_2^2 = \frac{\sum x_{2i}^2 - \frac{(\sum x_{2i})^2}{n_2}}{n_2 - 1}$$
$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

### 3. TO COME

#### 4. Relationships

The Chi-squared test has test statistic

$$X^2 = \sum_{all\ cells} \frac{(O_{ij} - E_{ij})^2}{E_{ij}}$$

Using a significance level of 0.05, this is compared with  $\chi^2(df; 0.05)$ , where  $df = (r-1)(c-1)$ .

TO COME

#### 5. Probability

A random variable X following a Binomial probability model with n trials and probability of success  $\theta$  has probability distribution

$$\Pr[X = x] = \frac{n!}{x!(n-x)!} \theta^x (1-\theta)^{n-x}, \quad x = 0, \dots, n.$$

and mean and variance given by

$$E(X) = n\theta, \quad \text{Var}(X) = n\theta(1-\theta).$$

If X follows a Normal probability model with mean  $\mu$  and variance  $\sigma^2$ , then  $Z = \frac{(X-\mu)}{\sigma}$  follows a N(0,1) distribution.

## 6. One Sample Confidence Intervals

An interval estimate for a population *proportion*, having approximate confidence 0.95, is

$$\hat{\theta} \pm 1.96 \times \sqrt{\frac{\hat{\theta}(1-\hat{\theta})}{n}}$$

An interval estimate for a population *mean*, having confidence 0.95, is

$$\bar{x} \pm t(n-1; 0.025) \times \frac{s}{\sqrt{n}}$$

An interval estimate for a population *variance*, having confidence 0.95, is

$$\left( \frac{(n-1)s^2}{\chi^2(n-1; 0.025)}, \frac{(n-1)s^2}{\chi^2(n-1; 0.975)} \right).$$

## 7. Two Sample Tests

The test statistics for the test of the Null Hypothesis  $H_0 : \mu_1 = \mu_2$  against the alternative  $H_1 : \mu_1 \neq \mu_2$  is

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{s_p^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

For a test having significance level 0.05, we compare this with  $\pm t(n_1 + n_2 - 2; 0.025)$ .

The test statistic for the test of the Null Hypothesis  $H_0 : \sigma_1^2 = \sigma_2^2$  against the alternative  $H_1 : \sigma_1^2 \neq \sigma_2^2$  is

$$\frac{s_1^2}{s_2^2} \left( \frac{\text{larger}}{\text{smaller}} \right)$$

For a test having significance level 0.05, we compare this with

$$F(n_1 - 1, n_2 - 1; 0.025)$$

8. TO COME
9. TO COME