

Maths Homework Template

Author

July 9, 2022

Short Math Guide can see <http://tug.ctan.org/info/short-math-guide>, and it's Chinese version here: <https://github.com/hoganbin/short-math-guide>.

1. ¶ Inline equations for simple math: $1 + 1 = 2$.

¶ Centered equations for more involved or important equations:

$$\frac{d}{dx} e^{a(x)} = a'(x) e^{a(x)} \quad (1)$$

$$\mathbf{p} = \hbar \mathbf{k} \quad (2)$$

$$\frac{\|\mathbf{Ax}\|_\infty}{\|\mathbf{x}\|_\infty} = \frac{\max_i \left| \sum_{j=1}^n A_{ij} x_{j1} \right|}{\max_i |x_{i1}|} \quad (3)$$

¶ An example of a matrix:

$$M = \begin{bmatrix} \frac{1 + \sqrt{3}i}{2} & e^{-i\omega t} \\ ie^{-i\omega t} & \frac{1 - \sqrt{3}i}{2} \end{bmatrix} \quad (4)$$

¶ An example of intergals:

$$\iiint \nabla \cdot \mathbf{A} \, dV = \oiint \mathbf{A} \, d\mathbf{S} \quad (5)$$

¶ An example of roots:

$$r = \sqrt[3]{\frac{3}{4\pi N}} \quad (6)$$

- 2.

$$\begin{aligned} I(x) &= \int_{-\frac{b}{2}}^{\frac{b}{2}} dI(x) \\ &= 4\alpha \frac{I_0}{b} \int_{-\frac{b}{2}}^{\frac{b}{2}} \cos^2 \left(\frac{\pi d}{\lambda D} \left(x + \frac{D}{R} \xi \right) \right) d\xi \\ &= 4\alpha I_0 \left(1 + \frac{1}{u} \sin u \cos \left(\frac{2\pi d}{\lambda D} x \right) \right), \quad u = \frac{\pi db}{\lambda R} \end{aligned}$$

- 3.

$$Y_n(y) = \begin{cases} 0, & n = 2k \\ \frac{8a^2}{\pi^3 n^3} \left(1 - \frac{\cosh \frac{n\pi y}{a}}{\cosh \frac{n\pi b}{2a}} \right), & n = 2k + 1 \end{cases}$$

4.

$$\begin{aligned} u(x, t) = & \frac{6Al(-1)^m}{ES(2m+1)^2\pi^2} \sin \frac{2m+1}{2l} \pi x \sin \frac{2m+1}{2l} \pi at \\ & + \frac{2A(-1)^{m+1}}{ES(2m+1)\pi} \left(x \cos \frac{2m+1}{2l} \pi x \sin \frac{2m+1}{2l} \pi at + at \sin \frac{2m+1}{2l} \pi x \cos \frac{2m+1}{2l} \pi at \right) \\ & + \frac{8Al(2m+1)}{ES\pi^2} \sum_{k=0, k \neq m}^{\infty} \frac{(-1)^k}{(2k+1)[(2m+1)^2 - (2k+1)^2]} \sin \frac{2k+1}{2l} \pi x \sin \frac{2k+1}{2l} \pi at \end{aligned}$$

5.

$$\begin{aligned} u(x, t) = & \sum_{n=0}^{\infty} X_n(x) Y_n(y) \\ = & \frac{8a^2}{\pi^3} \sum_{k=0}^{\infty} \frac{1}{(2k+1)^3} \left(1 - \frac{\cosh \frac{2k+1}{a} \pi y}{\cosh \frac{2k+1}{2a} \pi b} \right) \sin \frac{2k+1}{a} \pi x \end{aligned}$$

6.

$$\begin{aligned} \mathcal{L} \left\{ \frac{1 - \cos \omega t}{t} \right\} &= \int_p^{\infty} \left(\frac{1}{p} - \frac{1}{2(p + i\omega)} - \frac{1}{2(p - i\omega)} \right) dt \\ &= \ln p - \frac{1}{2} \ln(p + i\omega) - \frac{1}{2} \ln(p - i\omega) \Big|_p^{\infty} \\ &= \frac{1}{2} \ln \frac{p^2}{p^2 + \omega^2} \Big|_p^{\infty} \\ &= \frac{1}{2} \ln \frac{p^2 + \omega^2}{p^2} \end{aligned}$$

7.

$$\begin{aligned} [q, p]_{Q, P} &= \frac{\partial q}{\partial Q} \cdot \frac{\partial p}{\partial P} - \frac{\partial q}{\partial P} \cdot \frac{\partial p}{\partial Q} \\ &= \left(-\frac{\sqrt{1 - P^2 e^{2Q}}}{e^Q} - \frac{P^2 e^Q}{\sqrt{1 - P^2 e^{2Q}}} \right) \cdot \left(\frac{-e^Q}{\sqrt{1 - P^2 e^{2Q}}} \right) - \left(\frac{-P e^Q}{\sqrt{1 - P^2 e^{2Q}}} \right) \cdot \left(\frac{-P e^Q}{\sqrt{1 - P^2 e^{2Q}}} \right) \\ &= 1 + \frac{P^2 e^{2Q}}{1 - P^2 e^{2Q}} - \frac{P^2 e^{2Q}}{1 - P^2 e^{2Q}} \\ &= 1 \end{aligned}$$

8.

$$\begin{aligned} \frac{4\pi R_{\odot}^2 \cdot \sigma T_{\odot}^4}{4\pi d^2} &= 1352.83 \text{ W/m}^2 \\ T_{\odot} &= 5763.43 \text{ K} \\ \lambda_{\odot} &= \frac{b}{T_{\odot}} = 5027.86 \text{ \AA} \end{aligned}$$