

Per-SNP MR scatter plots — remaining six pairs
a
femoral neck → OA knee
IVW β = +0.058 (P = 0.00064)
MR-Egger β = +0.058
Weighted median β = +0.052
n = 25 instruments
SNP effect on OA (knee)
SNP effect on BMD femoral neck
b
lumbar spine → OA knee
IVW β = +0.095 (P = 7.7e-09)
MR-Egger β = +0.089
Weighted median β = +0.115
n = 21 instruments
SNP effect on OA (knee)
SNP effect on BMD lumbar spine
c
arms → OA knee
IVW β = +0.075 (P = 1.7e-06)
MR-Egger β = +0.075
Weighted median β = +0.026
n = 16 instruments
SNP effect on OA (knee)
SNP effect on BMD arms
d
legs → OA knee
IVW β = +0.060 (P = 2.3e-05)
MR-Egger β = +0.058
Weighted median β = +0.026
n = 24 instruments
SNP effect on OA (knee)
SNP effect on BMD legs
e
OA (knee) → Any fracture
IVW β = +0.027 (P = 0.39)
MR-Egger β = +0.031
Weighted median β = +0.019
n = 69 instruments
SNP effect on Any fracture
SNP effect on OA (knee)
f
OA (knee) → BMD femoral neck
IVW β = +0.042 (P = 0.19)
MR-Egger β = +0.048
Weighted median β = +0.023
n = 69 instruments
SNP effect on BMD femoral neck
SNP effect on OA (knee)