

a
femoral-neck BMD → knee OA (K=25)
MR-PRESSO outlier
IVW β=+0.017 (P=0.001)
MR-Egger β=+0.017
Weighted median β=+0.015
b
lumbar BMD → knee OA (K=21)
MR-PRESSO outlier
IVW β=+0.028 (P=7.7e-09)
MR-Egger β=+0.026
Weighted median β=+0.034
c
arms BMD → knee OA (K=16)
MR-PRESSO outlier
IVW β=+0.022 (P=1.7e-06)
MR-Egger β=+0.022
Weighted median β=+0.008
d
leg BMD → knee OA (K=24)
MR-PRESSO outlier
IVW β=+0.018 (P=2.3e-05)
MR-Egger β=+0.017
Weighted median β=+0.008
e
knee OA → fracture (K=69)
MR-PRESSO outlier
IVW β=+0.092 (P=0.395)
MR-Egger β=+0.106
Weighted median β=+0.063
f
knee OA → femoral-neck BMD (K=69)
IVW β=+0.142 (P=0.190)
MR-Egger β=+0.164
Weighted median β=+0.078
SNP effect on knee OA (log-OR per allele)
SNP effect on leg BMD (SD per allele)
SNP effect on knee OA (log-OR per allele)
SNP effect on knee OA (log-OR per allele)