

a
femoral-neck BMD → knee OA (K=25)
MR-PRESSO outlier
IVW β=+0.017 (P=0.001)
MR-Egger β=+0.017
Weighted median β=+0.015
b
lumbar BMD → knee OA (K=21)
MR-PRESSO outlier
IVW β=+0.028 (P=7.7e-09)
MR-Egger β=+0.026
Weighted median β=+0.034
c
arms BMD → knee OA (K=16)
MR-PRESSO outlier
IVW β=+0.022 (P=1.7e-06)
MR-Egger β=+0.022
Weighted median β=+0.008
d
leg BMD → knee OA (K=24)
MR-PRESSO outlier
IVW β=+0.018 (P=2.3e-05)
MR-Egger β=+0.017
Weighted median β=+0.008
e
knee OA → fracture (K=69)
MR-PRESSO outlier
IVW β=+0.092 (P=0.395)
MR-Egger β=+0.106
Weighted median β=+0.063
f
knee OA → femoral-neck BMD (K=69)
IVW β=+0.142 (P=0.190)
MR-Egger β=+0.164
Weighted median β=+0.078
SNP effect on knee OA (standardised effect per allele)
SNP effect on leg BMD (SD per allele)
SNP effect on knee OA (standardised effect per allele)
SNP effect on knee OA (standardised effect per allele)