

Per-SNP MR scatter plots — remaining six pairs
a
femoral neck → OA knee
IVW β = +0.017 (P = 0.00064)
MR-Egger β = +0.017
Weighted median β = +0.015
n = 25 instruments
SNP effect on OA (knee)
SNP effect on BMD femoral neck
b
lumbar spine → OA knee
IVW β = +0.028 (P = 7.7e-09)
MR-Egger β = +0.026
Weighted median β = +0.037
n = 21 instruments
SNP effect on OA (knee)
SNP effect on BMD lumbar spine
c
arms → OA knee
IVW β = +0.022 (P = 1.7e-06)
MR-Egger β = +0.022
Weighted median β = +0.031
n = 16 instruments
SNP effect on OA (knee)
SNP effect on BMD arms
d
legs → OA knee
IVW β = +0.018 (P = 2.3e-05)
MR-Egger β = +0.017
Weighted median β = +0.025
n = 24 instruments
SNP effect on OA (knee)
SNP effect on BMD legs
e
OA (knee) → Any fracture
IVW β = +0.092 (P = 0.39)
MR-Egger β = +0.106
Weighted median β = +0.118
n = 69 instruments
SNP effect on Any fracture
SNP effect on OA (knee)
f
OA (knee) → BMD femoral neck
IVW β = +0.142 (P = 0.19)
MR-Egger β = +0.164
Weighted median β = +0.113
n = 69 instruments
SNP effect on BMD femoral neck
SNP effect on OA (knee)