

CHAPTER 20

20.1 The integral can be evaluated analytically as,

$$I = \int_1^2 \left(x + \frac{1}{x}\right)^2 dx = \int_1^2 x^2 + 2 + x^{-2} dx = \left[\frac{x^3}{3} + 2x - \frac{1}{x}\right]_1^2 = \frac{2^3}{3} + 2(2) - \frac{1}{2} - \frac{1^3}{3} - 2(1) + \frac{1}{1} = 4.8333$$

The tableau depicting the implementation of Romberg integration to $\varepsilon_s = 0.5\%$ is

| iteration→ | 1 | 2 | 3 |
|-----------------------------|------------|------------|------------|
| $\varepsilon_t \rightarrow$ | 6.0345% | 0.0958% | 0.0028% |
| $\varepsilon_a \rightarrow$ | | 1.4833% | 0.0058% |
| 1 | 5.12500000 | 4.83796296 | 4.83347014 |
| 2 | 4.90972222 | 4.83375094 | |
| 4 | 4.85274376 | | |

Thus, the result is 4.83347014.

20.2 (a) The integral can be evaluated analytically as,

$$I = \left[-0.011x^5 + 0.215x^4 - 1.4x^3 + 3.15x^2 + 2x\right]_0^8 = 20.992$$

(b) The tableau depicting the implementation of Romberg integration to $\varepsilon_s = 0.5\%$ is

| iteration→ | 1 | 2 | 3 | 4 |
|-----------------------------|-------------|-------------|-------------|-------------|
| $\varepsilon_t \rightarrow$ | 87.8049% | 71.5447% | 0.0000% | 0.0000% |
| $\varepsilon_a \rightarrow$ | | 14.2857% | 4.4715% | 0.0000000% |
| 1 | 2.56000000 | 5.97333333 | 20.99200000 | 20.99200000 |
| 2 | 5.12000000 | 20.05333333 | 20.99200000 | |
| 4 | 16.32000000 | 20.93333333 | | |
| 8 | 19.78000000 | | | |

Thus, the result is exact.

(c) The transformations can be computed as

$$x = \frac{(8+0) + (8-0)x_d}{2} = 4 + 4x_d \quad \quad \quad dx = \frac{8-0}{2} dx_d = 4dx_d$$

These can be substituted to yield

$$I = \int_{-1}^1 \left[-0.055(4+4x_d)^4 + 0.86(4+4x_d)^3 - 4.2(4+4x_d)^2 + 6.3(4+4x_d) + 2\right] 4dx_d$$

The transformed function can be evaluated using the values from Table 20.1

$$I = 0.5555556f(-0.774596669) + 0.8888889f(0) + 0.5555556f(0.774596669) = 20.992$$

which is exact.

(d) The following script can be developed and saved as Prob2002Script.m:

PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.

```
format long g
Y = @(x) -0.055*x.^4+0.86*x.^3-4.2*x.^2+6.3*x+2;
I = quad(Y,0,8)
```

When it is run, the result is exact:

```
>> Prob2002Script
I =
                20.992
```

20.3 Although it's not required, the analytical solution can be evaluated simply as

$$I = \int_0^3 x e^{2x} dx = \left[0.25 e^{2x} (2x-1) \right]_0^3 = 504.53599$$

(a) The tableau depicting the implementation of Romberg integration to $\varepsilon_s = 0.5\%$ is

| iteration → | 1 | 2 | 3 | 4 |
|-----------------------------|---------------|--------------|--------------|--------------|
| $\varepsilon_t \rightarrow$ | 259.8216% | 31.8835% | 1.8912% | 0.0312% |
| $\varepsilon_a \rightarrow$ | | 43.2082% | 1.8397% | 0.0290545% |
| 1 | 1815.42957072 | 665.39980101 | 514.07794398 | 504.69324146 |
| 2 | 952.90724344 | 523.53556004 | 504.83987744 | |
| 4 | 630.87848089 | 506.00835760 | | |
| 8 | 537.22588842 | | | |

(b) The transformations can be computed as

$$x = \frac{(3+0) + (3-0)x_d}{2} = 1.5 + 1.5x_d \qquad dx = \frac{3-0}{2} dx_d = 1.5 dx_d$$

These can be substituted to yield

$$I = \int_{-1}^1 \left[(1.5 + 1.5x_d) e^{2(1.5+1.5x_d)} \right] 1.5 dx_d$$

The transformed function can be evaluated using the values from Table 20.1

$$\begin{aligned} I &= f(-0.577350269) + f(0.577350269) \\ f(-0.577350269) &= \left[(1.5 + 1.5(-0.577350269)) e^{2(1.5+1.5(-0.577350269))} \right] 1.5 = 3.379298 \\ f(0.577350269) &= \left[(1.5 + 1.5(0.577350269)) e^{2(1.5+1.5(0.577350269))} \right] 1.5 = 402.9157 \\ I &= 3.379298 + 402.9157 = 406.295 \end{aligned}$$

which represents a percent relative error of 19.47%.

(c) Using MATLAB

```
>> format long g
>> I = quad(@(x) x.*exp(2*x),0,3)
I =
        504.535991881658
```

```
>> I = quadl(@(x) x.*exp(2*x),0,3)
I =
    504.535991866404
```

20.4 The exact solution can be evaluated simply as

```
>> format long
>> erf(1.5)
ans =
    0.96610514647531
```

(a) The transformations can be computed as

$$x = \frac{(1.5+0) + (1.5-0)x_d}{2} = 0.75 + 0.75x_d \quad dx = \frac{1.5-0}{2} dx_d = 0.75dx_d$$

These can be substituted to yield

$$I = \frac{2}{\sqrt{\pi}} \int_{-1}^1 \left[e^{-(0.75+0.75x_d)^2} \right] 0.75dx_d$$

The transformed function can be evaluated using the values from Table 20.1

$$I = f(-0.577350269) + f(0.577350269) = 0.974173129$$

which represents a percent relative error of 0.835 %.

(b) The transformed function can be evaluated using the values from Table 20.1

$$I = 0.5555556f(-0.774596669) + 0.8888889f(0) + 0.5555556f(0.774596669) = 0.965502083$$

which represents a percent relative error of 0.062 %.

20.5 (a) The tableau depicting the implementation of Romberg integration to $\varepsilon_s = 0.5\%$ is

| iteration → | 1 | 2 | 3 | 4 |
|-----------------------------|---------------|---------------|---------------|---------------|
| $\varepsilon_a \rightarrow$ | | 17.8666% | 0.9589% | 0.0382084% |
| 1 | 348.00501404 | 1219.63999486 | 1440.68457469 | 1476.79729373 |
| 2 | 1001.73124965 | 1426.86928845 | 1476.23303250 | |
| 4 | 1320.58477875 | 1473.14779849 | | |
| 8 | 1435.00704356 | | | |

Note that if 8 iterations are implemented, the method converges on a value of 1480.56848.

(b) The transformations can be computed as

$$x = \frac{(30+0) + (30-0)x_d}{2} = 15 + 15x_d \quad dx = \frac{30-0}{2} dx_d = 15dx_d$$

These can be substituted to yield

$$I = 200 \int_{-1}^1 \left[\frac{15 + 15x_d}{5 + 15x_d} e^{-2(15 + 15x_d)/30} \right] 15 dx_d$$

The transformed function can be evaluated using the values from Table 20.1

$$I = f(-0.577350269) + f(0.577350269) = 1610.572$$

(c) Interestingly, the quad function encounters a problem and exceeds the maximum number of iterations

```
>> format long g
>> I = quad(@(z) 200*z/(5+z)*exp(-2*z/30), 0, 30)
Warning: Maximum function count exceeded; singularity likely.
> In quad at 92
I =
    1520.40857847296
```

The quadl function converges rapidly, but does not yield a very accurate result:

```
>> I = quadl(@(z) 200*z/(5+z)*exp(-2*z/30), 0, 30)
I =
    1483.68924281497
```

20.6 The integral to be evaluated is

$$I = \int_0^{1/2} \left(8e^{-t} \sin 2\pi t \right)^2 dt$$

Note that although it is not necessary, the integral can be evaluated analytically to yield

$$I = \left[-16e^{-2t} \frac{1 + 4\pi^2 - \cos(4\pi t) + 2\pi \sin(4\pi t)}{1 + 4\pi^2} \right]_0^{0.5}$$

which can be evaluated as 9.86406915. Therefore, the $I_{RMS} = 3.14071157$.

(a) The tableau depicting the implementation of Romberg integration to $\varepsilon_s = 0.1\%$ is

| iteration → | 1 | 2 | 3 | 4 |
|-----------------------------|------------|-------------|------------|------------|
| $\varepsilon_t \rightarrow$ | 100.0000% | 31.1763% | 1.6064% | 0.0156% |
| $\varepsilon_a \rightarrow$ | | 25.0000% | 2.0824% | 0.0253398% |
| 1 | 0.00000000 | 12.93932074 | 9.70561610 | 9.86561132 |
| 2 | 9.70449056 | 9.90772264 | 9.86311139 | |
| 4 | 9.85691462 | 9.86589959 | | |
| 8 | 9.86365335 | | | |

Therefore, the $I_{RMS} = 3.14095707$.

(b) The transformations can be computed as

$$x = \frac{(0.5 + 0) + (0.5 - 0)x_d}{2} = 0.25 + 0.25x_d \quad dx = \frac{0.5 - 0}{2} dx_d = 0.25 dx_d$$

These can be substituted to yield

$$I = \int_{-1}^1 \left[8e^{-(0.25+0.25x_d)} \sin 2\pi(0.25+0.25x_d) \right]^2 0.25dx_d$$

For the two-point application, the transformed function can be evaluated using the values from Table 20.1

$$I = f(-0.577350269) + f(0.577350269) = 7.678608$$

or an $I_{RMS} = 2.77103$.

For the three-point application, the transformed function can be evaluated using the values from Table 20.1

$$I = 0.5555556f(-0.774596669) + 0.8888889f(0) + 0.5555556f(0.774596669) = 10.02083$$

or an $I_{RMS} = 3.16557$.


(c)

```
>> format long g
>> I = quad(@(x) (8*exp(-x).*sin(2*pi*x)).^2,0,0.5)
I =
    9.86406918193917
```

or an $I_{RMS} = 3.14071157$.

20.7

```
clear,clc,clf
m=1000; DT=[0:300];
H(1)=0;
for i = 2:length(DT)
    H(i)=m*quad(@(T) 0.132+1.56e-4*T+2.64e-7*T.^2,-100,-100+DT(i));
end
plot(DT,H)
title('Heat required versus change in temperature')
xlabel('Change in temperature (C)'),ylabel('Heat required (cal)')
```



20.8 The integral to be evaluated is

$$I = \int_2^8 (9 + 5 \cos^2 0.4t)(5e^{-0.5t} + 2e^{0.15t}) dt$$

(a) The tableau depicting the implementation of Romberg integration to $\varepsilon_s = 0.1\%$ is

| iteration → | 1 | 2 | 3 | 4 |
|-----------------------------|--------------|--------------|--------------|--------------|
| $\varepsilon_a \rightarrow$ | | 8.2537% | 0.1298% | 0.0014429% |
| 1 | 437.99743327 | 329.28470773 | 336.26944122 | 335.95919795 |
| 2 | 356.46288911 | 335.83289538 | 335.96404550 | |
| 4 | 340.99039381 | 335.95584861 | | |
| 8 | 337.21448491 | | | |

(b)

```
>> format long g
>> Qc = @(t) (9+5*cos(0.4*t).^2).*(5*exp(-0.5*t)+2*exp(0.15*t));
>> I=quad(Qc,2,8)
```

```
I =
    335.962530076433
```

20.9 (a) The integral can be evaluated analytically as,

$$\int_{-2}^2 \left[\frac{x^3}{3} - 3y^2x + y^3 \frac{x^2}{2} \right]_0^4 dy$$

$$\int_{-2}^2 \frac{(4)^3}{3} - 3y^2(4) + y^3 \frac{(4)^2}{2} dy$$

$$\int_{-2}^2 21.33333 - 12y^2 + 8y^3 dy$$

$$\left[21.33333y - 4y^3 + 2y^4 \right]_{-2}^2$$

$$21.33333(2) - 4(2)^3 + 2(2)^4 - 21.33333(-2) + 4(-2)^3 - 2(-2)^4 = 21.33333$$

(b) The operation of the `dblquad` function can be understood by invoking `help`,

```
>> help dblquad
```

A session to use the function to perform the double integral can be implemented as,

```
>> dblquad(inline('x.^2-3*y.^2+x*y.^3'),0,4,-2,2)
```

```
ans =
    21.3333
```

20.10

```
>> F=@(x) (1.6*x-0.045*x.^2).*cos(-0.00055*x.^3+0.0123*x.^2+0.13*x);
>> W=quad(F,0,30)
W =
   -157.0871
```

20.11 The integral to be determined is

PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.

$$I = \int_0^{1/2} (6e^{-1.25t} \sin 2\pi t)^2 dt$$

Change of variable:

$$x = \frac{0.5+0}{2} + \frac{0.5-0}{2}x_d = 0.25 + 0.25x_d \quad dx = \frac{0.5-0}{2}dx_d = 0.25dx_d$$

$$I = \int_{-1}^1 (6e^{-1.25(0.25+0.25x_d)} \sin 2\pi(0.25+0.25x_d))^2 0.25 dx_d$$

Therefore, the transformed function is

$$f(x_d) = 0.25 (6e^{-1.25(0.25+0.25x_d)} \sin 2\pi(0.25+0.25x_d))^2$$

Five-point formula:

$$I = 0.236927f(-0.90618) + 0.478629f(-0.53847) + 0.568889f(0) \\ + 0.478629f(0.53847) + 0.236927f(0.90618) = 4.94153$$

Therefore, the RMS current can be computed as

$$I_{\text{RMS}} = \sqrt{4.94153} = 2.222955$$

20.12 (a)

```
>> F1=@(t) (sin(2*pi*t)).^2;
```

```
>> Irms=sqrt(quad(F1,0,1))
```

```
Irms =
```

```
0.7071
```

```
>> R=5; P=Irms^2*R
```

```
P =
```

```
2.5000
```

(b)

```
>> ib=@(t) sin(2*pi*t);
```

```
>> Vb=@(t) 5*ib(t)-1.25*ib(t).^3;
```

```
>> Pb=@(t) ib(t).*Vb(t);
```

```
>> P=quad(Pb,0,1)
```

```
P =
```

```
2.0313
```

The results are different. The reason for this can be seen by inspecting each of the power functions. For (a), the power function is

$$P = I^2 R = 5(\sin 2\pi t)^2$$

For (b), it is

$$P = IV = I(5I - 1.25I^3) = 5I^2 - 6.25I^4$$


$$P = 5(\sin 2\pi t)^2 - 6.25(\sin 2\pi t)^4$$

A plot can be developed of both functions along with their difference.

```

t=linspace(0,1);
Pa=@(t) 5*(sin(2*pi*t)).^2;
P1=Pa(t); P2=Pb(t);
delta=P2-P1;
subplot(2,1,1),plot(t,P1,t,P2,'--')
legend('(a)','(b)'),title('Power versus time')
subplot(2,1,2),plot(t,delta)
title('difference')

```



20.13 The average voltage can be computed as

$$\bar{V} = \frac{\int_0^{60} i(t)R(i) dt}{60}$$

We can use the formulas to generate values of $i(t)$ and $R(i)$ and their product for various equally-spaced times over the integration interval as summarized in the table below. The last column shows the integral of the product as calculated with Simpson's 1/3 rule.

| t | $i(t)$ | $R(i)$ | $i(t) \times R(i)$ | Simpson's 1/3 |
|-------|----------|-----------|--------------------|---------------|
| 0 | 3600.000 | 36469.784 | 131291223 | |
| 6 | 2950.461 | 29916.029 | 88266063 | 1075071847 |
| 12 | 2288.787 | 23235.215 | 53180447 | |
| 18 | 1726.549 | 17553.332 | 30306694 | 381186392 |
| 24 | 1260.625 | 12839.641 | 16185971 | |
| 30 | 878.355 | 8966.984 | 7876196 | 102019847 |
| 36 | 569.294 | 5830.320 | 3319166 | |
| 42 | 327.533 | 3370.360 | 1103904 | 15944048 |
| 48 | 151.215 | 1568.912 | 237242 | |
| 54 | 41.250 | 436.372 | 18000 | 618486 |
| 60 | 0.000 | 0.000 | 0 | |
| Sum → | | | | 1574840619 |


The average voltage can therefore be computed as

PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.

$$\bar{V} = \frac{1,574,840,619}{60} = 2.6247344 \times 10^7$$

20.14

```
clear,clc,clf
t = [0 0.2 0.4 0.6 0.8 1 1.2];
icurr = [0.2 0.3683 0.3819 0.2282 0.0486 0.0082 0.1441];
C=1e-5;
p=polyfit(t,icurr,5);
f= @(t) p(1)*t.^5+p(2)*t.^4+p(3)*t.^3+p(4)*t.^2+p(5)*t+p(6);
t=[0:1.2/100:1.2]; V(1)=0;
for i = 2:length(t)
    V(i)=quad(f,0,t(i))/C;
end
plot(t,V)
```



20.15 The work is computed as the product of the force times the distance, where the latter can be determined by integrating the velocity data,

$$W = F \int_0^t v(t) dt$$

Before solving this problem numerically, it can be solved analytically,

$$W = F \left[\int_0^5 4t dt + \int_5^{15} 20 + (5-t)^2 dt \right] = 200 \left[\left[2t^2 \right]_0^5 + \left[\frac{t^3}{3} - 5t^2 + 45t \right]_5^{15} \right]$$

$$= 200[50 + 533.333] = 200(583.333) = 116,666.7 \text{ N} \cdot \text{m}$$

Romberg integration gives

| ea→ | | 15.0000% | 0.2660% | 0.0127681% | 0.0004501% |
|-----|--------------|--------------|--------------|--------------|--------------|
| 1 | 900.00000000 | 562.50000000 | 587.50000000 | 582.73809524 | 583.41036415 |
| 2 | 646.87500000 | 585.93750000 | 582.81250000 | 583.40773810 | |
| 4 | 601.17187500 | 583.00781250 | 583.39843750 | | |
| 8 | 587.54882813 | 583.37402344 | | | |
| 16 | 584.41772461 | | | | |

Thus, we are converging on the exact result.

Interestingly, the MATLAB quad function does not perform as well. To illustrate this, a function can be developed to hold the integrand

```
function v=velocity(t)
if t <= 5
    v=4*t;
else
    v=20+(5-t).^2;
end
```

A script can then be written to evaluate the integral and compute the work:

```
clear,clc,clf
format long g
Work=200*quad(@velocity,0,15)

Work =
    117062.875572907
```

Thus, the work is computed as 117,062.9 N·m.


20.16 As in the plot, the initial point is assumed to be $e = 0$, $s = 40$. We can then use a combination of the trapezoidal and Simpsons rules to integrate the data as

$$I = (0.02 - 0) \frac{40 + 40}{2} + (0.05 - 0.02) \frac{40 + 37.5}{2} + (0.25 - 0.05) \frac{37.5 + 4(43 + 60) + 2(52) + 55}{12} = 0.8 + 1.1625 + 4.358333 + 5.783333 = 12.10417$$

20.17 The function to be integrated is

$$Q = \int_0^3 2 \left(1 - \frac{r}{r_0} \right)^{1/6} (2\pi r) dr$$

A plot of the integrand can be developed as



As can be seen, the shape of the function indicates that we must use fine segmentation to attain good accuracy. Here are the results of using a variety of segments.

| n | Q | n | Q |
|-----|---------|-------|---------|
| 2 | 25.1896 | 1024 | 44.7289 |
| 4 | 36.1635 | 2048 | 44.7361 |
| 8 | 40.9621 | 4096 | 44.7392 |
| 16 | 43.0705 | 8192 | 44.7407 |
| 32 | 44.0009 | 16384 | 44.7413 |

| | | | |
|-----|---------|--------|---------|
| 64 | 44.4127 | 32768 | 44.7416 |
| 128 | 44.5955 | 65536 | 44.7417 |
| 256 | 44.6767 | 131072 | 44.7418 |
| 512 | 44.7128 | 262144 | 44.7418 |

Therefore, the result to 4 significant figures appears to be 44.7418. The same evaluation can be performed simply with MATLAB

```
>> vA=@(r) 2*(1-r/3).^(1/6)*2*pi.*r;
>> Q=quad(vA,0,3)
Q =
    44.7418
```


20.18 The work is computed as

$$W = k \int_0^x F \, dx$$

The following script fits a 6th-order polynomial to the data and then evaluates the integral of this polynomial with the `quad` function. A plot of the polynomial fit is also displayed.

```
clear,clc,clf
x = [0 0.05 0.1 0.15 0.2 0.25 0.3 0.35];
F = [0 0.01 0.028 0.046 0.063 0.082 0.110 0.130];
k=300;
p=polyfit(x,F,6);
f=@(x) p(1)*x.^6+p(2)*x.^5+p(3)*x.^4+p(4)*x.^3+p(5)*x.^2+p(6)*x+p(7);
xp=linspace(0,max(x),100);
Fp=f(xp);
plot(xp,Fp,x,F,'o'),grid
I=quad(f,0,max(x))
Work=k*I

I =
    0.0202
Work =
    6.0654
```



20.19 The distance traveled is equal to the integral of velocity

$$y = \int_{t_1}^{t_2} v(t) dt$$

A table can be set up holding the velocities at evenly spaced times ($h = 1$) over the integration interval. The Simpson's 1/3 rule can then be used to integrate this data as shown in the last column of the table

| t | v | Simp 1/3 rule |
|--------------|------|-----------------|
| 0 | 0 | |
| 1 | 6 | 19.33333 |
| 2 | 34 | |
| 3 | 84 | 175.3333 |
| 4 | 156 | |
| 5 | 250 | 507.3333 |
| 6 | 366 | |
| 7 | 504 | 1015.333 |
| 8 | 664 | |
| 9 | 846 | 1699.333 |
| 10 | 1050 | |
| 11 | 1045 | 2090 |
| 12 | 1040 | |
| 13 | 1035 | 2070 |
| 14 | 1030 | |
| 15 | 1025 | 2050 |
| 16 | 1020 | |
| 17 | 1015 | 2030 |
| 18 | 1010 | |
| 19 | 1005 | 2010 |
| 20 | 1000 | |
| 21 | 1052 | 2105.333 |
| 22 | 1108 | |
| 23 | 1168 | 2337.333 |
| 24 | 1232 | |
| 25 | 1300 | 2601.333 |
| 26 | 1372 | |
| 27 | 1448 | 2897.333 |
| 28 | 1528 | |
| 29 | 1612 | 3225.333 |
| 30 | 1700 | |
| Sum → | | 26833.33 |

Since the underlying functions are second order or less, this result should be exact. We can verify this by evaluating the integrals analytically,

$$y = \int_0^{10} 11t^2 - 5t dt = \left[3.66667t^3 - 2.5t^2 \right]_0^{10} = 3416.667$$

$$y = \int_{10}^{20} 1100 - 5t dt = \left[1100t - 2.5t^2 \right]_{10}^{20} = 10,250$$

$$y = \int_{20}^{30} 50t + 2(t - 20)^2 dt = \left[\frac{2}{3}t^3 - 15t^2 + 800t \right]_{20}^{30} = 13,166.67$$

The total distance traveled is therefore $3416.667 + 10,250 + 13,166.67 = 26,833.33$.

Interestingly, the MATLAB `quad` function does not perform as well. To illustrate this, a function can be developed to hold the integrand

```
function v=vel(t)
if t <= 10
    v=11*t.^2-5*t;
elseif t <= 20
    v=1100-5*t;
else
    v=50*t+2*(t-20).^2;
end
```

A script can then be written to evaluate the integral and compute the work:

```
clear,clc
format long g
Distance=quad(@vel,0,30)

Distance =
    26835.4501169434
```

20.20 6-segment trapezoidal rule:

$$y = (30-0) \frac{0 + 2(101.439 + 216.213 + 346.916 + 496.983 + 671.095) + 875.867}{2(6)} = 11,352.9$$

6-segment Simpson's 1/3 rule:

$$y = (30-0) \frac{0 + 4(101.439 + 346.916 + 671.095) + 2(216.213 + 496.983) + 875.867}{3(6)} = 11,300.1$$

6-point Gauss quadrature: $y = 11,299.831051$

Romberg integration:

| | 1 | 2 | 3 | 4 |
|-----|--------------------------|-------------|-------------|--------------------|
| n | $\epsilon_a \rightarrow$ | | | |
| 1 | 13138.00101 | 4.0210% | 0.0097% | 0.0000288% |
| 2 | 11772.74279 | 11317.65672 | 11300.04046 | 11299.83245 |
| 4 | 11419.04180 | 11301.14147 | 11299.83570 | |
| 8 | 11329.69844 | 11299.91731 | | |

MATLAB script:

```
clear,clc
format long g
v=@(t) 1850*log(160000./(160000-2500*t))-9.81*t;
y=quad(v,0,30)

y =
    11299.8310550376
```

20.21 (a) Create the following M function:

```
>> y=@(x) 1/sqrt(2*pi)*exp(-(x.^2)/2);
```

PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.

```
>> Q=quad(y,-1,1)
Q =
    0.6827
>> Q=quad(y,-2,2)
Q =
    0.9545
```


Thus, about 68.3% of the area under the curve falls between -1 and 1 and about 95.45% falls between -2 and 2 .

(b) The inflection point is indicated by a zero second derivative. Recall from Chap. 4 (p. 103), that the second derivative can be approximated by

$$f''(x_i) \cong \frac{f(x_{i+1}) - 2f(x_i) + f(x_{i-1}))}{h^2}$$

The following script uses this formula to compute the second derivative and generate a plot of the results,


```
x=-2:.1:2;
y=1/sqrt(2*pi)*exp(-(x.^2)/2);
xx=zeros(length(x)-2);
d2ydx2=xx;
for i=2:length(x)-1
    xx(i-1)=x(i);
    d2ydx2(i-1)=(y(i-1)-2*y(i)+y(i+1))/(x(i)-x(i-1))^2;
end
plot(xx,d2ydx2);grid
```



Thus, inflection points ($d^2y/dx^2 = 0$) occur at -1 and 1 .

Note that in the next chapter we will introduce the `diff` function which provides an alternative way to make the same assessment. Here is a script that illustrates how this might be done:

```
x=-2:.1:2;
f=y(x);
d=diff(f)./diff(x);
xx=-1.95:.1:1.95;
d2=diff(d)./diff(xx);
xxx=-1.9:.1:1.9;
plot(xxx,d2,'o')
```



20.22

| | 1 | 2 | 3 |
|-----|--------------------------|------------|------------|
| n | $\epsilon_a \rightarrow$ | 7.9715% | 0.0997% |
| 1 | 1.34376994 | 1.97282684 | 1.94183605 |
| 2 | 1.81556261 | 1.94377297 | |
| 4 | 1.91172038 | | |

20.23 (a) Romberg:

| | 1 | 2 | 3 | 4 | 5 |
|-----|--------------------------|-----------|-----------|------------|---------------------|
| n | $\epsilon_a \rightarrow$ | 4.2665% | 0.0316% | 0.0001124% | 0.0000000% |
| 1 | 212.75103 | 256.53098 | 255.24166 | 255.26002 | 255.26003094 |
| 2 | 245.58600 | 255.32225 | 255.25974 | 255.26003 | |
| 4 | 252.88818 | 255.26364 | 255.26003 | | |
| 8 | 254.66978 | 255.26025 | | | |
| 16 | 255.11263 | | | | |

(b) MATLAB script:

```
clear,clc
format long g
g=9.81;m=80;cd=0.2;
v=@(t) sqrt(g*m/cd)*tanh(sqrt(g*cd/m)*t);
y=quad(v,0,8)
```

$y =$
255.260030099128

20.24 Equation 20.30 is

$$I = I(h_2) + \frac{1}{15} [I(h_2) - I(h_1)]$$

The integrals can be represented by

$$I(h_1) = (x_4 - x_0) \frac{f(x_0) + 4f(x_2) + f(x_4)}{2}$$

$$I(h_2) = (x_2 - x_0) \frac{f(x_0) + 4f(x_1) + f(x_2)}{6} + (x_4 - x_2) \frac{f(x_2) + 4f(x_3) + f(x_4)}{6}$$

Substituting these into Eq. (20.30) gives

$$I = (x_2 - x_0) \frac{f(x_0) + 4f(x_1) + f(x_2)}{6} + (x_4 - x_2) \frac{f(x_2) + 4f(x_3) + f(x_4)}{6} \\ + \frac{1}{15} \left[(x_2 - x_0) \frac{f(x_0) + 4f(x_1) + f(x_2)}{6} + (x_4 - x_2) \frac{f(x_2) + 4f(x_3) + f(x_4)}{6} - (x_4 - x_0) \frac{f(x_0) + 4f(x_2) + f(x_4)}{6} \right]$$

Note that if $h = x_4 - x_0$, $x_2 - x_0 = x_4 - x_2 = h/2$. Therefore,

$$I = \frac{h}{2} \frac{f(x_0) + 4f(x_1) + f(x_2)}{6} + \frac{h}{2} \frac{f(x_2) + 4f(x_3) + f(x_4)}{6} \\ + \frac{1}{15} \left[\frac{h}{2} \frac{f(x_0) + 4f(x_1) + f(x_2)}{6} + \frac{h}{2} \frac{f(x_2) + 4f(x_3) + f(x_4)}{6} - h \frac{f(x_0) + 4f(x_2) + f(x_4)}{6} \right]$$

Collecting terms

$$\frac{I}{h} = \frac{1}{12} f(x_0) + \frac{1}{15} \frac{1}{12} f(x_0) - \frac{1}{15} \frac{1}{6} f(x_0) + \frac{4}{12} f(x_1) + 4 \frac{1}{15} \frac{1}{12} f(x_1) + \frac{1}{12} f(x_2) + \frac{1}{12} f(x_2) + \frac{1}{15} \frac{1}{12} f(x_2) \\ + \frac{1}{15} \frac{1}{12} f(x_2) - 4 \frac{1}{15} \frac{1}{6} f(x_2) + 4 \frac{1}{15} \frac{1}{12} f(x_3) + \frac{4}{12} f(x_3) + \frac{1}{12} f(x_4) + \frac{1}{15} \frac{1}{12} f(x_4) - \frac{1}{15} \frac{1}{6} f(x_4)$$

or

$$\frac{I}{h} = 0.0777778f(x_0) + 0.3555556f(x_1) + 0.1333333f(x_2) + 0.3555556f(x_3) + 0.0777778f(x_4)$$

Multiplying by $90h$ gives Boole's rule


$$I = h \frac{7f(x_0) + 32f(x_1) + 12f(x_2) + 32f(x_3) + 7f(x_4)}{90}$$

20.25 Here is a script to solve the problem:

```
clear,clc,clf
format short g
r =[0 1100 1500 2450 3400 3630 4500 5380 6060 6280 6380];
rho = [13 12.4 12 11.2 9.7 5.7 5.2 4.7 3.6 3.4 3];
rp=[min(r):max(r)];
rhop=interp1(r,rho,rp,'pchip');
plot(rp,rhop)
rp=rp*1e3; %convert km to meters
rhop=rhop*1e6/1e3; %convert g/cm3 to kg/m3
Area=4*pi*rp.^2;
rpp=Area.*rhop;
Mass=trapz(rp,rpp)
```

When it is run, the result is

```
Mass =
    6.0905e+024
```

20.26

```
function [q,ea,iter]=romberg(func,a,b,es,maxit,varargin)
% romberg: Romberg integration quadrature
% q = romberg(func,a,b,es,maxit,p1,p2,...):
% Romberg integration.
% input:
% func = name of function to be integrated
% a, b = integration limits
% es = desired relative error (default = 0.000001%)
% maxit = maximum allowable iterations (default = 30)
% p1,p2,... = additional parameters used by func
% output:
% q = integral estimate
% ea = approximate relative error (%)
% iter = number of iterations
if nargin<3,error('at least 3 input arguments required'),end
if nargin<4|isempty(es), es=0.000001;end
if nargin<5|isempty(maxit), maxit=50;end
n = 1;
I(1,1) = trap(func,a,b,n,varargin{:});
iter = 0;
while iter<maxit
    iter = iter+1;
    n = 2^iter;
    I(iter+1,1) = trap(func,a,b,n,varargin{:});
    for k = 2:iter+1
        j = 2+iter-k;
        I(j,k) = (4^(k-1)*I(j+1,k-1)-I(j,k-1))/(4^(k-1)-1);
    end
    ea = abs((I(1,iter+1)-I(2,iter))/I(1,iter+1))*100;
    if ea<=es, break; end
end
q = I(1,iter+1);
```

Script to solve Example 20.1:

```
clear,clc
format long g
fx=@(x) 0.2+25*x-200*x^2+675*x^3-900*x^4+400*x^5;
[q,ea,iter]=romberg(fx,0,0.8)
```

Output:

```
q =
    1.640533333333334
ea =
    0
iter =
    3
```

Script to solve Prob 20.1:

PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.

```
clear,clc
format long g
fx=@(x) (x+1/x)^2;
[q,ea,iter]=romberg(fx,1,2,0.5)
```

Output:

```
q =
    4.833470143613
ea =
    0.00580951575223314
iter =
    2
```

20.27

```
function q = quadadapt(f,a,b,tol,varargin)
% Evaluates definite integral of f(x) from a to b
if nargin < 4 | isempty(tol),tol = 1.e-6;end
c = (a + b)/2;
fa = feval(f,a,varargin{:});
fc = feval(f,c,varargin{:});
fb = feval(f,b,varargin{:});
q = quadstep(f, a, b, tol, fa, fc, fb, varargin{:});
end

function q = quadstep(f,a,b,tol,fa,fc,fb,varargin)
% Recursive subfunction used by quadadapt.
h = b - a; c = (a + b)/2;
fd = feval(f,(a+c)/2,varargin{:});
fe = feval(f,(c+b)/2,varargin{:});
q1 = h/6 * (fa + 4*fc + fb);
q2 = h/12 * (fa + 4*fd + 2*fc + 4*fe + fb);
if abs(q2 - q1) <= tol
    q = q2 + (q2 - q1)/15;
else
    qa = quadstep(f, a, c, tol, fa, fd, fc, varargin{:});
    qb = quadstep(f, c, b, tol, fc, fe, fb, varargin{:});
    q = qa + qb;
end
end
```

Script to solve Example 20.1:

```
clear,clc
format long g
fx=@(x) 0.2+25*x-200*x^2+675*x^3-900*x^4+400*x^5;
q = quadadapt(fx,0,0.8)
```

Output:

```
q =
    1.640533333333334
```

Script to solve Prob 20.1:

```
clear,clc
format long g
fx=@(x) (x+1/x)^2;
[q,ea,iter]=romberg(fx,1,2,0.5)
```

Output:

$q =$
11299.8310550333