

# Group Homomorphism

Hengfeng Wei

hfwei@nju.edu.cn

Mar 21, 2017

# Group Homomorphism

1 Groups of Small Orders

2 Homomorphism

# Order of 4

# Order of 6

# Order of 8 (TJ 9.11)

# Group Homomorphism

1 Groups of Small Orders

2 Homomorphism

$$D_6 \cong D_3 \times \mathbb{Z}_2 \text{ (TJ 9.16)}$$

## (TJ 9.23)

$$G \times K \cong H \times K \not\Rightarrow G \cong H$$

$$G = \mathbb{Z}, \quad H = 1, \quad K = \prod_{n \in \mathbb{N}} \mathbb{Z}$$

“On Cancellation in Groups” by R. Hirshon, 1969

$$G \times K \cong H \times K \quad |K| < \infty \implies G \cong H$$



## (TJ 11.18)

- ▶  $\phi : G_1 \rightarrow G_2$
- ▶  $H_1 \triangleleft G_1$
- ▶  $\phi(H_1) = H_2$
- ▶  $G_1/H_1 \cong G_2/H_2$

$$G_1 = \mathbb{Z}_2 \quad G_2 = \{e\} \quad H_1 = \{0\} \quad H_2 = \{e\}$$

## (TJ 11.5)

$$\phi : \mathbb{Z}_{24} \rightarrow \mathbb{Z}_{18}$$

$$\begin{aligned}\phi(1) = a &\implies \phi(x) = ax \pmod{18} \\ &\implies \text{Ker}(\phi) = \mathbb{Z}_b \cap \mathbb{Z}_{24} : ab \equiv 0 \pmod{18}\end{aligned}$$

$$\phi(1) = ?$$

$$|\phi(1)||18 \wedge |\phi(1)||24 \implies |\phi(1)||6$$

$$\phi(1) = 0, 9, 6, 12, 3, 15$$

$$\phi_3(x) = 6x, \quad \text{Ker}(\phi_3) = 3\mathbb{Z}_{24}$$

# Normal subgroups

$$\mathbb{Z}/n\mathbb{Z}$$

$$(aH)(bH) = (abH)$$

$$\forall a, b \in G, \forall h_1, h_2 \in H, (ah_1) \in aH, (bh_2) \in bH : (ah_1)(bh_2) \in abH$$

$$\exists h_3 \in H, (ah_1)(bh_2) = abh_3 \iff h_1b = bh_3h_2^{-1}$$

$$\forall b \in G, \forall h_1 \in H : \exists h' \in H : h_1b = bh'$$

$$\forall g \in G, \forall h \in H : \exists h' \in H : hg = gh'$$

# Normal subgroups

$$\forall g \in G, \forall h \in H : \exists h' \in H : hg = gh'$$

$$\forall g \in G, Hg \subseteq gH$$

$$\forall g^{-1} \in G, Hg^{-1} \subseteq g^{-1}H$$

$$\forall h \in H, \exists h' \in H : hg^{-1} = g^{-1}h' \iff gh = h'g$$

$$\forall g \in G, Hg \subseteq gH$$

$$\forall g \in G, Hg = gH$$

## (TJ 10.13)

$$g \in G, C(g) = \{x \in G : gx = xg\}$$

$$Z(G) = \{x \in G : gx = xg, \forall g \in G\} \implies Z(G) \triangleleft G$$

$$C(g) \leq G$$

$$\langle g \rangle \triangleleft G \implies C(g) \triangleleft G$$

Proof.

$$\forall k \in G, x \in C(g) : k^{-1}xk \in C(g)$$

$$\forall k \in G, x \in C(g) : gk^{-1}xk = k^{-1}xkg$$

$$\langle g \rangle \triangleleft G \implies k\langle g \rangle = \langle g \rangle k \implies \exists t, gk = kg^t \iff gk^{-1} = k^{-1}g^t$$