

# CSE250A HW2

Name: Guanghao Chen

PID: A53276390

Section: A

## 2.1 Probabilistic inference

(a)

$$\begin{aligned}P(A = 1) &= P(A = 1, E = 0, B = 0) + P(A = 1, E = 0, B = 1) + P(A = 1, E = 1, B = 0) + P(A = 1, E = 1, B = 1) \\&= P(A = 1|E = 0, B = 0)P(E = 0)P(B = 0) + P(A = 1|E = 0, B = 1)P(E = 0)P(B = 1) \\&\quad + P(A = 1|E = 1, B = 0)P(E = 1)P(B = 0) + P(A = 1|E = 1, B = 1)P(E = 1)P(B = 1) \\&= 0.0025\end{aligned}$$

$$P(A = 0) = 1 - P(A = 1) = 0.9975$$

$$\begin{aligned}P(J = 1) &= P(J = 1, A = 0) + P(J = 1, A = 1) \\&= P(J = 1|A = 0)P(A = 0) + P(J = 1|A = 1)P(A = 1) \\&= 0.0521\end{aligned}$$

$$\begin{aligned}P(A = 0|E = 1) &= P(A = 0|B = 0, E = 1)P(B = 0) + P(A = 0|B = 1, E = 1)P(B = 1) \\&= (1 - 0.29)(1 - 0.001) + (1 - 0.95)0.001 \\&= 0.7093\end{aligned}$$

$$P(A = 1|E = 1) = 0.2907$$

Therefore,

$$\begin{aligned}P(E = 1|J = 1) &= \frac{0.7093 \times 0.002}{0.9975} \cdot \frac{0.05 \times 0.9975}{0.0521} + \frac{0.2907 \times 0.002}{0.0025} \cdot \frac{0.09 \times 0.0025}{0.0521} \\&= 0.0116\end{aligned}$$

(b)

$$\begin{aligned}P(E = 1|J = 1, B = 1) &= \frac{P(J = 1|B = 1, E = 1)P(E = 1|B = 1)}{P(J = 1|B = 1)} \\&= \frac{\sum_a P(J = 1, A = a|B = 1, E = 1)P(E = 1)}{\sum_{a,e} P(J = 1, A = a, E = e|B = 1)} \quad (\text{marginalization})\end{aligned}$$

Numerator:

$$\begin{aligned}&\sum_a P(J = 1, A = a|B = 1, E = 1)P(E = 1) \\&= P(E = 1) \times [P(J = 1|A = 0, B = 1, E = 1)P(A = 0|B = 1, E = 1) + P(J = 1|A = 1, B = 1, E = 1)P(A = 1|B = 1, E = 1)] \\&= P(E = 1) \times [P(J = 1|A = 0)P(A = 0|B = 1, E = 1) + P(J = 1|A = 1)P(A = 1|B = 1, E = 1)] \\&= 0.002 \times 0.8575 \\&= 0.0017\end{aligned}$$

Denominator:

$$\begin{aligned}
& \sum_{a,e} P(J = 1, A = a, E = e | B = 1) \\
&= P(J = 1, A = 0, E = 0 | B = 1) + P(J = 1, A = 0, E = 1 | B = 1) \\
&+ P(J = 1, A = 1, E = 0 | B = 1) + P(J = 1, A = 1, E = 1 | B = 1) \\
&= P(J = 1 | A = 0, E = 0, B = 1)P(A = 0 | E = 0, B = 1)P(E = 0) \\
&+ P(J = 1 | A = 1, E = 0, B = 1)P(A = 1 | E = 0, B = 1)P(E = 0) \\
&+ P(J = 1 | A = 0, E = 1, B = 1)P(A = 0 | E = 1, B = 1)P(E = 1) \\
&+ P(J = 1 | A = 1, E = 1, B = 1)P(A = 1 | E = 1, B = 1)P(E = 1) \\
&= P(J = 1 | A = 0)P(A = 0 | E = 0, B = 1)P(E = 0) \\
&+ P(J = 1 | A = 1)P(A = 1 | E = 0, B = 1)P(E = 0) \\
&+ P(J = 1 | A = 0)P(A = 0 | E = 1, B = 1)P(E = 1) \\
&+ P(J = 1 | A = 1)P(A = 1 | E = 1, B = 1)P(E = 1) \\
&= 0.8490
\end{aligned}$$

Therefore,

$$P(E = 1 | J = 1, B = 1) = \frac{0.0017}{0.8490} = 0.0020$$

**(c)**

$$\begin{aligned}
P(A = 1 | M = 0) &= \frac{P(M = 0 | A = 1)P(A = 1)}{P(M = 0)} \\
&= \frac{(1 - P(M = 1 | A = 1))P(A = 1)}{P(M = 0)} \\
&= \frac{0.3 \times 0.0025}{0.9883} \\
&= 0.00075887
\end{aligned}$$

**(d)**

$$\begin{aligned}
P(A = 1 | J = 0, M = 0) &= \frac{P(M = 0, J = 0 | A = 1)P(A = 1)}{P(J = 0, M = 0)} \\
&= \frac{P(J = 0 | A = 1)P(M = 0 | A = 1)P(A = 1)}{\sum_a P(J = 0, M = 0, A = a)} \\
&= \frac{0.1 \times 0.3 \times 0.0025}{0.95 \times 0.99 \times 0.9975 + 0.1 \times 0.3 \times 0.0025} \\
&= 0.00008
\end{aligned}$$

**(e)**

$$\begin{aligned}
P(M = 0) &= P(M = 0, A = 0) + P(M = 0, A = 1) \\
&= P(M = 0 | A = 0)P(A = 0) + P(M = 0 | A = 1)P(A = 1) \\
&= (1 - 0.01)0.9975 + (1 - 0.7)0.0025 \\
&= 0.9883
\end{aligned}$$

$$P(M = 1) = 0.0117$$

$$\begin{aligned}
P(A = 1 | M = 1) &= \frac{P(M = 1 | A = 1)P(A = 1)}{P(M = 1)} \\
&= \frac{0.70 \times 0.0025}{0.0117} \\
&= 0.1496
\end{aligned}$$

**(f)**

$$\begin{aligned}
P(A = 1|M = 1, B = 0) &= \frac{P(M = 1, B = 0|A = 1)P(A = 1)}{P(M = 1, B = 0)} \\
&= \frac{P(M = 1|A = 1)P(B = 0|A = 1)P(A = 1)}{\sum_a P(M = 1, B = 0|A = a)P(A = a)} \\
&= \frac{P(M = 1|A = 1)P(B = 0|A = 1)P(A = 1)}{P(M = 1|A = 0)P(B = 0|A = 0)P(A = 0) + P(M = 1|A = 1)P(B = 0|A = 1)P(A = 1)} \\
P(B = 0|A = 0) &= P(B = 0, E = 0|A = 0) + P(B = 0, E = 1|A = 0) \\
&= \frac{P(A = 0|B = 0, E = 0)P(B = 0)P(E = 0)}{P(A = 0)} + \frac{P(A = 0|B = 0, E = 1)P(B = 0)P(E = 1)}{P(A = 0)} \\
&= \frac{(1 - 0.001)(1 - 0.001)(1 - 0.002)}{0.9975} + \frac{(1 - 0.29)(1 - 0.001)0.002}{0.9975} \\
&= 0.999923 \\
P(B = 0|A = 1) &= \frac{(0.001)(1 - 0.001)(1 - 0.002)}{0.0025} + \frac{(0.29)(1 - 0.001)0.002}{0.0025} \\
&= 0.6306
\end{aligned}$$

Therefore,

$$P(A = 1|M = 1, B = 0) = 0.09962$$

Therefore,

Compared the result of (b) versus (a), it seems consistent with the commonsense pattern of reasoning, which means the prob will be less when John calls.

Compared the result of (d) versus (c), the result is consistent with the commonsense pattern, which means the prob will be less when John makes calls.

Compared the result of (f) versus (e), the result is consistent with the commonsense pattern, which means the prob will be less when actually there's no burglar.

## 2.2 Probabilistic reasoning

### (a)

Numerator of  $r_k$ :

$$\begin{aligned}
P(D = 1|S_1 = 1, S_2 = 1, \dots, S_k = 1) &= \frac{P(S_1 = 1, S_2 = 1, \dots, S_k = 1|D = 1)}{P(S_1 = 1, S_2 = 1, \dots, S_k = 1)} \quad (CI2) \\
&= \frac{\prod_{k=1} P(S_k = 1|D = 1)}{P(S_1 = 1, S_2 = 1, \dots, S_k = 1)} \\
&= \frac{1 \times \frac{f(1)}{f(2)} \times \frac{f(2)}{f(3)} \times \dots \times \frac{f(k-1)}{f(k)}}{P(S_1 = 1, S_2 = 1, \dots, S_k = 1)} \\
&= \frac{\frac{f(1)}{f(k)}}{P(S_1 = 1, S_2 = 1, \dots, S_k = 1)} \\
&= \frac{1}{\frac{2^k + (-1)^k}{2^k}} \\
&= \frac{2^k}{2^k + (-1)^k}
\end{aligned}$$

Denominator of  $r_k$ :

$$\begin{aligned}
 P(D = 0 | S_1 = 1, S_2 = 1, \dots, S_k = 1) &= \frac{P(S_1 = 1, S_2 = 1, \dots, S_k = 1 | D = 0)}{P(S_1 = 1, S_2 = 1, \dots, S_k = 1)} \quad (CI2) \\
 &= \frac{P(S_1 = 1 | D = 0) P(S_2 = 1 | D = 0) \dots P(S_k = 1 | D = 0)}{P(S_1 = 1, S_2 = 1, \dots, S_k = 1)} \\
 &= \frac{\frac{1}{2^k}}{P(S_1 = 1, S_2 = 1, \dots, S_k = 1)}
 \end{aligned}$$

Then

$$\begin{aligned}
 r_k &= \frac{\frac{1}{2^k + (-1)^k}}{\frac{1}{2^k}} \\
 &= \frac{2^k}{2^k + (-1)^k}
 \end{aligned}$$

Therefore, when  $k$  is an odd integer,  $r_k = \frac{2^k}{2^k - 1} > 1$ , and the patient will be diagnose as  $D = 1$  form.

When  $k$  is an even integer,  $r_k = \frac{2^k}{2^k + 1} < 1$ , and the patient will be diagnose as  $D = 0$  form.,

## (b)

As  $k$  values becomes larger, it means  $k \rightarrow \infty$ . Then

When  $k$  is an odd number,  $r_k = \frac{1}{1 - \frac{1}{2^k}}$ .

Then  $r_k$  will be 1.


Likewise,

$r_k$  will also be 1 when  $k$  is an even number.

Therefore, the result will be less certain when the  $k$  goes up.

## 2.3 Sigmoid function

### (a)



$$\begin{aligned}\sigma^{-1}(z) &= \frac{0 \times (1 + e^{-z}) + 1 \times (e^{-z})}{(1 + e^{-z})^2} \\ &= \frac{e^{-z}}{(1 + e^{-z})^2}\end{aligned}$$

At the same time

$$\begin{aligned}\sigma(z) \times \sigma(-z) &= \frac{1}{1 + e^{-z}} \times \frac{1}{1 + e^z} \\ &= \frac{1}{1 + e^{-z}} \times \frac{e^{-z}}{e^{-z} + 1} \\ &= \frac{e^{-z}}{(1 + e^{-z})^2}\end{aligned}$$

Therefore, it concludes that  $\sigma^{-1}(z) = \sigma(z) \times \sigma(-z)$

**(b)**

$$\begin{aligned}\sigma(z) + \sigma(-z) &= \frac{1}{1 + e^{-z}} + \frac{1}{1 + e^z} \\ &= \frac{1}{1 + e^{-z}} + \frac{e^{-z}}{e^{-z} + 1} \\ &= \frac{1 + e^{-z}}{1 + e^{-z}} = 1\end{aligned}$$

**(c)**

$$\begin{aligned}L(\sigma(z)) &= \log\left(\frac{\sigma(z)}{1 - \sigma(z)}\right) \\ &= \log\left(\frac{\frac{1}{1 + e^{-z}}}{1 - \frac{1}{1 + e^{-z}}}\right) \\ &= \log\left(\frac{1}{1 + e^{-z} - 1}\right) \\ &= \log\left(\frac{1}{e^{-z}}\right) \\ &= \log(e^z) = z\end{aligned}$$

**(d)**

Since  $p_i = P(Y = 1 | X_i, X_j = 0 \text{ for all } j \neq i)$ ,

Therefore  $p_i = \sigma(w_i)$ .

Furthur,


$L(p_i) = L(\sigma(w_i)) = w_i$  (according to the conclusion in problem (c))

## 2.4 Conditional independence

---

X	Y	Z
month	water	{sprinkler,rain}
month	water	{sprinkler,rain,accident}
month	accident	{sprinkler,water}
month	accident	{rain,water}
month	accident	{sprinkler,rain}
month	accident	{water}
month	accident	{sprinkler,rain,water}
sprinkler	rain	{month}
sprinkler	accident	{water}
sprinkler	accident	{water, month}
sprinkler	accident	{water, rain}
sprinkler	accident	{water, month, rain}
rain	accident	{water}
rain	accident	{water, month}
rain	accident	{water, rain}
rain	accident	{water, month, rain}

## 2.5 Markov blanket



Proof:

There're 5 cases for node Y.

**Case1: Node Y is the parent of X's parent**

For example, node 1 meets this case. The right top gray node is the evidence. Therefore, d-separation case 1 is satisfied, where left top gray node is flowed through. (node1  $\Rightarrow$  evidence  $\Rightarrow$  nodeX)

#### Case2: Node Y is the child of X's parent

For example, node 5 meets this case. The right top gray node is the evidence. Therefore, d-separation case 2 is satisfied, where right top gray node is diversified. (node5  $\Leftarrow$  evidence  $\Rightarrow$  nodeX)

#### Case3: Node Y is the parent of X's child's parent

For example, node 2 meets this case. There's a path satisfying conditional independence case1, where an evidence node is flowed through. (node2  $\Rightarrow$  **nodex's child's parent**  $\Rightarrow$  nodex's child  $\Rightarrow$  nodeX)

#### Case4: Node Y is the child of X's child's parent

On the contrary, node 3 is the child of node x's child's parent. So there's a path satisfying conditional independence case2, where an evidence node is diversified. (node2  $\Leftarrow$  **nodex's child's parent**  $\Rightarrow$  nodex's child  $\Rightarrow$  nodeX)

#### Case5: Node Y is the child of X's child

Node 4 is the child of node x's child. So there's a path satisfying conditional independence case1, where an evidence node is flowed through. (nodex  $\Rightarrow$  **nodex's child**  $\Rightarrow$  node4)

All in all, no matter node Y meets which case, it must have  $P(X, Y|BX) = P(X|B_X)P(Y|B_X)$ .

## 2.6 True or false

Value	Formula
True	$P(C, D A) = P(A S)$
False	$P(A D) = P(A B, D)$
True	$P(C, E) = P(C)P(E)$
False	$P(C, D, E) = P(C)P(D)P(E)$
False	$P(F, G) = P(F)P(G)$
True	$P(F, G D) = P(F D)P(G D)$
False	$P(A, D, G) = P(A)P(D A)P(G D)$
False	$P(B E) = P(B E, G)$
False	$P(C E) = P(C E, G)$

## 2.7 Subsets

### (a)

$$P(A|C) = P(A|C, E, F, B, D, G) \quad S = \{C, E, F, B, D, G\}$$

### (b)

$$P(C) = P(C|B, D, G) \quad S = \{B, D, G\}$$

### (c)

$$P(C|A) = P(C|A, B, D, G) \quad S = \{A, B, D, G\}$$

**(d)**

$$P(C|A, E) = P(C|A, E, D, B, G) \quad S = \{A, E, D, B, G\}$$

**(e)**

$$P(C|A, E, F) = P(C|A, E, F) \quad S = \{A, E, F\}$$

**(f)**

$$P(C|A, D, E, F) = P(C|A, D, E, F, G) \quad S = \{A, D, E, F, G\}$$

**(g)**

$$P(F) = P(F) \quad S = \{\}$$

**(h)**

$$P(F|C) = P(F|C, E) \quad S = \{C, E\}$$

**(i)**

$$P(F|C, D) = P(F|C, D, E, G) \quad S = \{C, D, E, G\}$$

**(j)**

$$P(B, G) = P(B, G|A, C, E) \quad S = \{A, C, E\}$$

## 2.8 Noisy-OR

---

**(a)**

$$left = 1 - 1 = 0$$

$$right = 1 - (1 - p_y) = p_y$$

$$\therefore left < right$$

**(b)**

$$left = 1 - (1 - p_x) = p_x$$

$$right = 1 - (1 - p_y) = p_y$$

$$\therefore left < right$$

**(c)**

$$left = 1 - (1 - p_x) = p_x$$

$$right = 1 - (1 - p_x)(1 - p_y) = (p_x + p_y - p_x p_y)$$

$$right - left = p_y - p_x p_y = p_y(1 - p_x) > 0$$

$$\therefore left < right$$



**(d)**

$left = right$  (d-separation III)

**(e)**

$left < right$

**(f)**

$left > right$

**(g)**

$$\begin{aligned} right &= P(X = 1, Y = 1, Z = 1) \\ &= P(Z = 1 | X = 1, Y = 1)P(X = 1, Y = 1) \\ &= P(Z = 1 | X = 1, Y = 1)P(X = 1)P(Y = 1) \end{aligned}$$

$$left = P(Z = 1)P(X = 1)P(Y = 1)$$

Compared to no evidence, it is more likely to make  $Z = 1$  when  $X = 1, Y = 1$ .

Therefore,  $left < right$ .

## 2.9 Polytree inference

---

**(a)**

$$P(B) = \sum_a P(B|A = a)P(A = a)$$

**(b)**

$$\begin{aligned} P(D|C) &= \sum_b P(D|B = b, C)P(B = b|C) \\ &= \sum_b P(D|B = b, C)P(B = b) \end{aligned}$$

Insert the result from (a).

$$P(D|C) = \sum_b P(D|B = b, C) \left[ \sum_a P(B|A = a)P(A = a) \right]$$

**(c)**

$$\begin{aligned} P(F|C, E) &= \sum_d P(F, D = d|C, E) \\ &= \sum_d P(F|C, D = d, E)P(D = d|C, E) \\ &= \sum_d P(F|D = d, E) \left\{ \sum_b P(D|B = b, C) \left[ \sum_a P(B|A = a)P(A = a) \right] \right\} \end{aligned}$$

**(d)**

$$\begin{aligned} P(E|C, F) &= \frac{P(F|C, E)P(E|C)}{P(F|C)} \\ &= \frac{P(F|C, E)P(E)}{P(F|C)} \end{aligned}$$

$$\begin{aligned}
P(F|C) &= \sum_e P(F, E = e|C) \\
&= \sum_e P(F|C, E = e)P(E = e|C) \\
&= \sum_e P(F|C, E = e)P(E = e)
\end{aligned}$$

Therefore,

$$P(E|C, F) = \frac{\sum_d \{P(F|D=d, E)\} \sum_b \{P(D|B=b, C) [\sum_a \{P(B|A=a)P(A=a)\}]\}}{\sum_d \{P(F|D=d, E)\} \sum_b \{P(D|B=b, C) [\sum_a \{P(B|A=a)P(A=a)\}]\}} \times P(E) \sum_e \{ \sum_d \{P(F|D=d, E)\} \sum_b \{P(D|B=b, C) [\sum_a \{P(B|A=a)P(A=a)\}]\} P(E=e) \}$$

**(e)**

In order to calculate the result of  $P(E|C, F)$ , we need to traverse n possible values for variable E to calculate  $P(F|C)$ . Then in order to calculate  $P(F|C)$ , we need to traverse n possible values for variable D in term  $P(F|C, E)$ . Further, we need to traverse n possible values for variable B in calculating  $P(D = d|C)$ . Finally, we need to traverse n possibilities for variable A. The whole process is recursive from the top to the bottom.

All in all, the calculation complexity is polynomial and the degree is 4.