

Photometric Stereo

Computer Vision I CSE252A Lecture 8

CS252A, Fall 2017

Computer Vision I

Announcement

- Read Chapter 2 of Forsyth & Ponce
- Office hours tomorrow:
 - 3-5, CSE 4127
 - 5-6, CSE B260A
- Piazza
- Next lecture

CS252A, Fall 2017

Computer Vision I


Shape-from-X

- Where X is
 - Shading
 - Photometric stereo
 - Stereo
 - Motion
 - Texture
 - Blur
 - Focus
 - Structured light
 -

CS252A, Fall 2017

Computer Vision I

Shading reveals 3-D surface geometry



CS252A, Fall 2017

Computer Vision I


Two shape-from-X methods that use shading

- Shape-from-shading: Use just one image to recover shape. Requires knowledge of light source direction and BRDF everywhere. Too restrictive to be useful without priors/learning.
- Photometric stereo: Single viewpoint, multiple images under different lighting.
 1. Arbitrary known BRDF
 2. Lambertian BRDF, known lighting
 3. Lambertian BRDF, unknown lighting.

CS252A, Fall 2017

Computer Vision I

Two shape-from-X methods that use shading



- Photometric stereo: Single viewpoint, multiple images under different lighting.
 1. Arbitrary known BRDF
 2. Lambertian BRDF, known lighting
 3. Lambertian BRDF, unknown lighting.

CS252A, Fall 2017

Computer Vision I

Photometric Stereo Rigs: One viewpoint, changing lighting



Because of single viewpoint, a pixel location sees the same point of the scene across all images

CS252A, Fall 2017

Computer Vision I

An example of photometric stereo



CS252A, Fall 2017

Computer Vision I

Multi-view stereo vs. Photometric Stereo: Assumptions

- Multi-view (binocular) Stereo
 - Multiple images
 - Dynamic scene
 - Multiple viewpoints
 - Fixed lighting
- Photometric Stereo
 - Multiple images
 - Static scene
 - Fixed viewpoint
 - Multiple lighting conditions



CS252A, Fall 2017

Computer Vision I

Photometric Stereo:


1. General BRDF and Reflectance Map
2. Lambertian Surface, known lighting
3. Lambertian surface

CS252A, Fall 2017

Computer Vision I

BRDF

- Bi-directional Reflectance Distribution Function
 $\rho(\theta_{in}, \phi_{in}; \theta_{out}, \phi_{out})$
- Function of
 - Incoming light direction:
 θ_{in}, ϕ_{in}
 - Outgoing light direction:
 θ_{out}, ϕ_{out}
- Ratio of incident irradiance to emitted radiance









CS252A, Fall 2017


Computer Vision I

1. General BRDF and Reflectance Maps

CS252A, Fall 2017

Computer Vision I








Three Source Photometric stereo:
Step1

Offline:
Using source directions & BRDF, construct reflectance map for each light source direction. $R_1(p,q)$, $R_2(p,q)$, $R_3(p,q)$
This can be done analytically or by taking images of spheres

Online:

- Acquire three images with the known light source directions. $E_1(x,y)$, $E_2(x,y)$, $E_3(x,y)$
- For each pixel location (x,y) , find (p,q) as the intersection of the three curves
 $R_1(p,q)=E_1(x,y)$
 $R_2(p,q)=E_2(x,y)$
 $R_3(p,q)=E_3(x,y)$
- The estimated (p,q) gives the surface normal at pixel (x,y) . Over image, the normal field $[p(x,y), q(x,y)]$ is estimated

CS252A, Fall 2017 Computer Vision I



Step 2: Recovering the surface $f(x,y)$


Many methods: Simplest approach

Assumption: Surface and derivative are continuous

- From estimated $\mathbf{n}=(n_x, n_y, n_z)$, $p=n_x/n_z$, $q=n_y/n_z$
- Integrate $df/dx=p$ along a row $(x,0)$ to get $f(x,0)$
- Then integrate $df/dy=q$ along each column starting with value of the first row

CS252A, Fall 2017 Computer Vision I

What might go wrong?



- Height $z(x,y)$ is obtained by integration along a curve from (x_0, y_0) .

$$z(x, y) = z(x_0, y_0) + \int_{(x_0, y_0)}^{(x, y)} (pdx + qdy)$$

- If one integrates the derivative field along any closed curve, one expects to get back to the starting value.
- Might not happen because of noisy estimates of (p,q)

CS252A, Fall 2017

Computer Vision I

What might go wrong?

Integrability. If $f(x,y)$ is the height function, we expect that

$$\frac{\partial}{\partial y} \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} \frac{\partial f}{\partial y}$$



In terms of estimated gradient (p,q) , this means:

$$\frac{\partial p}{\partial y} = \frac{\partial q}{\partial x}$$

where $p=n_x/n_z$, $q=n_y/n_z$ with $\mathbf{n}=[n_x \ n_y \ n_z]$

But N (and in turn p,q) are estimated independently at each, equality is not going to exactly hold

Computer Vision I

Horn's Method

["Robot Vision, B.K.P. Horn, 1986]

- Formulate estimation of surface height $z(x,y)$ from gradient field by minimizing cost functional:
- $$\iint_{\text{Image}} (z_x - p)^2 + (z_y - q)^2 dx dy$$
- where (p,q) are estimated components of the gradient while z_x and z_y are partial derivatives of best fit surface
- Solved using calculus of variations – iterative updating
 - $z(x,y)$ can be discrete or represented in terms of basis functions.
 - Integrability is naturally satisfied.

CS252A, Fall 2017


Computer Vision I

II. Photometric Stereo: Lambertian Surface, Known Lighting

CS252A, Fall 2017

Computer Vision I

Lambertian Surface



At image location (u,v) , the intensity of a pixel $x(u,v)$ is:

$$\begin{aligned} e(u,v) &= [a(u,v) \hat{n}(u,v)] \cdot [\hat{s}_0 \hat{s}] \\ &= b(u,v) \cdot s \end{aligned}$$

where

- $a(u,v)$ is the albedo of the surface projecting to (u,v) .
- $n(u,v)$ is the direction of the surface normal.
- s_0 is the light source intensity.
- s is the direction to the light source.

CS252A, Fall 2017

Computer Vision I

Lambertian Photometric stereo

- If the light sources s_1 , s_2 , and s_3 are known, then we can recover b from as few as three images. (Photometric Stereo: Silver 80, Woodham81).

$$[e_1 \ e_2 \ e_3] = \mathbf{b}^T [s_1 \ s_2 \ s_3]$$

- i.e., we measure e_1 , e_2 , and e_3 and we know s_1 , s_2 , and s_3 . We can then solve for \mathbf{b} by solving a linear system.
- $\mathbf{b}^T = [e_1 \ e_2 \ e_3] [s_1 \ s_2 \ s_3]^{-1}$
- Normal is: $\mathbf{n} = \mathbf{b}/\|\mathbf{b}\|$, albedo is: $|\mathbf{b}|$

CS252A, Fall 2017

Computer Vision I

What if we have more than 3 Images?

Linear Least Squares

$[e_1 \ e_2 \ e_3 \dots e_n] =$ Let the residual be

$$\mathbf{b}^T [\mathbf{s}_1 \ \mathbf{s}_2 \ \mathbf{s}_3 \dots \mathbf{s}_n]$$

Squaring this:

$$r^2 = r^T r = (\mathbf{e} - \mathbf{S}\mathbf{b})^T (\mathbf{e} - \mathbf{S}\mathbf{b}) \\ = \mathbf{e}^T \mathbf{e} - 2\mathbf{b}^T \mathbf{S}^T \mathbf{e} + \mathbf{b}^T \mathbf{S}^T \mathbf{S} \mathbf{b}$$

Rewrite as

$$\mathbf{e} = \mathbf{S}\mathbf{b}$$

where

\mathbf{e} is n by 1

\mathbf{b} is 3 by 1

\mathbf{S} is n by 3


Solving for \mathbf{b} gives

$$\mathbf{b} = (\mathbf{S}^T \mathbf{S})^{-1} \mathbf{S}^T \mathbf{e}$$

CS252A, Fall 2017

Computer Vision I


Input Images



CS252A, Fall 2017


Computer Vision I

Recovered albedo



Computer Vision I


Recovered normal field



CS252A, Fall 2017


Computer Vision I

Surface recovered by integration



Computer Vision I

Lambertian Photometric Stereo



CS252A, Fall 2017

Reconstruction with Albedo Map

Computer Vision I


Without the albedo map



CS252A, Fall 2017

Computer Vision I


Another person



CS252A, Fall 2017

Computer Vision I

No Albedo map



CS252A, Fall 2017

Computer Vision I

III. Photometric Stereo with unknown lighting and Lambertian surfaces


Covered in Illumination cone slides

What is the set of images of an object under all possible lighting conditions?

In answering this question, we'll arrive at a photometric stereo method for reconstructing surface shape w/ unknown lighting.

CSE252A

The Space of Images



- Consider an n -pixel image to be a point in an n -dimensional space, $\mathbf{x} \in \mathbb{R}^n$.
- Each pixel value is a coordinate of \mathbf{x} .
- Many results will apply to linear transformations of image space (e.g. filtered images)
- Other image representations (e.g. Cayley-Klein spaces, See Koenderink's "pixel f#@king paper")

CSE252A

Model for Image Formation

Lambertian Assumption with shadowing:

$$x = \max(\mathbf{B} \mathbf{s}, \mathbf{0}) \quad \mathbf{B} = \begin{bmatrix} -\mathbf{b}_1^T \\ -\mathbf{b}_2^T \\ \dots \\ -\mathbf{b}_n^T \end{bmatrix}_{n \times 3}$$

where

- \mathbf{x} is an n -pixel image vector
- \mathbf{B} is a matrix whose rows are unit normals scaled by the albedos
- $\mathbf{s} \in \mathbb{R}^3$ is vector of the light source direction scaled by intensity

3-D Linear subspace

The set of images of a Lambertian surface with no shadowing is a subset of 3-D linear subspace.

[Moses 93], [Nayar, Murase 96], [Shashua 97]

$$\mathcal{L} = \{\mathbf{x} \mid \mathbf{x} = \mathbf{B}\mathbf{s}, \forall \mathbf{s} \in \mathbb{R}^3\}$$

where \mathbf{B} is a n by 3 matrix whose rows are product of the surface normal and Lambertian albedo

CSE252A

Still Life

Original Images

Basis Images

CSE252A

Rendering Images: $\sum_i \max(\mathbf{B}\mathbf{s}_i, \mathbf{0})$

1 Light	2 Lights	3 Lights

CSE252A

How do you construct subspace?

$[\mathbf{x}_1 \mathbf{x}_2 \mathbf{x}_3] \rightarrow \mathbf{B}$

- Any three images w/o shadows taken under different lighting span \mathcal{L}
- Not orthogonal
- Orthogonalize with Gram-Schmidt

CSE252A

How do you construct subspace?

$[\mathbf{x}_1 \mathbf{x}_2 \mathbf{x}_3 \mathbf{x}_4 \mathbf{x}_5] \rightarrow \mathbf{B}$

With more than three images, perform least squares estimation of \mathbf{B} using Singular Value Decomposition (SVD)

CSE252A

Matrix Decompositions

- Definition: The factorization of a matrix \mathbf{M} into two or more matrices $\mathbf{M}_1, \mathbf{M}_2, \dots, \mathbf{M}_n$, such that $\mathbf{M} = \mathbf{M}_1 \mathbf{M}_2 \dots \mathbf{M}_n$.
- Many decompositions exist...
 - **QR** Decomposition
 - **LU** Decomposition
 - **LDU** Decomposition
 - Etc.

CSE252A

Singular Value Decomposition

Excellent ref: "Matrix Computations," Golub, Van Loan

- Any m by n matrix \mathbf{A} may be factored such that $\mathbf{A} = \mathbf{U}\Sigma\mathbf{V}^T$
- $[\mathbf{m} \times \mathbf{n}] = [\mathbf{m} \times \mathbf{m}][\mathbf{m} \times \mathbf{n}][\mathbf{n} \times \mathbf{n}]$
- \mathbf{U} : m by m , orthogonal matrix
 - Columns of \mathbf{U} are the eigenvectors of $\mathbf{A}\mathbf{A}^T$
- \mathbf{V} : n by n , orthogonal matrix,
 - columns are the eigenvectors of $\mathbf{A}^T\mathbf{A}$
- Σ : m by n , diagonal with non-negative entries ($\sigma_1, \sigma_2, \dots, \sigma_s$) with $s=\min(m,n)$ are called the singular values
 - Singular values are the square roots of eigenvalues of both $\mathbf{A}\mathbf{A}^T$ and $\mathbf{A}^T\mathbf{A}$
 - Result of SVD algorithm: $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_s$

CSE252A

SVD Properties

- In Matlab, $[\mathbf{u} \ \mathbf{s} \ \mathbf{v}] = \text{svd}(\mathbf{A})$, and you can verify that: $\mathbf{A} = \mathbf{u}\mathbf{s}\mathbf{v}^T$
- $r = \text{Rank}(\mathbf{A}) = \#$ of non-zero singular values.
- \mathbf{U}, \mathbf{V} give us orthonormal bases for the subspaces of \mathbf{A} :
 - 1st r columns of \mathbf{U} : Column space of \mathbf{A}
 - Last $m - r$ columns of \mathbf{U} : Left nullspace of \mathbf{A}
 - 1st r columns of \mathbf{V} : Row space of \mathbf{A}
 - last $n - r$ columns of \mathbf{V} : Nullspace of \mathbf{A}
- For $d \leq r$, the first d column of \mathbf{U} provide the best d -dimensional basis for columns of \mathbf{A} in least squares sense.

CSE252A

Thin SVD

- Any m by n matrix \mathbf{A} may be factored such that $\mathbf{A} = \mathbf{U}\Sigma\mathbf{V}^T$
- If $m > n$, then one can view Σ as:

$$\begin{bmatrix} \Sigma \\ 0 \end{bmatrix}$$
- Where $\Sigma' = \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_s)$ with $s = \min(m, n)$, and lower matrix is $(n-m \times m)$ of zeros.
- Alternatively, you can write:

$$\mathbf{A} = \mathbf{U}'\Sigma'\mathbf{V}'^T$$
- In Matlab, thin SVD is: $[\mathbf{U} \ \mathbf{S} \ \mathbf{V}] = \text{svds}(\mathbf{A})$

CSE252A


Estimating \mathbf{B} with SVD

1. Construct data matrix







$$\mathbf{D} = [x_1 \ x_2 \ x_3 \dots \ x_n]$$
2. $[\mathbf{U} \ \mathbf{S} \ \mathbf{V}] = \text{svds}(\mathbf{D})$
 - If data had no noise, then $\text{rank}(\mathbf{D})=3$, and the first three singular values (\mathbf{S}) would be positive and rest would be zero.
 - Take first three column of \mathbf{U} as \mathbf{B} .

CSE252A


Face Basis



CSE252A



Some natural ideas & questions




- Can the cones of two different objects intersect?
- Can two different objects have the same cone?
- How big is the cone?
- How can cone be used for recognition?

CSE252A

Do Ambiguities Exist?

Can two objects of differing shapes produce the same illumination cone?



CSE252A

Do Ambiguities Exist? Yes

- Cone is determined by linear subspace L
- The columns of B span L
- For any $A \in GL(3)$, $B^* = BA$ also spans L .
- For any image of B produced with light source S , the same image can be produced by lighting B^* with $S^* = A^{-1}S$ because


$$X = B^*S^* = BAA^{-1}S = BS$$
- When we estimate B using SVD, the rows are NOT generally normal * albedo.
-

CSE252A

Surface Integrability

In general, B^* does not have a corresponding surface.

Linear transformations of the surface normals in general do not produce an integrable normal field.




CSE252A

GBR Transformation


Only Generalized Bas-Relief transformations satisfy the integrability constraint:

$$A = G^T = \begin{bmatrix} \lambda & 0 & -\mu \\ 0 & \lambda & -\nu \\ 0 & 0 & 1 \end{bmatrix}$$



CSE252A

Generalized Bas-Relief Transformations



Objects differing by a GBR have the same illumination cone.

Without knowledge of light source location, one can only recover surfaces up to GBR transformations.

CSE252A

Uncalibrated photometric stereo


1. Take n images as input, perform SVD to compute B^* .
2. Find some A such that B^*A is close to integrable.
3. Integrate resulting gradient field to obtain height function $f^*(x,y)$.

Comments:

- $f^*(x,y)$ differs from $f(x,y)$ by a GBR.
- Can use specularities to resolve GBR for non-Lambertian surface.

CSE252A

What about cast shadows for nonconvex objects?




P.P. Reubens in Opticorum Libri Sex, 1613
CSE252A

GBR Preserves Shadows

Given a surface f and a GBR transformed surface f' then for every light source s which illuminates f there exists a light source s' which illuminates f' such that the **attached** and **cast shadows** are identical.

GBR is the **only** transform that preserves shadows.

[Kriegman, Belhumeur 2001]



CSE252A

Bas-Relief Sculpture



CSE252A

Codex Urbinas



As far as light and shade are concerned low relief fails both as sculpture and as painting, because the shadows correspond to the low nature of the relief, as for example in the shadows of foreshortened objects, which will not exhibit the depth of those in painting or in sculpture in the round.

Leonardo da Vinci
Treatise on Painting
(Kemp)

CSE252A