

Reflectance, Lights

Computer Vision I
CSE 252A
Lecture 7

CS252A, Fall 2017

Computer Vision I

Computer Vision I

What I learned about this week



- Large Synoptic Survey Telescope
- 3.2Gigapixel camera
- 189 CCD's, each with 16 megapixels
- Pixels are $10\mu\text{m}$
- Filter wheel
- 6200 lbs
- <https://www.lsst.org/about/camera>

CS252A, Fall 2017

Computer Vision I

What I learned about this week

Portrait mode on the Pixel 2 and Pixel 2 XL smartphones

Tuesday, October 17, 2017

Posted by Marc Levoy, Principal Engineer and Yael Pritch, Software Engineer

Portrait mode, a new feature of the new Pixel 2 and Pixel 2 XL smartphones, allows anyone to take professional-looking shallow depth-of-field images. This feature helped both devices earn DxO's highest mobile camera ranking, and works with both the rear-facing and front-facing cameras, even though neither is dual-camera (normally required to obtain this effect). Today we discuss the machine learning and computational photography techniques behind this feature.



<https://research.googleblog.com/2017/10/portrait-mode-on-pixel-2-and-pixel-2-xl.html>

CS252A, Fall 2017

Computer Vision I

Announcements

- HW1 due Friday
- Some point around HW1

CS252A, Fall 2017

Computer Vision I

Projective Transformation (Homography)

- 3×3 linear transformation of homogenous coordinates
- Points map to points,
- lines map to lines

$$\begin{bmatrix} u_1 \\ u_2 \\ u_3 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

Special cases

- Affine transformation: $[a_{31} \ a_{32} \ a_{33}] = [0 \ 0 \ 1]$
- Rigid: Affine and upper-left 2×2 submatrix is orthogonal

CS252A, Fall 2017

Computer Vision I

HW1: Warping by Homography

$$\mathbf{t} = \mathbf{H}\mathbf{s}$$

s: source image coordinates
t: target image coordinates



CS252A, Fall 2017

Computer Vision I


BRDF

With assumptions in previous slide

- Bi-directional Reflectance Distribution Function
 $\rho(\theta_{in}, \phi_{in}; \theta_{out}, \phi_{out})$
- Ratio of emitted radiance to incident irradiance (units: sr⁻¹)
- In local coordinate system at x
 - Incoming light direction:
 θ_{in}, ϕ_{in}
 - Outgoing light direction:
 θ_{out}, ϕ_{out}

$$\rho(x; \theta_{in}, \phi_{in}; \theta_{out}, \phi_{out}) = \frac{L_o(x; \theta_{out}, \phi_{out})}{L_i(x; \theta_{in}, \phi_{in}) \cos \theta_{in} d\omega_i}$$


Where ρ is sometimes denoted f_r .



Computer Vision I

The Reflection Equation

Emitted radiance in direction f_r for incident radiance L_i .



$$L_r(x, \omega_r) = \int_{H^2} f_r(x, \omega_i \rightarrow \omega_r) L_i(x, \omega_i) \cos \theta_i d\omega_i$$

CS348B Lecture 10

where $\omega_i = (\theta_i, \phi_i)$

Pat Hanrahan, Spring 2002

Properties of BRDFs

1. Non-negative: $\rho(\theta_{in}, \phi_{in}; \theta_{out}, \phi_{out}) \geq 0$
2. Helmholtz Reciprocity Principle:
 $\rho(\theta_{in}, \phi_{in}; \theta_{out}, \phi_{out}) = \rho(\theta_{out}, \phi_{out}; \theta_{in}, \phi_{in})$
3. Total energy leaving a surface must be less than total energy arriving at the surface


$$\int_{\Omega_i} L_i(x, \theta_i, \phi_i) \cos \theta_i d\omega_i \geq \int_{\Omega_o} \left[\int_{\Omega_i} \rho(\theta_i, \phi_i; \theta_o, \phi_o) L_i(x, \theta_i, \phi_i) \cos \theta_i d\omega_i \right] \cos \theta_o d\omega_o$$

CS252A, Fall 2017

Computer Vision I

Important class of BRDFs: Isotropic BRDF

$$\rho(\theta_i, \phi_i; \theta_o, \phi_o) = \rho_r(\theta_i, \theta_o, \phi_i - \phi_o)$$



Isotropic BRDF's are symmetric about the surface normal. If the surface is rotated about the normal for the same incident and emitting directions, the value of the BRDF is the same.

CS252A, Fall 2017

Computer Vision I

Anisotropic BRDF

$\rho(\theta_i, \phi_i; \theta_o, \phi_o)$ is an arbitrary function




CS252A, Fall 2017

Computer Vision I

Lambertian (Diffuse) Surface

- BRDF is a constant called the albedo. $\rho(x; \theta_{in}, \phi_{in}; \theta_{out}, \phi_{out}) = K$
- Emitted radiance is NOT a function of outgoing direction – i.e. constant in all directions.
- For lighting coming in single direction ω_i , emitted radiance is proportional to cosine of the angle between normal and light direction

$$L_r = K \hat{N} \cdot \omega_i$$




$$L_r(x, \omega_r) = \int_{H^2} f_r(x, \omega_i \rightarrow \omega_r) L_i(x, \omega_i) \cos \theta_i d\omega_i$$

CS252A, Fall 2017

Computer Vision I

Lambertian reflection




Light emitted equally in all directions.

CS252A, Fall 2017

Computer Vision I

Specular Reflection: Smooth Surface




- $\mathbf{N}, \omega_i, \omega_o$ are coplanar
- $\theta_i = \theta_o$

Speculum – Latin for “Mirror”

CS252A, Fall 2017

Computer Vision I


Rough Specular Surface



CS252A, Fall 2017

Computer Vision I

Phong Model



CS348B Lecture 10

Pat Hanrahan, Spring 2002

Non-Lambertian reflectance



CS252A, Fall 2017


Computer Vision I

Lots of BRDF Models

Common Models

- Lambertian
- Phong
- Physics-based
 - Specular [Blinn 1977], [Cook-Torrance 1982], [Ward 1992]
 - Diffuse [Hanrahan, Kreuger 1993]
 - Generalized Lambertian [Oren, Nayar 1995]
 - Thoroughly Pitted Surfaces [Koenderink et al 1999]
- Phenomenological
 - [Koenderink, Van Doorn 1996]

Arbitrary Reflectance



- Non-parametric model
- Anisotropic
- Non-uniform over surface
- BRDF Measurement [Dana et al, 1999], [Marschner]

Specialized

- Hair, skin, threads, paper [Jensen et al]

CS252A, Fall 2017

Computer Vision I

General BRDF: e.g. Velvet



Portrait of Sir Thomas More, Hans Holbein the Younger, 1527

[After Koenderink et al, 1998]

CSE 252A, Fall 2017

Computer Vision I


Ways to measure BRDFs

CSE 252A, Fall 2017

Computer Vision I

Gonioreflectometers


- Three degrees of freedom spread among light source, detector, and/or sample



-CSE 252A

Gonioreflectometers

- Three degrees of freedom spread among light source, detector, and/or sample



-CSE 252A

Gonioreflectometers

- Can add fourth degree of freedom to measure anisotropic BRDFs



-CSE 252A

BRDF Not Always Appropriate



BRDF



BSSRDF

<http://graphics.stanford.edu/papers/bssrdf/>
(Jensen, Marschner, Levoy, Hanrahan)

Light sources and shading


- A light source is a power source that emits light (instead of reflecting it)
- Laser – a single ray
- Point source – like a light bulb
- Line source – fluorescent light bulb
- Area source

-CSE 252A

Nearby point source model


$$L = \rho_d(x) \left(\frac{\hat{N}(x) \cdot S(x)}{r(x)^2} \right)$$

- \hat{N} is the surface normal
- ρ_d is diffuse (Lambertian) albedo
- S is source vector - a vector from x to the source, whose length is the intensity term



-CSE 252A

Nearby Line Source



Intensity due to line source varies with inverse distance, if the source is long enough

-CSE 252A

Distant Point Source Model

- Assume that all points in the model are close to each other with respect to the distance to the source. Then the source vector doesn't vary much, and the distance doesn't vary much either, and we can roll the constants together to get:

$$L = \rho_d(x) N(x) \cdot S(x)$$

- N is the surface normal
- ρ_d is diffuse (Lambertian) albedo
- S is source vector - a vector in source direction, whose length is the intensity term



-CSE 252A


Lighting at infinity

- Direction is a three vector s , with $|s| = I$.
- Described as function on a sphere: radiance as a function of direction $r(s)$
- Single point source is a delta function at some direction
- Multiple point sources: sum of delta functions



-CSE 252A

Diffuse lighting at infinity: Spherical Harmonics



-CSE 252A

