

Chapter 5 - Integral Dependence and Valuations

Flandre

20 Jun 2026

Exercise 5.1

Use *Theorem 5.10*, trivial.

Exercise 5.2

Use *Theorem 5.10*, trivial.

Exercise 5.3

Because

$$\operatorname{Im}(f \otimes 1) = (\operatorname{Im} f) \otimes_A C$$

Therefore $B' \otimes_A C$ is integral over $\operatorname{Im} f \otimes_A C$, as B' is integral over $\operatorname{Im} f$.

Remark 5.1

Indeed, *Exercise 5.3* includes *Proposition 5.6(ii)* as special case, as $S^{-1}B \cong S^{-1}A \otimes_A B$.

Exercise 5.4

Following from the hinted counterexample, the answer is no.

Exercise 5.5

(i) Since $x^{-1} \in B$, $\exists a_{n-1}, \dots, a_0$, s.t.

$$x^{-n} + a_{n-1}x^{-n+1} + \dots + a_1x^{-1} + a_0 = 0.$$

Hence

$$x^{-1} = -a_{n-1} - a_{n-2}x - \dots - a_1^{n-2} - a_0x^{n-1} \in A$$

(ii) Use *Corollary 5.8*, trivial.

Exercise 5.6

Indeed. Denote the integral ring homomorphism

$$f_k : A \longrightarrow B_k$$

and we have

$$\prod_{k=1}^n f_k : A^n \longrightarrow \prod_{k=1}^n B_k$$

as an integral ring homomorphism.

(Elimination of terms to make the elements integral is easy, as we can nest those polynomials by terms.)

Exercise 5.7