

Computeranimation

Lesson 1 – Transformations

Motivation

Topics

- **Rigid Transformation**
- Animation
- Collision
- Dynamic
- Mass-Spring Simulation
- Rigging and Skeletal Animation
- Motion Capturing using RGB-D Sensor

Rigid Bodies

Math Primer

Positions

Spaces

Matrices

Transformations




Application

Rigid Bodies

Definition:


- Bodies that do not change by any influences
- In physics they exist only theoretically, deformation is neglected

→ Each pair of points has always constant distance




Rigid Bodies

- Rigid bodies usually are modelled as a set of discrete points




Rigid Bodies

- Rigid bodies usually are modelled as a set of discrete points
- The points have constant relative positions



Rigid Bodies

- Rigid bodies usually are modelled as a set of discrete points
- The points have constant relative positions
- The single points' masses sum up to the rigid body's total mass



rigid body with total mass M_{total}

point mass with single mass m_i

$$M_{total} = \sum_{i=0}^{N-1} m_i$$

Rigid Bodies

Math Primer

Positions

Spaces

Matrices

Transformations


Application

Math Primer

Position

- A **unique** location within a **geometrical space**
- This space is defined by a **coordinate system**
- A Coordinate System consists of an **origin** ($\vec{0}$) and ...
- ... spanning vectors (e.g. *unit vectors* in *Cartesian Systems*)

- (a) rectilinear systems
- (b) rectilinear & orthogonal systems
- (c) curvilinear orthogonal grid
- (d) curvilinear grid




P Position within the coordinate system

Math Primer


Position

- Points in a space usually are defined via coordinates $\in \mathbb{R}^2$ or $\in \mathbb{R}^3$


\mathbb{R}^2



\mathbb{R}^3 – right handed



\mathbb{R}^3 – left handed



Rigid Bodies

Math Primer

Positions

Spaces

Matrices


Transformations

Application

Math Primer

Spaces

- Within a single Coordinate System a point is *uniquely* defined by its **coordinates**





• $P = \begin{pmatrix} x_P \\ y_P \end{pmatrix}$

Math Primer

Spaces


- Within a single Coordinate System a point is uniquely defined by its **coordinates**
- Things become complicated when more point of views are needed
- The position of P **depends** on the point of view




Math Primer

Spaces

- Positions often need to be expressed in different spaces



"The Astronaut wants to know the position of the landing place in world's local space"



Rigid Bodies

Math Primer

Positions

Spaces

Matrices

Transformations

Application

Math Primer

Matrices

- A Table of Elements (a_{ij}) arranged in a rectangular structure of M rows and N columns

$$\begin{bmatrix} a_{11} & \cdots & a_{1N} \\ \vdots & \ddots & \vdots \\ a_{M1} & \cdots & a_{MN} \end{bmatrix}$$

Math Primer

Special Matrices

Symmetric Matrices

$$A = A^T = \begin{bmatrix} 1 & 10 & 30 \\ 10 & 2 & 64 \\ 30 & 64 & 3 \end{bmatrix}$$

Quadratic Matrices

$$M = N \rightarrow \begin{bmatrix} a_{11} & \cdots & a_{1M} \\ \vdots & \ddots & \vdots \\ a_{M1} & \cdots & a_{MM} \end{bmatrix}$$

Orthogonal Matrices

$$|\det (A)| = 1$$

Math Primer


Orthogonal Matrices

- Of special interest for computer graphics (spaces, rotations, ...)
- Most important property: $A^{-1} = A^T \rightarrow A^T A = I$

Rotation Matrix

- A simple 2D matrix that rotates a point about the angle α

$$R_\alpha = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$




Math Primer

Matrix-Vector-Multiplication

- Represents a transformation to a vector.
- Example (rotation about 45°):

$$R_{45^\circ} = \begin{bmatrix} \cos 45^\circ & -\sin 45^\circ \\ \sin 45^\circ & \cos 45^\circ \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \rightarrow \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \cdot 1 - \frac{1}{\sqrt{2}} \cdot 0 \\ \frac{1}{\sqrt{2}} \cdot 1 + \frac{1}{\sqrt{2}} \cdot 0 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$



Rigid Bodies

Math Primer

Positions

Spaces

Matrices

Transformations

Application

Transformations

Spaces

- Expressing a single Point (defined in one Coordinate System) in another system is called **Transformation**

Transformations

Affine Transformations

Linear Transformations

Rigid Transformations

Transformations

Affine Transformation

$$f: X \rightarrow Y \quad x \rightarrow A \cdot x + b$$

A: linear Transformation
b: vector in Y

preserve...

- ... points
- ... preserve lines
- ... preserve planes
- ... parallel lines remain parallel

not necessarily preserve...

- ... angles
- ... ratios of distances between points on a straight line

Transformations

Affine Transformation


Example:

$$\begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 \cdot 0 + 1 \cdot 1 - 1 \\ 2 \cdot 0 + 1 \cdot 1 - 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} -1 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 \cdot 0 + 1 \cdot 0 - 1 \\ 2 \cdot 0 + 1 \cdot 0 - 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$


$$\begin{bmatrix} 2 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 0 \end{bmatrix} + \begin{bmatrix} -1 \\ -1 \end{bmatrix} = \begin{bmatrix} 0 \cdot 2 + 1 \cdot 0 - 1 \\ 2 \cdot 2 + 1 \cdot 0 - 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 3 \end{bmatrix}$$




Transformations

Affine Transformation


Classes of Affine Transformations




translate




rotate about origin



scale about origin



shear




reflection


Transformations

Rigid Transformations


A sub-category of affine transformations




translate




rotate about origin



scale about origin



shear



reflection

preserves the distance between every pair of points

Transformations

Rigid Transformation Pipeline

- Since rigid transformations (amongst others) are linear, they can be put together

only rotation:
$$\begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \mathbf{R}_{-45^\circ} \cdot (\mathbf{R}_{+45^\circ} \cdot \begin{bmatrix} x \\ y \end{bmatrix}) \rightarrow \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \cdot \left(\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} \right)$$

only translation:
$$\begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \left(\begin{bmatrix} x \\ y \end{bmatrix} + \vec{t}_0 \right) + \vec{t}_1 \rightarrow \begin{bmatrix} x \\ y \end{bmatrix} + (\vec{t}_0 + \vec{t}_1)$$

both:
$$\begin{bmatrix} x \\ y \end{bmatrix} \rightarrow \mathbf{R}_1 \cdot \left(\mathbf{R}_0 \cdot \begin{bmatrix} x \\ y \end{bmatrix} + \vec{t}_0 \right) + \vec{t}_1$$

Rigid Bodies

Math Primer

Positions

Spaces

Matrices

Transformations

Application