Exact Recovery of a Linear Differential Operator from a Single Solution

Identification, Factor Decomposition, the Interaction Term, and a Nonlinear Extension

October 2026

Abstract

Given one exact solution u of an unknown differential equation in one space and one time variable, we ask what u "remembers" about the equation. We answer in four steps. (i) *Identification*: for constant coefficients, u determines its governing operator P if and only if no partial quotient $(P/P_i)(\partial)u$ is an exponential-polynomial; no finite amount of data can certify this, and a black-box test is impossible, yet it becomes decidable with a modest extra hypothesis. (ii) *Decomposition*: u splits into one component per irreducible *factor* of the symbol, with an exponential-polynomial ambiguity of known finite dimension; the split is generally *not* possible per additive *term* of the operator. (iii) *Interaction*: for evolution equations $u_t = \sum_n A_n u$ with commuting terms the part of the solution due to the interplay of the terms has the exact form $\prod_n(\mathrm{Id} + \Delta_n) - \mathrm{Id} - \sum_n \Delta_n$ applied to the initial datum, it costs $O(N)$ per value but $\Theta(2^N)$ as a full table, and it is not a continuous function of the data in Sobolev norms. (iv) *Nonlinear* equations (Burgers, KdV–Burgers, Fisher–KPP) admit analogous statements, with sharp asymptotics across a shock.

We present the mathematics, the resulting algorithm component by component, a worked example, and the precise limits of the method. Every result carries a verification status.

Contents

1 Introduction **3**

1.1 The question . . . 3
1.2 How to read the status labels . . . 3
1.3 Overview of the results . . . 3
1.4 Notation at a glance . . . 4
1.5 Roadmap . . . 4

2 Mathematical preliminaries **5**

2.1 Derivatives as polynomials . . . 5
2.2 Annihilators, ideals, and what a solution "knows" . . . 6
2.3 Exponential-polynomials . . . 8

3 One variable: the ODE prototype **8**

3.1 Identification by the Wronskian . . . 9

3.2 The companion matrix and spectral form . . . 9
3.3 Tensor construction versus module construction . . . 10

4 Identification **10**
4.1 The main criterion . . . 11
4.2 The spectral form . . . 12
4.3 Limits of what can be certified . . . 13

5 Decomposition into components **14**
5.1 Why per-term splitting fails . . . 14
5.2 Why factors do work . . . 15
5.3 An explicit projector for evolution factors . . . 17
5.4 Where decomposition fails . . . 17
5.5 Regularity: when are the components in H^m? . . . 17

6 Terms inside a factor and the interaction term **19**
6.1 Definition via the flows of the single terms . . . 20
6.2 Exact formula in the commuting case . . . 20
6.3 Cost: $O(N)$ versus 2^N . . . 22
6.4 Why E_{int} cannot be removed or regularized stably . . . 23
6.5 Time-dependent perturbation theory . . . 23
6.6 Non-commuting terms . . . 24

7 Nonlinear equations **24**
7.1 Interaction term for flows of single terms . . . 25
7.2 Before, at, and after the shock . . . 25
7.3 Identification relative to a dictionary . . . 27
7.4 Linear versus nonlinear: the same idea, two structures . . . 27

8 The algorithm **28**
8.1 Inputs and data models . . . 28
8.2 The pipeline . . . 29
8.3 The steps in detail . . . 29
8.4 Worked example . . . 31
8.5 Complexity summary . . . 32

9 Limits, open problems, and status **33**

A Glossary of tools **34**

B Primer for readers new to the concepts **34**

C Scope: what this paper contains and what it does not **36**

D Notes for the author **37**

1 Introduction

In this section	
Starting from	one function $u(t,x)$ known exactly on a region $\Omega \subset \mathbb{R}^2$ (no noise), which solves $P(\partial_t, \partial_x)u = 0$ for a constant-coefficient operator P that is *not* given.
Why this section	to state precisely what we want from u (three questions) and to give the reader a map of the answers before the details.
Role in the paper	frames Q1–Q3, which are answered in Sections 4, 5 and 6; Sections 7–9 extend and assemble them.

1.1 The question

Let $u(t,x)$ be a function on a region $\Omega \subset \mathbb{R}^2$, known without noise, and suppose it solves a linear equation $P(\partial_t, \partial_x)\, u = 0$. We do not assume P is known. Three questions arise naturally.

- **Q1. Identification.** Can the operator be reconstructed from u, and when is the answer unique?
- **Q2. Decomposition.** If P is built from pieces, can u be split into the parts that each piece generates?
- **Q3. Interaction.** If the pieces are the *terms* of the operator (diffusion, advection, ...), what part of u is due to *no single term* but to their interplay, and can it be computed exactly and cheaply?

The setting is deliberately analytic: no statistical model, no learned weights, no noise. The aim is an *exact* procedure, and a precise statement of what is impossible.

1.2 How to read the status labels

Every statement carries one of the labels [proved] (complete proof, independently reviewed), [proved mod. X] (complete except for a cited published theorem X whose source was checked), [conditional] (depends on an unverified lemma), [open], or [false] (a counterexample exists). The reviewers were independent AI agents, not human referees; fifteen core algebraic theorems were additionally machine-checked in Lean 4 (Section 9). Grey tags such as (ledger C3-T1) refer to entries in the project's claim ledger.

1.3 Overview of the results

Figure 1 summarizes the logical chain, read left to right: each box is a task, and each arrow is a theorem that allows the next task to be carried out. Table 1 lists the main statements with their status; the rightmost column uses the labels just explained, and every row points to the theorem where it is proved.




Figure 1: Logical structure of the method. Each arrow is a theorem with explicit hypotheses, so a step can only be trusted under the hypotheses of the theorem preceding it. The three linear stages (blue) are followed by the nonlinear extension (red); the algorithm that chains them is in Section 8.

Topic	Result	Status
Identification	$\mathrm{Ann}(u) = (P) \iff (P/P_i)(\partial)u \notin \mathrm{EP}$ for all irreducible $P_i \mid P$ (Thm. 4.1)	[proved]
Spectral form	$\mathrm{Ann}(u) = (P) \iff \sigma_p(u) = \emptyset$ and $G_u \sim P$ (Prop. 4.3)	[proved]
Data limits	no finite certificate; black box impossible; decidable with a bound or a holonomic model (Sec. 4.3)	[proved] / [proved mod. (AC)]
Decomposition	$\ker P = \sum_j \ker Q_j$ on simply connected Ω, ambiguity $\delta(P)$ (Thm. 5.2)	[proved]
Sobolev regularity	components in H^m for all $u \iff$ sum of kernels closed ($\gamma_N > 0$) and dense (Thm. 5.5)	[proved]
Terms vs. factors	splitting per additive term is false (Prop. 5.1)	[proved][E9]
Interaction term	$E_{\mathrm{int}} = \prod(\mathrm{Id} + \Delta_n) - \mathrm{Id} - \sum \Delta_n$; $O(N)$ value, $\Theta(2^N)$ table (Thm. 6.1)	[proved]
Stability	$u_0 \mapsto E_{\mathrm{int}}$ is not continuous in H^m (Prop. 6.3)	[proved]
Nonlinear	Burgers: closed form before the shock, $\Theta(\nu)$ in L^1 after (Sec. 7)	[proved] / [proved mod. Kružkov]
Dictionary identification	exact only modulo differential consequences of the equation (Thm. 7.2)	[proved]

Table 1: Main results, in the order in which they are proved. Columns: topic, statement, verification status. "Proved" means a complete proof that survived an independent review (see Section 9 for what this means).

1.4 Notation at a glance

Table 2 collects the symbols used throughout, with the place where each is defined; the appendix glossary (Appendix A) covers the named theorems.

1.5 Roadmap

Section 2 fixes the algebraic language (symbols, annihilators, exponential-polynomials). Section 3 introduces, in one variable, the objects that reappear later (minimal annihilator, companion matrix, spectral projectors). Section 4 answers Q1: what u determines. Section 5 answers Q2: how u splits once the operator is known. Section 6 answers Q3: what is left over when

Symbol	Meaning	Defined in
(t, x), Ω	time and space variables; open region of $\mathbb{R}^2$ on which u is known	Sec. 1
$\partial = (\partial_t, \partial_x)$	partial derivatives	Sec. 2
$\mathcal{R} = \mathbb{C}[\partial_t, \partial_x]$	ring of constant-coefficient operators ($\cong \mathbb{C}[\tau, \xi]$)	Def. 2.1
P, P_i, Q_j	the operator (its symbol), its irreducible factors, the coprime factors $P_j^{m_j}$	Sec. 2, 5
$V(P)$	characteristic variety, zeros of the symbol in $\mathbb{C}^2$	Def. 2.1
$\mathrm{Ann}(u)$	ideal of all operators annihilating u	Def. 2.2
G_u, J_u	$G_u = \gcd \mathrm{Ann}(u)$ and the cofactor, $\mathrm{Ann}(u) = G_u J_u$	Lemma 2.3
EP	exponential-polynomials	Def. 2.4
$\sigma_p(u)$	point spectrum $= V(J_u)$	Prop. 4.3
$\delta(P)$	dimension of the ambiguity of the factor decomposition	Thm. 5.2
c_F, γ_N	Friedrichs cosine; smallest nonzero eigenvalue of S^*S	Thm. 5.5
A_n, $S_n(t)$, Δ_n	terms of the evolution, their flows e^{tA_n}, increments $S_n - \mathrm{Id}$	Sec. 6
E_{int}	interaction term	Eq. (3)
T_n, $\mathcal{D}$	nonlinear terms; dictionary of jet monomials	Sec. 7

Table 2: Notation guide. A few letters are used with more than one meaning in different sections (for instance S for the sum map of Theorem 5.5 and for the semigroup $S(t)$; D for d/dx in Section 3 and for a diffusion constant elsewhere; J for ideals); the context always fixes the meaning.

one splits by terms, how much it costs to compute it, and why it cannot be stabilized. Section 7 extends the questions to nonlinear equations. Section 8 assembles everything into an algorithm with a worked example, and Section 9 states the limits.

2 Mathematical preliminaries

In this section	
Starting from	constant coefficients; functions or distributions on $\Omega \subset \mathbb{R}^2$ with complex values. No algebra beyond polynomials is assumed; Appendix B explains ideals, annihilators, exponential-polynomials and the Chinese remainder theorem with concrete examples.
Why this section	three notions carry the whole theory: the *symbol* of an operator (derivatives become polynomials), the *annihilator* of a function (all the equations it satisfies), and *exponential-polynomials* (the "trivial" solutions).
Role in the paper	foundation for Sections 4 and 5. Needs Fourier/Laplace transforms, polynomial factorization and the linear algebra of commuting matrices.

2.1 Derivatives as polynomials

Write ∂_t, ∂_x for the partial derivatives. A constant-coefficient operator is a polynomial in them,

$$P(\partial_t, \partial_x) = \sum_{j,k} c_{jk}\, \partial_t^j \partial_x^k, \qquad c_{jk} \in \mathbb{C}.$$

The exponentials $e_{\tau,\xi}(t,x) = e^{\tau t+\xi x}$ are eigenfunctions of every derivative,

$$\partial_t e_{\tau,\xi} = \tau\, e_{\tau,\xi}, \qquad \partial_x e_{\tau,\xi} = \xi\, e_{\tau,\xi}, \qquad \text{hence} \qquad P(\partial)\, e_{\tau,\xi} = P(\tau,\xi)\, e_{\tau,\xi}.$$

Definition 2.1 (Operator ring, symbol, characteristic variety)

$\mathcal{R} = \mathbb{C}[\partial_t, \partial_x] \cong \mathbb{C}[\tau,\xi]$ is the commutative ring of constant-coefficient operators; the polynomial $P(\tau,\xi)$ is the *symbol* of $P(\partial)$. The *characteristic variety* is $V(P) = \{(\tau,\xi) \in \mathbb{C}^2 : P(\tau,\xi) = 0\}$.

So $e_{\tau,\xi}$ solves $P(\partial)u = 0$ exactly when $(\tau,\xi) \in V(P)$ (Fourier case: $\xi = i\kappa$). Figure 2 shows two examples.



Figure 2: Real slices of characteristic varieties. A *single exponential* corresponds to a single point of the variety; a rich, non-exponential solution "fills" the whole curve.

2.2 Annihilators, ideals, and what a solution "knows"

A single u is usually annihilated by many operators (the function e^x is killed by $\partial_x - 1$ and by $\partial_x^2 - 1$ and ...).

Definition 2.2 (Annihilator)

$\mathrm{Ann}(u) = \{Q \in \mathcal{R} :\ Q(\partial)u = 0\}$.

(Readers new to ideals: see Appendix B, §B.1.) For instance $\mathrm{Ann}(e^x)$ contains $\partial_x - 1$ and its multiples such as $\partial_x^2 - 1 = (\partial_x - 1)(\partial_x + 1)$. It is an *ideal* of $\mathcal{R}$: if $Qu = 0$ and $S \in \mathcal{R}$, then $(SQ)u = S(Qu) = 0$; and sums of annihilators annihilate. Ideals are the polynomial analogue of the multiples of an integer: in one variable they are generated by a single polynomial, in two variables they may need several, and this is the source of all the subtleties below. We use three facts about $\mathbb{C}[\tau,\xi]$.

(F1) **Unique factorization** (the polynomial analogue of prime factorization): $P = \prod_i P_i^{e_i}$ with P_i irreducible (the "primes") and e_i their multiplicities, unique up to constants; gcd is defined as for integers.
Used for: extracting the common part of an ideal (Lemma 2.3) and splitting an operator into factors (Section 5).

(F2) **Bézout:** two polynomials in τ,ξ without common factor have only finitely many common zeros.
Geometrically: two curves in $\mathbb{C}^2$ with no common component meet in isolated points.
Used for: showing that what is left of an ideal after removing its gcd is "small" (Lemma 2.3).

(F3) **Nullstellensatz** (Hilbert): if q vanishes on $V(I)$, the set of common zeros of all elements of the ideal I, then $q^k \in I$ for some k (a power is needed because zeros do not see multiplicities). In particular an ideal without zeros is the whole ring.
Used for: translating statements about zero sets into statements about ideals (proof of Theorem 4.1, Proposition 4.3).

All three are classical [1, 6]; they hold over $\mathbb{C}$, which is algebraically closed (the Nullstellensatz fails over $\mathbb{R}$).

In the next lemma

- $\mathcal{R} = \mathbb{C}[\tau, \xi]$ is the ring of Definition 2.1;
- $I \subset \mathcal{R}$ is any ideal (in our use, $I = \mathrm{Ann}(u)$);
- $g = \gcd(I)$ is the polynomial of largest degree that divides every element of I, defined up to a nonzero constant;
- $J = \{f/g : f \in I\}$ is the ideal of the quotients;
- $V(J) \subset \mathbb{C}^2$ is the set of common zeros of all elements of J.

The lemma says that an ideal always splits into a "curve part" g and a residue J that vanishes only at finitely many points.

Lemma 2.3 (Splitting off the gcd) (ledger C3-L2) [proved]

Every nonzero ideal $I \subset \mathcal{R}$ factors as $I = g \cdot J$ with $g = \gcd(I)$ and $V(J)$ finite.

Proof sketch. Every $f \in I$ is divisible by g, so $J = \{f/g\}$ is well defined; it is again an ideal and $I = gJ$. The elements of J have no common factor (a common factor h would make gh a common divisor of I).

Pick $0 \neq h_1 \in J$; if h_1 is constant, $J = \mathcal{R}$ and $V(J) = \emptyset$. Otherwise let $p_1, \dots, p_r$ be its pairwise non-associate irreducible factors. For each i there is $h_i' \in J$ with $p_i \nmid h_i'$ (otherwise p_i would be a common factor). Put

$$h_2 = \sum_i h_i' \prod_{k \neq i} p_k \ \in J.$$

Modulo p_i only the i-th summand survives and it is not divisible by p_i, so no p_i divides h_2: the two elements $h_1, h_2 \in J$ have no common factor. By (F2) they have finitely many common zeros, and $V(J) \subseteq V(h_1, h_2)$ is finite. □

For $I = \mathrm{Ann}(u)$ we write $G_u = \gcd \mathrm{Ann}(u)$ (defined up to a constant) and J_u for the cofactor ideal: $\mathrm{Ann}(u) = G_u J_u$. Intuitively G_u is the "genuinely infinite-dimensional" part of the equation of u, and J_u the finitely many exponentials that happen to be present. These two objects drive Section 4.

Two extreme examples

- For $u = e^x$, seen as a function of (t, x), $\mathrm{Ann}(u) = (\tau,\ \xi - 1)$ is the ideal of the single point $(0, 1)$: $G_u = 1$ and $J_u = \mathrm{Ann}(u)$, so u has no infinite part.
- For $u = f(x + t)$ with f generic (not an exponential-polynomial), $\mathrm{Ann}(u) = (\tau - \xi)$: $G_u = \tau - \xi$ and $J_u = \mathcal{R}$ (Section 4).

2.3 Exponential-polynomials

Notation used in this subsection

- Recall that $\mathcal{R} = \mathbb{C}[\partial_t, \partial_x] \cong \mathbb{C}[\tau, \xi]$ (Definition 2.1) acts on functions by differentiation: $Q \in \mathcal{R}$ sends w to $Q(\partial)w$.
- For a function w,

$$\mathcal{R}w = \operatorname{span}\{\partial_t^j \partial_x^k w : \ j,k \geq 0\} = \{Q(\partial)w : \ Q \in \mathcal{R}\}$$

  is the set of all derivatives of w and their linear combinations; it is a vector space over $\mathbb{C}$ and $\dim_{\mathbb{C}}$ denotes its dimension as such.
- The map $Q \mapsto Q(\partial)w$ from $\mathcal{R}$ onto $\mathcal{R}w$ has kernel $\operatorname{Ann}(w)$, which is why $\mathcal{R}w \cong \mathcal{R}/\operatorname{Ann}(w)$: the quotient identifies two operators that act in the same way on w.
- In a sum $\sum_j p_j(t,x)e^{\tau_j t+\xi_j x}$, each p_j is a polynomial in (t,x) and each pair $(\tau_j, \xi_j) \in \mathbb{C}^2$ is an *exponent*, i.e. a point of $\mathbb{C}^2$ as in Section 2.

Definition 2.4 (EP) (see Appendix B, §B.2)

EP is the set of finite sums $\sum_j p_j(t,x)\, e^{\tau_j t+\xi_j x}$ with polynomials p_j.

Lemma 2.5 (Finite dimension criterion) (ledger C3-L3) [proved]

$w \in \text{EP} \iff \dim_{\mathbb{C}}(\mathcal{R}w) < \infty$, where $\mathcal{R}w = \operatorname{span}\{\partial_t^j \partial_x^k w\}$ is the span of all derivatives of w. In that case $\mathcal{R}w \cong \mathcal{R}/\operatorname{Ann}(w)$ and $V(\operatorname{Ann} w)$ is the finite set of exponents $\{(\tau_j, \xi_j)\}$.

Idea of proof. ($\Rightarrow$) Derivatives of $p\, e^{\tau t+\xi x}$ are (polynomial of degree $\leq \deg p) \cdot e^{\tau t+\xi x}$. ($\Leftarrow$) ∂_t and ∂_x act as commuting matrices on the finite-dimensional space $\mathcal{R}w$. Commuting matrices are simultaneously triangularizable (the multivariable Jordan form); their joint eigenvalues are the exponents (τ_j, ξ_j), and the nilpotent parts produce the polynomial prefactors. □

> **Why it matters.** Lemma 2.5 is the computational heart of the method: "is w an exponential-polynomial?" is a *finite-dimensional linear algebra question*. It is the test used in Theorem 4.1 and in Step 3 of the algorithm.

3 One variable: the ODE prototype

In this section

Starting from	one variable x, an ODE $p(D)u = 0$ with constant coefficients, $D = d/dx$, p known.
Why this section	in one variable every object of the later sections (minimal annihilator, spectral projectors, companion matrix) appears with explicit formulas and no geometry of curves.
Role in the paper	prototype for Section 4 (identification) and Section 5 (the Lagrange projectors reappear as Eq. (2)).

This section introduces, in the simplest setting, objects that reappear in Sections 4 and 5: the minimal annihilator, the companion matrix and the spectral projectors.

Setting and symbols

- We consider an ODE $p(D)u = 0$ in one variable x, with $D = d/dx$ (in this section only; elsewhere D is a diffusion constant).
- The polynomial $p(\lambda) = \prod_j (\lambda - \lambda_j)^{m_j}$ has distinct roots $\lambda_j \in \mathbb{C}$ with multiplicities $m_j \geq 1$, so that $m = \deg p = \sum_j m_j$ is the *order* of the equation (not to be confused with the Sobolev index m of Section 5).
- The set of solutions is the *kernel* $\ker p(D)$, and

$$\ker p(D) = \Big\{ \sum_j q_j(x) e^{\lambda_j x} \,:\, \deg q_j < m_j \Big\},$$

where each q_j is a polynomial of degree $< m_j$.

Why this form

For one root, $(D - \lambda)(q\, e^{\lambda x}) = q' e^{\lambda x}$: applying $D - \lambda$ lowers the degree of q by one. Hence $(D - \lambda_j)^{m_j}$ kills $q\, e^{\lambda_j x}$ exactly when $q^{(m_j)} = 0$, i.e. $\deg q < m_j$.

Each root therefore contributes the m_j functions $x^k e^{\lambda_j x}$ $(0 \leq k < m_j)$: these are the natural basis functions of the equation, finitely many, and together they form a basis of $\ker p(D)$, a space of dimension m.

Example

$p(\lambda) = (\lambda - 1)^2(\lambda + 2)^3$ has $\lambda_1 = 1$, $m_1 = 2$, $\lambda_2 = -2$, $m_2 = 3$, order $m = 5$. A basis of the solutions is

$$e^x, \quad xe^x, \quad e^{-2x}, \quad xe^{-2x}, \quad x^2 e^{-2x},$$

so every solution is

$$c_1 e^x + c_2 x e^x + c_3 e^{-2x} + c_4 x e^{-2x} + c_5 x^2 e^{-2x}.$$

3.1 Identification by the Wronskian

Proposition 3.1 (ODE identification)

Let $u \in \ker p(D)$. The minimal annihilator of u has degree equal to the number of independent basis functions actually present in u; it equals p if and only if $u, u', \ldots, u^{(m-1)}$ are linearly independent (nonvanishing Wronskian), i.e. iff every root of p appears in u with its full multiplicity.

Reason. $\dim \operatorname{span}\{u, u', u'', \ldots\}$ is the degree of the minimal annihilator (Lemma 2.5 in one variable); a missing root lowers it.

3.2 The companion matrix and spectral form

Write $p(\lambda) = \lambda^m + a_{m-1}\lambda^{m-1} + \cdots + a_0$ (monic) and put $X = (u, u', \ldots, u^{(m-1)})^\top$. Then $X' = CX$ with

$$C = \begin{pmatrix} 0 & 1 & & \\ & 0 & \ddots & \\ & & \ddots & 1 \\ -a_0 & -a_1 & \cdots & -a_{m-1} \end{pmatrix}, \qquad X(x) = e^{xC} X(0).$$

The eigenvalues of C are the roots of p. For distinct roots the matrix exponential has the spectral form

$$e^{xC} = \sum_{j=1}^{m} e^{\lambda_j x}\, \Pi_j, \qquad \Pi_j = \prod_{k \neq j} \frac{C - \lambda_k \mathrm{Id}}{\lambda_j - \lambda_k}, \tag{1}$$

where the Π_j are the Lagrange projectors onto the eigenspaces. The last row of C carries the whole equation; C is a *matrix over the base field* for an ODE, but, as soon as the coefficients depend on x, or there is a second variable, its last row has entries that are operators and the object becomes a matrix over a noncommutative or polynomial ring. The invariant statement is that the whole equation corresponds to the cyclic module $\mathcal{R}/\operatorname{Ann}(u)$ (Section 2); the companion matrix is a way to compute with it relative to one chosen direction.

Proposition 3.2 (Projectors of the companion matrix) (ledger II-T2) [proved]

The projectors Π_j are explicit rational functions of the coefficients, with poles exactly where roots collide; they are polynomial in the coefficients if and only if the corresponding varieties are disjoint. They are *not* orthogonal in general: the companion matrix is not normal for $m = 2$ [false] (the contrary was an early conjecture).

Hence Hilbert-space orthogonality arguments need an inner product adapted to the problem, not the standard one on the coefficient vector.

3.3 Tensor construction versus module construction

One may encode a PDE in a higher-order tensor of derivatives, $J = \partial_{x_{i_1}} \otimes \cdots \otimes \partial_{x_{i_n}}$, or by the companion form. The comparison, (ledger Gate 3): the *principal symbol tensor* $\sigma_m(P) \in \operatorname{Sym}^m(\mathbb{C}^2)$ only sees the highest-order part of P (for the heat operator it sees ξ^2 alone), so it is an invariant "at infinity"; the *cyclic module* $\mathcal{R}/\operatorname{Ann}(u)$ is the complete invariant. All subsequent results use the module.

Takeaway. This section shows how an ODE's polynomial encodes its solution space. Its roots determine the exponential rates, their multiplicities determine the polynomial prefactors, the companion matrix represents differentiation, and spectral projectors isolate the different modes. The later sections generalize these ideas to multiple commuting derivatives, where the same structure survives but requires more algebraic machinery: the minimal annihilator becomes the ideal $\operatorname{Ann}(u)$ (Section 4) and the Lagrange projectors reappear as Eq. (2) (Section 5).

4 Identification

In this section	
Starting from	$\Omega \subset \mathbb{R}^2$ open and connected, $u \in \mathcal{D}'(\Omega)$, constant coefficients, and an operator P with $P(\partial)u = 0$ is available (any annihilator).
Why this section	to decide whether u pins down the equation it solves, i.e. whether P can be recovered from u alone.
Role in the paper	first link of the chain in Figure 1: it defines G_u and the test "$\in$ EP" that Sections 5 and 8 reuse. *Result:* a necessary and sufficient criterion (Theorem 4.1) and its limits.

4.1 The main criterion

Let $\Omega \subset \mathbb{R}^2$ be open and *connected* (connectedness is necessary: on disconnected Ω different components can satisfy different equations), $u \in \mathcal{D}'(\Omega)$ with $P(\partial)u = 0$ and $P = \prod_i P_i^{e_i}$ the irreducible factorization.

Terminology

- $\mathrm{EP}(\Omega)$ denotes the restrictions to Ω of exponential-polynomials (Definition 2.4).
- P is *reduced* if all $e_i = 1$ (no repeated factor).
- P is *monic in* τ if, written as a polynomial in τ with coefficients in $\mathbb{C}[\xi]$, its leading coefficient is a nonzero constant; then $k = \deg_\tau P$ is its order in time and the *t-jet* of u is the vector $(u, \partial_t u, \dots, \partial_t^{k-1} u)$.
- In (a) we remove one copy of P_i from P; in (b) we remove the whole power $P_i^{e_i}$, which makes (b) sensitive to multiplicities.

Theorem 4.1 (Identification criterion) (ledger C3-T1) [proved]

(a) $\mathrm{Ann}(u) = (P) \iff \left(\frac{P}{P_i}\right)(\partial)\, u \notin \mathrm{EP}(\Omega)$ for every irreducible factor P_i of P.

(b) $V(\mathrm{Ann}\, u) = V(P)$ iff $(P/P_i^{e_i})(\partial)u \notin \mathrm{EP}$ for all i.

(c) For P reduced the statements (a) and (b) coincide; for P monic in τ the test can be carried out on the t-jet $u, \partial_t u, \dots, \partial_t^{k-1} u$ (ledger II-C1).

Intuition

Strip one irreducible factor off P and look at what is left: $w_i = (P/P_i)(\partial)u$ lies in $\ker P_i$. If w_i is a "generic" member of that kernel (not a finite sum of exponentials), then no smaller operator can kill u. If w_i is an exponential-polynomial, u contains a finite-dimensional piece that is also killed by an operator of lower order, and u cannot tell the large equation from the small one.

Proposition 4.2 (Uniqueness of the identified operator)

If $\mathrm{Ann}(u) = (P)$, then P is determined by u up to a nonzero constant factor: every constant-coefficient operator Q with $Q(\partial)u = 0$ is a multiple $Q = SP$ with $S \in \mathcal{R}$, and P is, up to a constant, the *only* annihilator of minimal degree. If instead $\mathrm{Ann}(u) \supsetneq (P)$, the equation is *not* identified: P cannot be recovered from u, and the only object intrinsic to u is the generator G_u of the infinite part of $\mathrm{Ann}(u)$ (Section 4.2).

Proof. $Q \in (P)$ means $Q = SP$ for some $S \in \mathcal{R}$, which is the first claim. Since degrees add, $\deg Q = \deg S + \deg P \geq \deg P$ for $Q \neq 0$, with equality iff S is constant. Two generators of the same principal ideal differ by a unit of $\mathcal{R}$, i.e. by a nonzero constant. □

Example 4.1 (Heat equation). Take $P = \tau - D\xi^2$ with $D > 0$, i.e. $u_t = D u_{xx}$. P is irreducible, so $P/P_i = 1$ and the criterion reads: $\mathrm{Ann}(u) = (P)$ iff $u \notin \mathrm{EP}$. Three cases show what this means.

(i) *One Fourier mode.* Let $u = e^{-D\kappa^2 t + i\kappa x}$ with $\kappa \in \mathbb{R}$ fixed. Then $u_t = -D\kappa^2 u$ and $u_{xx} = -\kappa^2 u$, so $P(\partial)u = 0$. But also $(\partial_x - i\kappa)u = 0$ and $(\partial_t + D\kappa^2)u = 0$, so

$$\mathrm{Ann}(u) = (\tau + D\kappa^2,\ \xi - i\kappa),$$

the maximal ideal of the point $(-D\kappa^2, i\kappa)$ of $V(P)$. It is strictly larger than (P); indeed

$$P = (\tau + D\kappa^2) - D\,(\xi - i\kappa)(\xi + i\kappa) \in \mathrm{Ann}(u).$$

(ii) *A finite superposition of modes.* Let $u = \sum_{j=1}^{r} c_j e^{-D\kappa_j^2 t + i\kappa_j x}$ with distinct κ_j and $c_j \neq 0$. Then $u \in \mathrm{EP}$, so $\mathrm{Ann}(u) \neq (P)$: it is the ideal of the r points $(-D\kappa_j^2, i\kappa_j)$, and contains many operators that are not multiples of P.

(iii) *The heat kernel.* Let $u = t^{-1/2} e^{-x^2/(4Dt)}$ for $t > 0$ (up to the normalizing constant $(4\pi D)^{-1/2}$). Its x-derivatives are polynomials of growing degree times the same Gaussian, so they span an infinite-dimensional space and $u \notin \mathrm{EP}$ (Lemma 2.5). By the criterion $\mathrm{Ann}(u) = (P)$, and by Proposition 4.2 every constant-coefficient equation satisfied by the heat kernel is a multiple of $\tau - D\xi^2$.

Message. A "poor" solution (in EP) remembers only finitely many exponentials and cannot tell the heat equation from many others; a "rich" solution (outside EP) remembers the whole equation.

Proof of the easy direction of (a): $\Rightarrow$, by contraposition. Suppose $w_i \in \mathrm{EP}$ for some i.

1. By Lemma 2.5, $J = \mathrm{Ann}(w_i)$ has a finite zero set.
2. Since $V(P_i)$ is a curve (infinite), it contains a point at which some $q \in J$ does not vanish; hence q does not vanish identically on $V(P_i)$, and by (F3) $P_i \nmid q$ (if $P_i \mid q$ then q would vanish on $V(P_i)$; conversely if q vanished there, $q^k \in (P_i)$ and, P_i being irreducible, $P_i \mid q$).
3. Then $(P/P_i)\, q \in \mathrm{Ann}(u)$ because $(P/P_i) q\, u = q\, w_i = 0$;
4. but $(P/P_i) q \notin (P)$, since divisibility by P would force $P_i \mid q$.

So $\mathrm{Ann}(u) \supsetneq (P)$. □

The converse uses the structure of ideals (Lemma 2.3) and the Nullstellensatz, and is given in the project's proof package[E6].

> **Why it matters.** Theorem 4.1 turns "does u identify P?" into finitely many membership tests "$\in \mathrm{EP}$", each a finite-dimensional computation (Lemma 2.5). What remains unsettled is whether those tests can actually be carried out from the data one has; this is the subject of Section 4.3.

4.2 The spectral form

Theorem 4.1 says when (P) is the whole annihilator. Proposition 4.3 restates this through the two pieces of $\mathrm{Ann}(u)$ given by Lemma 2.3.

Notation

- $G_u = \gcd \mathrm{Ann}(u)$ is the "infinite part" of $\mathrm{Ann}(u)$, J_u is the cofactor ($\mathrm{Ann}(u) = G_u J_u$), and
$$\sigma_p(u) = V(J_u) \subset \mathbb{C}^2$$
is the *point spectrum* of u: a finite set of exponents (τ_0, ξ_0), i.e. of exponentials $e^{\tau_0 t + \xi_0 x}$ that are "present" in u. ("Spectrum" is used in an algebraic sense, not as the spectrum of an operator on a Hilbert space.)
- We write $G_u \sim P$ when G_u and P agree up to a nonzero constant factor.

Proposition 4.3 (Spectral characterization) (ledger P4-T2) [proved]

Let P be an annihilator of u. Then
$$\mathrm{Ann}(u) = (P) \iff \sigma_p(u) = \emptyset \text{ and } G_u \sim P.$$
Further $\mathcal{R}u \cap \mathrm{EP} \cong \mathcal{R}/J_u$.

Derivation of the equivalence. ($\Rightarrow$) If $\mathrm{Ann}(u) = (P)$, the gcd of the ideal is P, so $G_u \sim P$ and $J_u = \mathcal{R}$; then $\sigma_p(u) = V(\mathcal{R}) = \emptyset$.

($\Leftarrow$) If $\sigma_p(u) = V(J_u) = \emptyset$, the Nullstellensatz (F3) gives $1 \in J_u$, i.e. $J_u = \mathcal{R}$, hence $\mathrm{Ann}(u) = G_u\mathcal{R} = (G_u) = (P)$ because $G_u \sim P$.

The isomorphism $\mathcal{R}u \cap \mathrm{EP} \cong \mathcal{R}/J_u$ is not rederived here: it follows from $\mathcal{R}u \cong \mathcal{R}/\mathrm{Ann}(u)$ (Lemma 2.5) by passing to the finite-dimensional part (ledger P4-T2)[E11]. □

Illustration. For $u = e^{\tau_0 t + \xi_0 x}$ one has $\mathrm{Ann}(u) = (\tau - \tau_0, \xi - \xi_0)$, so $G_u = 1$ and $\sigma_p(u) = \{(\tau_0, \xi_0)\}$: a single exponential has no "infinite part" and a one-point spectrum. For $u = e^{\tau_1 t + \xi_1 x} + e^{\tau_2 t + \xi_2 x}$ with distinct exponents, $G_u = 1$ and $\sigma_p(u)$ has two points. For a solution of $P(\partial)u = 0$ with $\mathrm{Ann}(u) = (P)$ one has $\sigma_p(u) = \emptyset$ and $G_u \sim P$.

What G_u is good for

Thus G_u is the correct "operator intrinsic to u": it depends only on u, whereas P depends on which annihilator was chosen. It is recovered from *any* annihilator P by at most $\sum_i e_i$ membership tests "$\in$ EP" (ledger C3-T6) (one test for each copy of an irreducible factor that may have to be dropped; see Step 4 of the algorithm).

Why G_u and not the full variety

A tempting stronger formulation replaces G_u by the variety $V(\mathrm{Ann}\, u)$. It is correct only for reduced P, because zero sets do not see multiplicities.

Example: $P = \tau^2$ and $u = e^{x^2}$. Then $P(\partial)u = u_{tt} = 0$. Since $\partial_x^k e^{x^2}$ is a polynomial of degree exactly k times e^{x^2}, no nonzero polynomial in ∂_x alone kills u, hence

$$\mathrm{Ann}(u) = (\tau).$$

Here $\sigma_p(u) = \emptyset$ but $G_u = \tau \not\sim P$, so $\mathrm{Ann}(u) \neq (P)$, while $V(\mathrm{Ann}\, u) = \{\tau = 0\} = V(P)$: the variety criterion would wrongly say that u identifies τ^2[E12].

4.3 Limits of what can be certified

Theorem 4.1 is a statement about u, not about finite data. Table 3 says in which data situations the test of Theorem 4.1 can be carried out and in which it cannot, and what each case means for the algorithm, whose Step 3 is exactly this test (Section 8).

A few terms used in the table

Jet of order k	of u at a point is the list of all partial derivatives of u of order $\leq k$ at that point (its Taylor coefficients).
Zero-test oracle	is a procedure that, given an expression built from the data, tells whether it is identically zero.
Holonomic ideal	describes u by finitely many linear differential equations with polynomial coefficients whose solution space is finite-dimensional.
Weyl algebra	is the ring of differential operators with polynomial coefficients.

Situation	What can or cannot be done	What it means for the algorithm	Status
$u \in \mathrm{EP}$	a jet of order $2d-1$ at one point determines $\mathrm{Ann}(u)$; the threshold (a Bézout bound) cannot be improved in general (ledger C3-T3)[E8]	finite local data suffice: compute the jet at one point, solve a finite linear-algebra problem and return $\mathrm{Ann}(u)$ with a guarantee; the number of derivatives needed is known in advance	[proved]
$u \notin \mathrm{EP}$	no finite set of numbers (jets, samples) proves $\mathrm{Ann}(u) = (P)$, since every finite jet is realized by an exponential-polynomial (ledger C3-P1)	the algorithm can only *refute* "$u \in \mathrm{EP}$", never confirm the opposite: its output must read "not refuted at the given resolution", not "identified"	[proved] (jets); cited (Malgrange) for general data on convex sets
black box (jets plus a zero-test oracle)	"$w \in \mathrm{EP}$?" is undecidable (ledger P4-T5)	no general guarantee is possible; an extra hypothesis is needed (the next two rows)	[proved]
bound on $\dim(\mathcal{R}u \cap \mathrm{EP})$ known	decidable (ledger P4-T4)	the test of Theorem 4.1 becomes a finite computation and the algorithm can stop with a certain verdict	[proved]
u real-analytic with a known holonomic ideal over $\overline{\mathbb{Q}}$	decidable, if jets at an algebraic point are computable in a computable field (ledger OP-ALG-T1)	use the exact symbolic route (data model M3, Gröbner bases); decidable under the stated computability assumption	[proved mod. assumption (AC)]
variable coefficients (Weyl algebra)	restriction to the derivatives of one u never identifies anything (the algebra is simple) (ledger P4-C1); a finite-dimensional class of coefficients has a criterion with non-similar factors (ledger OP-ALG-T4)	do not apply the test as it stands; the algorithm is restricted to a fixed finite-dimensional class of coefficients	[proved]

Table 3: When the identification test can be carried out, and what each case means for the algorithm.

5 Decomposition into components

In this section

Starting from	the operator is known and factored, $P = \prod_j P_j^{m_j}$ with pairwise coprime $Q_j = P_j^{m_j}$; for the main theorem Ω is bounded, connected and simply connected.
Why this section	once P is known, to split u into the parts generated by each piece of P, so that the pieces can be analysed separately.
Role in the paper	second link of the chain. *Result:* the split exists for *factors* of the symbol (with a finite ambiguity) and fails for additive *terms*, which motivates Section 6.

5.1 Why per-term splitting fails

Proposition 5.1 (Terms do not give components) (ledger R2-T3)

If $L = L_1 + L_2$ (constant coefficients), then

$$\ker L \cap (\ker L_1 + \ker L_2) \;\subseteq\; \ker(L,\; L_1 L_2),$$

the common kernel of the two operators L and L_1L_2 (the kernel of the ideal they generate). This space is finite-dimensional as soon as $V(L_1) \cap V(L_2)$ is finite, because $V(L, L_1L_2) = V(L_1) \cap V(L_2)$.

[E9]

Proof. If $u = a + b$ with $L_1 a = 0$ and $L_2 b = 0$, then

$$L_1L_2u = L_2L_1a + L_1L_2b = 0$$

because the operators commute; and $Lu = 0$ by assumption.

For the varieties: at a point where $L_1L_2 = 0$, either $L_1 = 0$ and then $L = 0$ forces $L_2 = 0$, or $L_2 = 0$ and then $L_1 = 0$; conversely $L_1 = L_2 = 0$ gives $L = L_1L_2 = 0$. □

Example 5.1 (Heat equation)**.** Take $L = \partial_t - D\partial_x^2$ with $L_1 = \partial_t$ and $L_2 = -D\partial_x^2$ (the sign makes $L_1 + L_2 = L$).

- Solutions of $L_1u = 0$ do not depend on t:
$$\ker L_1 = \{f(x)\}.$$
- Solutions of $L_2u = 0$ are linear in x (integrate $u_{xx} = 0$ twice, the constants may depend on t):
$$\ker L_2 = \{a(t) + b(t)x\}.$$

So the elements of $\ker L_1 + \ker L_2$ have the form $f(x) + a(t) + b(t)x$, and all are killed by $\partial_t\partial_x^2$ (∂_x^2 kills $a + bx$ and ∂_t kills f). A heat solution is not: for $u = e^{-D\kappa^2 t}\cos\kappa x$,

$$\partial_x^2 u = -\kappa^2 u, \qquad \partial_t\partial_x^2 u = D\kappa^4 u \neq 0.$$

So a generic heat solution is **not** a sum of a solution of $\partial_t u = 0$ and one of $\partial_x^2 u = 0$.

Sharpness. The inclusion above cannot be sharpened to the common kernel $\ker(L_1, L_2)$: $u = x^2/2 + Dt$ lies in $\ker L_1 + \ker L_2$ and solves the heat equation ($u_t - Du_{xx} = D - D = 0$), but $L_1u = D \neq 0$. These exceptional solutions are polynomials and form a finite-dimensional space.

The consequence is structural: any attempt to represent u as a sum of independent contributions of the *terms* of the operator must leave a remainder (Section 6).

5.2 Why factors do work

Example 5.2 (Wave equation)**.** Let $c > 0$ be the wave speed ($c = 1$ in the figures) and

$$P = \tau^2 - c^2\xi^2 = (\tau - c\xi)(\tau + c\xi) =: Q_1Q_2, \qquad Q_1 = \partial_t - c\partial_x, \quad Q_2 = \partial_t + c\partial_x.$$

Direct check:

$$Q_1\, f(x + ct) = cf' - cf' = 0, \qquad Q_2\, g(x - ct) = -cg' + cg' = 0.$$

So $\ker Q_1 = \{f(x + ct)\}$ (a profile moving left) and $\ker Q_2 = \{g(x - ct)\}$ (a profile moving right), and d'Alembert's solution $u = f(x + ct) + g(x - ct)$ is exactly $\ker P = \ker Q_1 + \ker Q_2$.

The two pieces are ambiguous only up to a constant ($f \mapsto f + k$, $g \mapsto g - k$), reflecting $\dim \mathcal{R}/(Q_1, Q_2) = 1$:

$$Q_2 - Q_1 = 2c\,\xi, \qquad Q_1 + Q_2 = 2\tau \quad \Longrightarrow \quad (Q_1, Q_2) = (\tau, \xi), \qquad \mathcal{R}/(\tau, \xi) \cong \mathbb{C}$$

(Figure 2, left: the two lines meet in one point).

Why factors work and terms do not

One variable. If two polynomials in one variable are coprime, there are a, b with $aQ_1 + bQ_2 = 1$ (Bézout identity). Then any u with $Q_1Q_2u = 0$ splits explicitly as

$$u = bQ_2u + aQ_1u, \qquad bQ_2u \in \ker Q_1, \quad aQ_1u \in \ker Q_2,$$

and the sum of the two kernels is direct.

Two variables. "Coprime" only means that the common zeros are finitely many (F2), not that $(Q_1, Q_2) = \mathcal{R}$; the finite quotient $\mathcal{R}/(Q_1, Q_2)$ measures the failure of the one-variable picture and gives the ambiguity (Theorem 5.2).

Terms. Per-term splitting fails for a different reason: the terms L_1, L_2 are not factors of L, and the sum $\ker L_1 + \ker L_2$ lies in the kernel of the *different* operator L_1L_2 (Proposition 5.1).




Figure 3: Left: d'Alembert solution at $t = -2, 0, 2$ with $c = 1$; $f(x+ct) = 1/(1+(x+ct)^2)$ (a lorentzian bump) moves left, $g(x-ct) = e^{-(x-ct)^2}$ (a Gaussian) moves right. Right: the two families of characteristics, one per factor Q_1, Q_2. Each factor of the symbol carries one rigidly translating component.

Theorem 5.2 (Factor decomposition) (ledger C12-T1′) [proved]

Let $\Omega \subset \mathbb{R}^2$ be bounded, connected and simply connected, and $P = P_1^{m_1} \cdots P_N^{m_N}$ with $Q_j = P_j^{m_j}$ pairwise coprime. Then every $u \in \ker P$ (in $\mathcal{D}'$ or C^∞) can be written

$$u = \sum_{j=1}^{N} u_j, \qquad u_j \in \ker Q_j(\partial),$$

and the ambiguity of the decomposition is an exponential-polynomial space of dimension exactly

$$\delta(P) = \sum_{i<j} \dim \mathcal{R}/(Q_i, Q_j).$$

Intuition

This is the Chinese remainder theorem (concrete examples in Appendix B, §B.3; see Figure 2, left, for the geometric overlap). For integers coprime to each other, $\mathbb{Z}/ab \cong \mathbb{Z}/a \times \mathbb{Z}/b$. For polynomials in two variables, "coprime" does not imply $(Q_i, Q_j) = \mathcal{R}$: the ideal may have a finite zero set (the points where the curves $Q_i = 0$ and $Q_j = 0$ cross, see Figure 2), and the finite quotient $\mathcal{R}/(Q_i, Q_j)$ measures the overlap. Algebraically the proof uses the exact sequence

$$0 \to \mathcal{R}/(Q_1Q_2) \to \mathcal{R}/(Q_1) \oplus \mathcal{R}/(Q_2) \to \mathcal{R}/(Q_1, Q_2) \to 0$$

and the fact that $C^\infty(\Omega)$ is an injective module (a consequence of the Malgrange–Ehrenpreis theorem [2, 3]), which turns it into an exact sequence of function spaces.

Why it matters. A "component" of u is a *factor* of the symbol, not a term of the operator. The overlap between components is finite-dimensional and exactly measured by $\delta(P)$.

5.3 An explicit projector for evolution factors

For an evolution operator with $P = \prod_{j=1}^{N}(\tau - a_j(\xi))$ and distinct a_j, a component is an eigenfunction of ∂_t: $\partial_t u_j = a_j(\partial_x)u_j$. Applying $\prod_{k\neq j}(\partial_t - a_k(\partial_x))$ to $u = \sum_l u_l$ kills every u_l with $l \neq j$ (it contains the factor $\partial_t - a_l$) and multiplies u_j by $\prod_{k\neq j}(a_j - a_k)(\partial_x)$. Hence

$$u_j = \prod_{k\neq j} \frac{\partial_t - a_k(\partial_x)}{a_j(\partial_x) - a_k(\partial_x)}\, u, \tag{2}$$


the Lagrange formula of Eq. (1) with C replaced by ∂_t. The result is a Fourier multiplier in x that is rational in ξ, with poles where $a_j(\xi) = a_k(\xi)$; this matches the characterization "projectors rational in ξ iff the factors are coprime" (ledger II-T2/T3).

5.4 Where decomposition fails

Proposition 5.3 (Holes) (ledger C12-P10) [proved]

On a domain with a hole, the decomposition fails for some P.

Figure 4 shows the standard obstruction. In the theorem the hypothesis "simply connected" means $H^1_{dR}(\Omega) = 0$, exactly what removes this obstruction; a cohomological criterion for non-simply connected Ω is also available (ledger OP-FUN, C12-Q6).



$u = \log|z-\alpha|^2$ solves $\Delta u = 0$ on the annulus, but $u \neq h(z) + \overline{g(z)}$ with h, g single-valued holomorphic: the logarithm has monodromy (winding number $\neq 0$).

Figure 4: The Laplacian $\partial_x^2 + \partial_y^2 = (\partial_x + i\partial_y)(\partial_x - i\partial_y)$ factors, but the factor decomposition fails on a punctured disc: the obstruction is topological ($H^1_{dR} \neq 0$).

5.5 Regularity: when are the components in H^m?

The decomposition is an algebraic statement; analysis enters when we want components in a Sobolev space. For two components the governing condition is geometric: the two component spaces M_1 and M_2 must be separated by a positive angle (Figure 5); the next theorem makes this precise for any number of components. Two difficulties appear along the way; they are collected in the box after the theorem.

Proposition 5.4 (Loss of a derivative) (ledger C12-C1) [false]

"If $u \in H^m(\Omega)$ then all $u_j \in H^m$" is false: one derivative can be lost. In other words, the pieces of a smooth function may be rougher than the function itself.

Theorem 5.5 (Regularity criterion) (ledger C12-T3, C12-P7) [proved]

All components lie in H^m for every $u \in \ker_{H^m} P$ iff the sum $\sum_j \ker_{H^m} Q_j$ is *closed* and *dense* in $\ker_{H^m} P$. Closedness is equivalent to $\gamma_N > 0$, where

$$\gamma_N = \inf \sigma\big(S^*S\big|_{(\ker S)^\perp}\big),$$
$$S : \bigoplus_{j=1}^{N} \ker Q_j \to H^m, \qquad S(u_1, \dots, u_N) = \sum_{j=1}^{N} u_j.$$

For $N = 2$ this is the condition $c_F < 1$ on the Friedrichs cosine. Density holds on every bounded convex Ω, for every P and m (ledger OPF5B-T1) [proved mod. Hörmander [5], Stein].

Reading γ_N

- The space $\bigoplus_j \ker Q_j$ consists of the N-tuples $v = (u_1, \dots, u_N)$ with norm $\|v\|^2 = \sum_j \|u_j\|^2_{H^m}$;
- $Sv = \sum_j u_j$ is their sum;
- S^* is the adjoint of S;
- $(\ker S)^\perp$ removes the tuples that sum to zero (the ambiguity of Theorem 5.2);
- $\inf \sigma$ is the bottom of the spectrum.

Since S^*S restricted to $(\ker S)^\perp$ is self-adjoint and nonnegative, the bottom of its spectrum is the minimum of the Rayleigh quotient

$$\frac{\langle S^*Sv, v\rangle}{\|v\|^2} = \frac{\|Sv\|^2}{\|v\|^2}.$$

Hence the formula of the theorem is equivalent to the more intuitive

$$\boxed{\gamma_N = \inf_{\substack{v \in (\ker S)^\perp \\ v \neq 0}} \frac{\|Sv\|^2_{H^m}}{\|v\|^2}}$$

In words

γ_N is the worst case, over all tuples of pieces, of the ratio between the size of the sum and the size of the pieces.

- If $\gamma_N > 0$ the pieces cannot nearly cancel each other, and
$$\|v\| \le \gamma_N^{-1/2} \|Sv\| :$$
the pieces are controlled by the sum, so recovering them from u is stable.
- If $\gamma_N \to 0$ there are large pieces whose sum is tiny, and the recovery is unstable.

The case $N = 2$. One finds $\gamma_2 = 1 - c_F$, where

$$c_F = \sup\{|\langle a, b\rangle| : \ a \in M_1 \ominus (M_1 \cap M_2),\ b \in M_2 \ominus (M_1 \cap M_2),\ \|a\| = \|b\| = 1\}$$

is the Friedrichs cosine: the part of v lying in $M_1 \cap M_2$ contributes the ratio 2 and does not affect the infimum, and on the rest

$$\|a + b\|^2 \ge (1 - c_F)(\|a\|^2 + \|b\|^2)$$

with near-equality attained. Hence $c_F < 1 \iff \gamma_2 > 0$, as stated.




Figure 5: Friedrichs angle θ between two component spaces $M_1 = \ker Q_1$ and $M_2 = \ker Q_2$. A larger angle means a better-conditioned recovery of the pieces from their sum.

For $N = 2$, Figure 5 reads as follows: $c_F = \cos\theta$; if $\theta > 0$ the sum $M_1 + M_2$ is closed and the pieces depend continuously on u; as $\theta \to 0$ the split becomes unstable. For $N \geq 3$ the pairwise angles do not suffice; γ_N, the smallest nonzero eigenvalue of S^*S, is the right conditioning number: it is the conditioning of the linear map "components $\mapsto$ sum", exactly like the smallest nonzero singular value in a least-squares problem.

Two problems with regularity

Problem 1: stability [false]. Algebraically the split always exists (Theorem 5.2), but the pieces may be less regular than u. When the angle between the component spaces tends to 0 (that is, $\gamma_N \to 0$) the map "$u \mapsto$ pieces" becomes unstable, and one derivative can be lost (Proposition 5.4). For the algorithm: check $\gamma_N > 0$ before using the components in H^m (Step 5).

Problem 2: three or more components [open]. For $N \geq 3$ the pairwise cosines $c_F(i,j)$ do not decide whether the sum of kernels is closed: an abstract Hilbert-space counterexample exists (ledger S-CHECK-4), and whether it can occur for differential operators is open (ledger C12-Q2).

Why it matters. Algebraically the split always exists (Theorem 5.2); analytically, the components are well behaved exactly when the component spaces are separated by a positive angle (Theorem 5.5). The next section stops asking for a split into factors and asks what survives when one splits by additive terms instead.

6 Terms inside a factor and the interaction term

In this section	
Starting from	an evolution equation $u_t = Au$, $A = A_1 + \cdots + A_N$ (the terms), with flows $S_n(t) = e^{tA_n}$; the terms commute, except in the last subsection of this section (non-commuting terms).
Why this section	Proposition 5.1 shows that splitting by terms is never exact; here we compute exactly what is left over (the interaction term E_{int}), its cost, and its stability.
Role in the paper	third link of the chain, answering Q3. *Result:* an exact product formula, a cost dichotomy $O(N)$ versus $\Theta(2^N)$, and a negative stability theorem.

Here the equation is an evolution $u_t = Au$ with $A = A_1 + \cdots + A_N$, each A_n a *term* (diffusion, advection, reaction, ...).

From summing operators to the interaction term

The solution is

$$u(t) = e^{tA}u_0 = S(t)u_0,$$

where $S(t) = e^{tA}$ is the *flow* (semigroup) generated by A. If term n acted alone, it would produce $S_n(t)u_0 = e^{tA_n}u_0$, a change of $S_n(t)u_0 - u_0$.

The naive guess is that the terms act independently, so that the total change $u(t) - u_0$ is the sum of the individual changes. It is not: the flow of a sum is not the sum of the flows. For advection ($A_1 = -c\partial_x$, a translation) and diffusion ($A_2 = D\partial_x^2$, a smoothing) the solution is translated *and* smoothed, and this joint action is not "only translated" plus "only smoothed". What is missing is the *interaction term* E_{int} defined next.

The section proceeds in three steps:

1. define E_{int} (6.1);
2. compute it exactly when the terms commute, using Fourier variables (6.2);
3. measure its cost (6.3) and its stability (6.4).

Subsections 6.5 and 6.6 treat the cases without a closed formula.

6.1 Definition via the flows of the single terms

Let $S(t) = e^{tA}$ and $S_n(t) = e^{tA_n}$. The contribution that term n *alone* would make to the solution is $S_n(t)u_0 - u_0$. The remainder is the interaction term

$$E_{\text{int}}^{\text{add}}(t) = u(t) - u_0 - \sum_{n=1}^{N} \big(S_n(t)u_0 - u_0\big). \tag{3}$$

(The superscript "add" records that this is the *additive* definition; below we write E_{int} for $E_{\text{int}}^{\text{add}}$ unless the distinction matters.)

6.2 Exact formula in the commuting case

Why constant coefficients commute, and what Δ_n is

Constant-coefficient operators are polynomials in ∂_t, ∂_x (Section 2) and the ring $\mathcal{R}$ is commutative, so any two terms commute: $A_1A_2 = A_2A_1$. For commuting operators

$$e^{t(A_1+A_2)} = e^{tA_1}e^{tA_2}, \qquad \text{hence} \qquad S = \prod_n S_n;$$

without commutativity this fails (Section 6.6).

We call

$$\Delta_n = S_n(t) - \mathrm{Id}$$

the *increment* produced by term n alone, where Id is the identity operator and t is fixed; thus $S_n = \mathrm{Id} + \Delta_n$, and $\Delta_n u_0 = S_n(t)u_0 - u_0$ is exactly the contribution of term n in (3). Products of operators are compositions.

Fourier variables

For constant coefficients it is convenient to pass to Fourier variables. With

$$\hat{u}(\kappa) = \int u(x)e^{-i\kappa x}\,\mathrm{d}x,$$

where $\kappa \in \mathbb{R}$ is the frequency:

- the derivative ∂_x becomes multiplication by $i\kappa$;
- each term A_n becomes multiplication by its *symbol* $a_n(\kappa)$ (a polynomial in $i\kappa$);
- e^{tA_n} becomes multiplication by the number $e^{ta_n(\kappa)}$.

Operators turn into numbers and compositions into products of numbers, so for a fixed κ every computation below is scalar.

Theorem 6.1 (Interaction term) (ledger P1-T2, P1-T3) [proved]

Assume the A_n commute (e.g. constant coefficients). Write $\Delta_n = S_n(t) - \mathrm{Id}$. Then

$$E_{\text{int}}^{\text{add}} = \Big[\prod_{n=1}^{N}(\mathrm{Id}+\Delta_n) - \mathrm{Id} - \sum_{n=1}^{N}\Delta_n\Big]u_0. \tag{4}$$

For $N = 2$ this is simply $E_{\text{int}} = \Delta_1\Delta_2 u_0 = (S_1 - \mathrm{Id})(S_2 - \mathrm{Id})u_0$. In Fourier variables, with $\hat{A}_n = a_n(\kappa)$,

$$\widehat{E_{\text{int}}}(t,\kappa) = \Big[\prod_n \big(e^{ta_n(\kappa)}\big) - 1 - \sum_n \big(e^{ta_n(\kappa)} - 1\big)\Big]\hat{u}_0(\kappa).$$

Proof. Commutation gives $S = e^{t\sum A_n} = \prod_n S_n$. Substitute $S_n = \mathrm{Id} + \Delta_n$ in (3). For $N = 2$, expand the product:

$$\begin{aligned} S_1S_2 &= (\mathrm{Id}+\Delta_1)(\mathrm{Id}+\Delta_2) \\ &= \mathrm{Id} + \Delta_1 + \Delta_2 + \Delta_1\Delta_2, \end{aligned}$$

and $\sum_n(S_n - \mathrm{Id}) = \Delta_1 + \Delta_2$, so the remainder in (3) is $\Delta_1\Delta_2 u_0$.

For general N, the Δ_n commute, so

$$\prod_n(\mathrm{Id}+\Delta_n) = \sum_{S\subseteq\{1,\dots,N\}}\prod_{n\in S}\Delta_n;$$

removing the subsets with $|S| = 0$ and $|S| = 1$ leaves

$$E_{\text{int}} = \sum_{|S|\geq 2}\Delta_S\, u_0, \qquad \Delta_S = \prod_{n\in S}\Delta_n.$$

In words: the interaction is the sum of products of at least two increments, and appears only when at least two terms act together. □

A concrete case. Take advection–diffusion, $u_t = -cu_x + Du_{xx}$, with Gaussian datum $u_0 = g(x;\sigma^2)$, where $g(x;v) = e^{-x^2/2v}/\sqrt{2\pi v}$. The two flows act as a translation and a Gaussian smoothing:

$$S_1u_0 = g(x-ct;\sigma^2), \qquad S_2u_0 = g(x;s^2), \qquad u = g(x-ct;s^2), \qquad s^2 = \sigma^2 + 2Dt.$$

Hence

$$E_{\text{int}} = g(x-ct;s^2) - g(x-ct;\sigma^2) - g(x;s^2) + g(x;\sigma^2).$$

Figure 6(a) plots it: it is comparable in size to u itself, so it cannot be discarded as a small correction. In Fourier variables the ratio $|\widehat{E_{\text{int}}}|/|\hat{u}_0| = |(e^{-ic\kappa t} - 1)(e^{-D\kappa^2 t} - 1)|$ is at most 2; it vanishes at $\kappa = 0$ and whenever $c\kappa t \in 2\pi\mathbb{Z}$ (the translation returns to the identity), see Figure 6(b).




Figure 6: Advection–diffusion with Gaussian datum ($\sigma^2 = 1$, $c = 2$, $D = \frac{1}{2}$). (a) u_0, $u(t)$ and E_{int} at $t = 1$. (b) The Fourier ratio $|\widehat{E_{\text{int}}}|/|\hat{u}_0|$ for three times.

6.3 Cost: $O(N)$ versus 2^N

Expanding (4) in the Δ_n produces one monomial per subset of $\{1, \dots, N\}$ with at least two elements: $2^N - N - 1$ cross terms (Figure 7: the red nodes are the cross terms). Writing the *complete table* of cross terms is therefore exponential. Evaluating a *value* is cheap, because we never expand the product. The algorithm uses exactly the product (Fourier) form of Theorem 6.1,

$$\widehat{E_{\text{int}}}(t, \kappa) = \Big[\prod_{n=1}^{N} e^{t a_n(\kappa)} - 1 - \sum_{n=1}^{N} \big(e^{t a_n(\kappa)} - 1 \big) \Big] \hat{u}_0(\kappa),$$

which for a fixed κ needs the N numbers $e^{t a_n(\kappa)}$, $N - 1$ products and N subtractions: $O(N)$ operations once $\hat{u}_0(\kappa)$ is known.

> **Two remarks on cost**
>
> **What "value" means.** Here "value" means one Fourier value; recovering $E_{\text{int}}(t, x)$ at a point of space needs an inverse Fourier transform, whose cost is not counted in the $O(N)$.
>
> **Where 2^N comes from.** The count $2^N - N - 1$ refers only to writing out all cross terms one by one; the product form never does so (Step 6 of the algorithm).




Figure 7: Subset lattice for $N = 3$ terms: one node per subset S of terms, one monomial Δ_S per node with $|S| \geq 2$. The count $2^N - N - 1$ is what makes the full table exponential.

Proposition 6.2 (Complexity) (ledger P1-T3) [proved]

In the commuting exponential-type class, a value of E_{int} costs $O(M(N+\sum d))$ for M evaluation points; aggregated by order $O(MN^2)$; the full table $\Theta(M2^N(L+1)^2)$.

[E1]

6.4 Why E_{int} cannot be removed or regularized stably

The cost question being settled, we ask whether E_{int} can be suppressed by a regularization, or at least depends continuously on the datum.

Proposition 6.3 (Not continuous in H^m) (ledger P1-P1(e)) [false]

The map $u_0 \mapsto E_{\text{int}}(t)$ is *not* continuous in $H^m(\Omega)$ for advection–diffusion, for any m and any t; the same holds for any two non-proportional terms (ledger P1-Q6).

Consequently:

- **A sparsity (or L^2) penalty cannot make E_{int} vanish.** Outside a finite-dimensional space of exponential-polynomials there is no representation of u as a sum of per-term pieces (ledger P2-T2) [proved]. For advection–diffusion this space is span$\{1,\ x-ct,\ (x-ct)^2+2Dt\}$: three functions.
- **No stable regularization** of the interaction exists (ledger P2-T4): it is a bilinear quantity built from unbounded operators, and no uniformly Lipschitz regularization of such maps exists.
- The only *continuous* splitting found is a balance convention (ledger P3-T6) which requires an external principle to say how to distribute the cross terms.

6.5 Time-dependent perturbation theory

Why this subsection

The product formula (4) needs commuting terms. When terms do not commute, or when boundary conditions are imposed, there is no closed formula, but E_{int} is still defined by (3). Perturbation theory expands the flow in powers of the interaction. From this expansion one reads

(i) that E_{int} is well defined for all times (convergence),
(ii) approximations obtained by truncating the series order by order, and
(iii) the size of E_{int} for small t.

Split $A = A_1 + B$, where A_1 is the principal term (for instance diffusion) and B, the remaining terms, is treated as a perturbation of A_1. The Dyson–Phillips series is

$$\begin{aligned} S(t) &= \sum_{k\geq 0} S^{(k)}(t), \\ S^{(0)}(t) &= e^{tA_1}, \\ S^{(k+1)}(t) &= \int_0^t e^{(t-s)A_1}\, B\, S^{(k)}(s)\, \mathrm{d}s, \end{aligned} \tag{5}$$

the same expansion that gives the interaction picture of quantum mechanics [8].

Reading the series

- $S^{(0)} = e^{tA_1}$ is the flow of A_1 alone;

- $S^{(k)}$ collects the contributions in which B acts k times, at intermediate times $s \in (0,t)$, with the flow of A_1 in between.

The terms with $k \geq 1$ are therefore the part of the flow in which B acts on the evolution generated by A_1, that is, the interaction. (When A_1 and B commute, the first-order term is

$$\int_0^t e^{(t-s)A_1} B e^{sA_1}\, \mathrm{d}s = t\, B e^{tA_1},$$

consistent with Section 6.)

What it gives. The series converges for all t when A_1 generates an even-order dissipative semigroup (e.g. diffusion) and B is a lower-order L^∞ perturbation, also with Dirichlet conditions (ledger P3-T1) [proved mod. Miyadera–Voigt [7]]: so E_{int} is well defined and can be approximated by truncation even without a closed formula. Each $S^{(k)}$ is the order-k part of the interaction. For small times, for boundary-induced interaction, with k the order of contact of the datum with the boundary, $\|E_{\text{int}}\| \sim t^{k/2+3/4}$ (ledger P3-T3, OP-F5A) [proved][E10]: the smoother the datum is against the boundary (larger k), the faster the interaction becomes negligible as $t \to 0$.

6.6 Non-commuting terms

If $[A_1, A_2] \neq 0$, then $e^{t(A_1+A_2)} \neq e^{tA_1}e^{tA_2}$; by Baker–Campbell–Hausdorff,

$$e^{tA_1}e^{tA_2} = \exp\Big(t(A_1+A_2) + \tfrac{t^2}{2}[A_1,A_2] + O(t^3)\Big).$$

Proposition 6.4 (Lie-algebra dichotomy) (ledger P1-T4, P1-T5) [proved]

If the Lie algebra generated by the terms is solvable, an exact formula with finitely many scalar functions exists (Wei–Norman). Otherwise only a local expansion exists, and **no finite product of the flows of the individual terms** reproduces $e^{t(X+\varepsilon Y)}$.

Why it matters. For commuting terms the interaction is explicit and cheap to evaluate, but it cannot be made to vanish nor be computed stably in H^m; for non-commuting terms even the explicit form is lost in general. These are the linear-case limits against which the nonlinear extension of Section 7 must be read.

7 Nonlinear equations

In this section	
Starting from	a scalar equation in one space variable, $u_t = \sum_n T_n(u)$, each T_n linear or polynomial in the jet $(u, u_x, u_{xx}, \dots)$; Ω bounded and simply connected; the exact solution is known.
Why this section	to test which of the notions of Sections 4–6 survive when the terms are nonlinear.
Role in the paper	extension of the chain. *Result:* weaker, relative statements (exact expansion of E_{int}, sharp asymptotics across a shock for Burgers, identification relative to a dictionary). Benchmarks: Burgers, KdV–Burgers, Fisher–KPP; KdV and NLS only as controls.

We consider $u_t = \sum_n T_n(u)$ with each T_n linear or polynomial in the jet $(u, u_x, u_{xx}, \dots)$. Benchmarks: Burgers $u_t = -uu_x + \nu u_{xx}$, KdV–Burgers, Fisher–KPP; integrable cases (KdV, NLS) serve as controls.

7.1 Interaction term for flows of single terms

With S_n the (nonlinear) flow of $u_t = T_n(u)$ alone, (3) still defines $E_{\text{int}}^{\text{add}}$. Its Taylor expansion in t:

Theorem 7.1 (Jet expansion) (ledger S4T-T1) [proved]

$E_{\text{int}}^{\text{add}} = \dfrac{t^2}{2} \sum_{n \neq m} T_n'[\, T_m(u_0)\,] + O(t^3)$, where $T_n'[v]$ is the Fréchet derivative of T_n at u_0 in the direction v.

Example 7.1 (Burgers). $T_1 = -uu_x$, $T_2 = \nu u_{xx}$. Then $T_1'[v] = -(vu_x + uv_x)$ and $T_2'[v] = \nu v_{xx}$, so

$$T_1'[T_2 u_0] = -\nu\,(u_0'' u_0' + u_0 u_0'''),$$
$$T_2'[T_1 u_0] = -\nu\,(3u_0' u_0'' + u_0 u_0'''),$$

and half the sum times t^2 gives

$$\begin{aligned} E_{\text{int}} &= \tfrac{1}{2}\big(T_1'[T_2 u_0] + T_2'[T_1 u_0]\big)t^2 + O(t^3) \\ &= -\nu\big(u_0 u_0''' + 2u_0' u_0''\big)t^2 + O(t^3). \end{aligned}$$

7.2 Before, at, and after the shock

The interaction term of Burgers changes qualitatively with time, and each regime calls for different tools:

Before shock formation smoothness permits perturbative expansions and derivative-based estimates.

At breaking the inviscid gradient blows up, invalidating estimates that require smoothness.

After shock formation pointwise behavior can be discontinuous, so L^1 estimates become more informative than pointwise bounds.

Figure 8 recalls why a shock appears at all in the flow of the single term $-uu_x$: along each characteristic u is constant and the characteristic moves with speed u, so faster parts overtake slower ones. For $u_0 = \sin x$ the characteristics are

$$x = \xi + t \sin \xi$$

(the parameter s in the figure plays the role of the time t); since

$$\frac{\partial x}{\partial \xi} = 1 + t \cos \xi$$

vanishes first at $\xi = \pi$, the breaking time is $t^* = 1$.

Before the shock ($t < t^*$): smooth regime. Smoothness permits perturbative expansions and derivative-based estimates.

- E_{int} has a closed form at first order in ν, with remainder $O(\nu^2 \varepsilon^2 t^3)$ (ledger S4T-T2) [proved]; it blows up at the breaking time t^* (ledger S4T-P3). (This is the same family of expansions as Theorem 7.1 and Example 7.1.)
- The Cole–Hopf substitution $u = -2\nu\varphi_x/\varphi$ turns Burgers into the heat equation $\varphi_t = \nu\varphi_{xx}$; the inverse scattering transform linearizes KdV and NLS. These are used as the "exact linearization" of the nonlinear solution [9].




Figure 8: Characteristic solution of the inviscid flow of $-uu_x$. At $s = 1$ the profile has a vertical tangent: the gradient blows up. Beyond that the curve folds; the physical (entropy) solution replaces the fold by a jump at $x = \pi$ (equal-area rule), shown dashed. This breaking is what makes E_{int} discontinuous after the shock.

At breaking ($t = t^*$): estimates that need smoothness fail. The inviscid gradient u_x blows up (vertical tangent, $s = 1$ in Figure 8), so every statement that needs bounds on $u_x, u_{xx}, \dots$ stops being usable at t^*: the closed form of E_{int} blows up, and estimates that require smoothness, such as those behind the jet expansion of Theorem 7.1, are no longer valid.

After the shock ($t > t^*$): discontinuous regime. The flow of $-uu_x$ alone forms a shock, so E_{int} jumps by amounts independent of ν; this is forced by the definition. A function with a jump is not in H^1 in one dimension, so the Sobolev norms of Sections 5 and 6 are no longer the natural ones: pointwise values are unreliable, while the *measure* of the discrepancy is well behaved. The meaningful quantities are therefore L^1 sizes, collected below:

$$\|E_{\text{int}}\|_{L^1} = \Theta(\nu), \tag{6}$$

$$\sup_{t \le T} \|E_{\text{int}}\|_{L^1} = \nu |\log \nu| + O(\nu), \tag{7}$$

$$\|S - S_1 S_2\|_{L^1} = \Theta(\sqrt{\nu}). \tag{8}$$

The first line holds at fixed t, each shock contributing $4\nu \log\big(2\cosh(\rho/2)\big)$; the second has exact leading coefficient 1. Sources: the first line is (ledger OPA-T1) [proved mod. Kružkov uniqueness [10]]; the second is (ledger OPF5A-T1/T2); the third is (ledger OPF5D-T6) [proved]. The second line holds for data with a single birth of shock such as $\sin x$, and is conditional on a localization lemma in general [conditional].

Regime	What happens	Tools and results
Before the shock, $t < t^*$	the profile is smooth, characteristics do not cross	jet expansion (Thm. 7.1), first-order closed form of E_{int}, Cole–Hopf
At breaking, $t = t^*$	the inviscid gradient blows up (vertical tangent)	smoothness-based estimates fail; E_{int} blows up
After the shock, $t > t^*$	the single-term flow is discontinuous (equal-area rule)	L^1 estimates: $\Theta(\nu)$, $\nu\lvert\log\nu\rvert$, $\Theta(\sqrt{\nu})$

Table 4: The three regimes of the Burgers interaction term and the tools that apply in each.

7.3 Identification relative to a dictionary

Nonlinear terms cannot be handled by the ideal $\mathrm{Ann}(u)$ of Section 2; instead one fixes a finite candidate list. Choose a dictionary $\mathcal{D}$ of monomials of degree $\le d$ in the jet, of order $\le k$, and ask which polynomial relations $F(u, u_x, \dots) = 0$ the data satisfy. Notation: V is the span of $\mathcal{D}$, $I(u)$ the ideal of all differential relations satisfied by u, $[F]$ the differential ideal generated by F, and $\mathcal{J}_u$ the jet-level obstruction (a different object from the cofactor J_u of Section 2, hence the calligraphic letter)[E13].

Theorem 7.2 (Dictionary criterion) (ledger S4ID-T1) [proved]

For smooth u on any Ω there is an exact sequence $0 \to [F] \cap V \to I(u) \cap V \to \mathcal{J}_u \cap \psi(V) \to 0$. In particular

$$I(u) \cap V = [F] \cap V \iff \mathcal{J}_u \cap \psi(V) = 0.$$

The correct statement is therefore "the relations are exactly F *modulo its differential consequences*"; the literal "$I(u) \cap V = \langle F \rangle$" is false in complete dictionaries (they contain $\partial_x F$, etc.) (ledger S4ID-P1). Concrete criteria: for Burgers with N shocks, u identifies the equation in the evolutionary dictionary iff $d \le N$ (ledger S4ID-T2, OP-T7); for N-soliton KdV the identifying polynomial has total degree $(N+2)2^{N-1}$ (generic $N = 2$: 8) (ledger OP-K15) [proved mod. Novikov 1974 [11]]. Finally, "closed form $\Rightarrow$ does not identify" is itself a theorem (ledger S4ID-T4). [E2]

7.4 Linear versus nonlinear: the same idea, two structures

Sections 2–6 and the present one answer the same question: *what does a single solution u remember about its equation?* In both settings we collect all relations satisfied by u and ask when these relations are generated by the equation we were given. What changes is the algebraic structure in which the relations live.

	Linear approach (Sections 2–6)	Nonlinear approach (this section)
Space of candidate equations	the ring $\mathcal{R} = \mathbb{C}[\tau, \xi]$ of constant-coefficient operators	a finite *dictionary* $\mathcal{D}$ of jet monomials of degree $\le d$ and order $\le k$; $V = \mathrm{span}\,\mathcal{D}$
Relations satisfied by u	$\mathrm{Ann}(u)$, an ideal of $\mathcal{R}$	$I(u)$, the ideal of all differential relations
What the equation generates	the principal ideal (P)	the differential ideal $[F]$
Criterion for identification	$\mathrm{Ann}(u) = (P)$ (Theorem 4.1)	$I(u) \cap V = [F] \cap V$, i.e. F *modulo its differential consequences* (Theorem 7.2)
What escapes the equation	the cofactor J_u: the finitely many exponentials in u ($\sigma_p(u) = V(J_u)$)	the jet-level obstruction $\mathcal{J}_u \cap \psi(V)$ (exact sequence of Theorem 7.2)
Solutions that do not identify	$u \in \mathrm{EP}$	closed-form solutions ((ledger S4ID-T4))
Concrete criteria	finite linear algebra (Lemma 2.5)	Burgers with N shocks identifies iff $d \le N$; N-soliton KdV: degree $(N+2)2^{N-1}$

Table 5: The linear and nonlinear identification problems side by side.

Where the parallel breaks

Linear case. The solutions form a vector space and the annihilators an ideal: one can add, multiply by operators and factor, which is why the Nullstellensatz applies and the test is a finite-dimensional computation.

Nonlinear case. There is no linear structure on the solutions: one must fix a finite space V of candidates in advance, and the identification is *relative to* V, never absolute. The literal statement

$$I(u) \cap V = \langle F \rangle$$

is false in complete dictionaries, which contain $\partial_x F$, so the correct statement is "modulo differential consequences".

Takeaway. Both approaches ask what one solution determines about its equation. The linear one answers with the annihilator ideal $\mathrm{Ann}(u)$ and its generator G_u; the nonlinear one answers with a dictionary and the differential ideal of relations, and only up to differential consequences. In the algorithm (Section 8) Steps 1–6 use the structure of $\mathrm{Ann}(u)$, while Step 7 switches to the dictionary and returns a relative identification.

8 The algorithm

In this section	
Starting from	input of one of the data models M1–M3 below, and for each step the hypotheses of the theorem that justifies it.
Why this section	to turn the theorems into one procedure with explicit inputs, outputs, and guarantees.
Role in the paper	operational summary of Sections 4–7. Table 6 lists the steps, Figure 9 the control flow, and a worked example makes them concrete.

8.1 Inputs and data models

The reconstruction is exact, hence the input must be of an exact kind. Three data models were analysed (ledger O7):

M1 closed-form u (e.g. sums of exponentials and Gaussians);
M2 the jet of u at a point, to a given order;
M3 u real-analytic on a connected Ω, given through a holonomic ideal with coefficients in $\overline{\mathbb{Q}}$ (assumption (AC) on computing jets).

In model M1 the integral-ring identification never holds (ledger C3-C2), and relative identification is used.

8.2 The pipeline

Step	Task	Result relied upon	Output
0	Input	data models M1–M3	exact u
1	Annihilator	jet Gram matrix kernel	candidate P
2	Factor	unique factorization (F1)	irreducible factors P_i, e_i
3	EP test	Lemma 2.5, Thm. 4.1	verdict per factor
4	Reduce	Prop. 4.3, ledger C3-T6	G_u, J_u, certificate
5	Decompose	Thm. 5.2, Eq. (2)	components u_j, ambiguity $\delta(P)$
6	Terms	Thm. 6.1	log-symbol, E_{int}
7	Nonlinear branch	Thm. 7.2, Sec. 7	relative identification, flow-based E_{int}

Table 6: The steps of the algorithm at a glance: what each step does, which result justifies it, and what it returns.

8.3 The steps in detail

Step 1: obtain an annihilator

Does Let $\Phi(u) = (T_1 u, \dots, T_M u)^\top$ be the jet over a dictionary of M candidate terms and $\mathbb{G}(u) = \int_\Omega \Phi\Phi^* \, \mathrm{d}\mu$ its Gram matrix (positive semidefinite). A relation $\lambda \cdot \Phi = 0$ holds iff $\lambda \in \ker \mathbb{G}(u)$.

How Exact arithmetic (rational or algebraic coefficients), Gaussian elimination: no learning, no statistics.

Returns a candidate P. If $\ker \mathbb{G}$ has dimension > 1, u satisfies several dictionary relations, which is itself information about G_u and J_u.

[E3]

Step 2: factor

Does factor $P(\tau, \xi) \in \overline{\mathbb{Q}}[\tau, \xi]$ into irreducibles with multiplicities (a standard computer-algebra operation).

Returns the factors P_i and exponents e_i.

Step 3: exponential-polynomial test

Does for each i compute $w_i = (P/P_i)(\partial)u$ and decide whether $\dim \mathcal{R} w_i < \infty$; by Theorem 4.1(a) this is exactly the identification test.

How depends on the data model:

- M1: a syntactic check (is the closed form a finite sum of polynomials times exponentials?);
- M3: a Gröbner basis of the holonomic annihilator, checking for finite holonomic rank in the sense of Lemma 2.5 (ledger OP-ALG-T1);
- M2: undecidable from the jet alone (ledger P4-T5); one needs an upper bound $B \geq \dim(\mathcal{R}u \cap \mathrm{EP})$, after which comparing the first B derivatives suffices (ledger P4-T4).




Figure 9: The algorithm as a flowchart. Left column: steps 0–3 certify what the operator is. If every factor test fails (no leftover is an exponential-polynomial) the operator is identified; otherwise step 4 reduces it to G_u. Right column: steps 4–7 extract structure from u once the operator is known; the dashed arrow shows that an identified operator proceeds directly to decomposition.

Step 4: reduce to G_u

Does drop from P every factor P_i whose test returned "EP" (with the correct multiplicity bookkeeping: at most $\sum_i e_i$ tests in total (ledger C3-T6)).

Returns G_u, the intrinsic operator; the dropped part is J_u, whose finite zero set is the point spectrum (the exponentials contained in u). Certificate: "$\mathrm{Ann}(u) = (P)$" if nothing was dropped, else "u identifies only G_u and contains these exponentials".

Step 5: decompose

Evolution factors $\tau - a_j(\xi)$: apply (2).
General coprime factors use the exact-sequence construction of Theorem 5.2; the components are determined up to the finite exponential-polynomial space of dimension $\delta(P)$, fixed by a normalization (for the wave equation, by fixing the constant, Example 5.2).
Check before using the components in H^m, verify $\gamma_N > 0$ (Theorem 5.5).

Step 6: terms and interaction

Does inside an evolution factor with symbol $a(\xi) = \sum_n a_n(\xi)$, read the terms as the coefficients of $\log \hat{u}(t,\kappa) = t\sum_n a_n(\kappa) + \log \hat{u}_0(\kappa)$ (linear in the terms, cost $O(N)$); evaluate E_{int} by (4).
Non-commuting classify the generated Lie algebra: solvable $\Rightarrow$ exact formula; otherwise a local expansion only.

Step 7: nonlinear case

Replace Step 3 by the dictionary criterion of Theorem 7.2 and Step 6 by the flow-based E_{int} of Section 7.

8.4 Worked example

We follow Steps 1–6 on the wave equation with $c = 1$ and the (non-exponential) solution

$$u(t,x) = \underbrace{\frac{1}{1+(x+t)^2}}_{f(x+t)} + \underbrace{e^{-(x-t)^2}}_{g(x-t)}.$$

Each box below states the goal, the computation, and the outcome of one step.

Step 1 — annihilator

Goal. Find an equation satisfied by u.
Computation. Take the dictionary $\{u_{tt}, u_{xx}, u_{tx}, u_t, u_x, u\}$; the relations $\lambda \cdot \Phi = 0$ form the line spanned by $u_{tt} - u_{xx} = 0$.
Outcome. $P = \tau^2 - \xi^2$.

Step 2 — factor

Computation.

$$P = Q_1 Q_2, \qquad Q_1 = \tau - \xi, \quad Q_2 = \tau + \xi.$$

Outcome. Two irreducible, simple factors.

Step 3 — EP test, factor Q_2 removed

Goal. Decide whether $w_1 = Q_2(\partial)u = (\partial_t + \partial_x)u$ is an exponential-polynomial.
Computation. The two parts behave differently:

$$(\partial_t + \partial_x) f(x+t) = 2f'(x+t), \qquad (\partial_t + \partial_x) g(x-t) = -g' + g' = 0.$$

With $f(s) = \dfrac{1}{1+s^2}$ one has $f'(s) = -\dfrac{2s}{(1+s^2)^2}$, hence

$$w_1 = 2f'(x+t) = -\frac{4(x+t)}{\big(1+(x+t)^2\big)^2}.$$

Outcome. w_1 is rational, not a finite sum of polynomials times exponentials, so $w_1 \notin$ EP.

Step 3 — EP test, factor Q_1 removed

Computation. Symmetrically, with $g(s) = e^{-s^2}$, $g'(s) = -2se^{-s^2}$:

$$w_2 = Q_1(\partial)u = (\partial_t - \partial_x)u = -2g'(x-t) = 4(x-t)\, e^{-(x-t)^2}.$$

Outcome. $w_2 \notin$ EP (a Gaussian times a polynomial is not an exponential-polynomial).

Step 4 — reduce

Outcome. Both tests say $\notin$ EP, so nothing is dropped: $\mathrm{Ann}(u) = (\tau^2 - \xi^2)$, the point spectrum is empty, and the wave equation is identified.

Step 5 — decompose

Computation. $(Q_1, Q_2) = (\tau, \xi)$, so $\delta(P) = 1$.
Outcome. The components are $u_1 = f(x+t)$, recovered from w_1 by integration in x at fixed t up to a constant, and $u_2 = g(x-t)$; they are determined up to a constant k with $u_1 + u_2$ fixed. Whether such components lie in H^m for every solution is governed by Theorem 5.5 and is not asserted here.

Step 6 — terms

Not applicable to the wave equation (no evolution factor with several terms). For the advection–diffusion case of Figure 6, the log-symbol is

$$\log \hat{u}(t,\kappa) = \log \hat{u}_0(\kappa) + t\,(-ic\kappa - D\kappa^2),$$

its two coefficients are the two terms, and E_{int} follows from (4) at cost $O(N) = O(2)$ per point.

8.5 Complexity summary

Table 7 collects the cost and feasibility of each component.

Component	Cost / feasibility	Source
Annihilator from a dictionary	Gaussian elimination, $O(M^3)$	Step 1
Factorization	standard computer-algebra operation	Step 2
EP test (M1)	syntactic	Step 3
EP test (M3)	Gröbner/holonomic, decidable mod (AC)	(ledger OP-ALG-T1)
EP test (M2, black box)	impossible without a bound	(ledger P4-T5)
Components (evolution)	one rational Fourier multiplier per factor	(2)
E_{int} value / by order / full table	$O(N)$ / $O(N^2)$ / $\Theta(2^N)$	(ledger P1-T3)

Table 7: Cost and feasibility of the algorithm's components.

9 Limits, open problems, and status

In this section

Starting from	no new assumptions.
Why this section	to state what is impossible, what is open, and how reliable the results are.
Role in the paper	delimits the claims of the paper; Appendix C compares the paper with the full research.

What is impossible (all [proved]).

- finite certificates of identification, and black-box identification (Section 4.3);
- per-term decomposition of the solution (Proposition 5.1[E9]);
- stable regularization of E_{int} (Proposition 6.3);
- identification by restricting variable-coefficient operators to one solution;
- decomposition on domains with holes (Figure 4);
- a finite product of single-term flows reproducing non-commuting dynamics.

Open or conditional.

- identification for analytic KdV data with nonzero reflection coefficient (ledger INT-Q1) [open];
- time-uniform shock estimate with several shock births (ledger OPA-Q1) [conditional];
- density of the sum of kernels on nonconvex domains [open];
- exact values $\delta_N(K)$ of an invariant for general N (ledger OPF5C-Q1) [open];
- nonlinear asymptotics with derivatives (ledger S4ID-P8) [conditional];
- closedness of the sum of kernels for $N \geq 3$ differential factors (ledger C12-Q2) [open].

Honest assessment. Many ingredients are classical (Malgrange–Ehrenpreis, Nullstellensatz, Cole–Hopf, inverse scattering). The contribution is the *package*: sharp characterizations, negative theorems, and counterexamples, each with a verified status. Independent reviewers were AI agents. Fifteen theorems of the algebraic core (the E_{int} identity for commuting flows, the comaximal factor decomposition with explicit projector, companion normality for $k = 2$, irreducibility results) are formalized in Lean 4/Mathlib without `sorry`; the analytic parts (distributions, Sobolev spaces, semigroups) are not, since Mathlib lacks the libraries. Several citations were verified only on secondary sources; a result relied upon should be traced to its primary source. Out of scope so far: a variable potential treated as noise (Schrödinger/Dirac), matrix-valued coefficients (4×4 Dirac), and any statistical preprocessing.

A Glossary of tools

Each entry states a classical tool in one sentence and points to the section where the paper uses it.

Name	Statement in one sentence	Used in
Chinese remainder theorem	Coprime moduli split a quotient ring into a product; the algebra behind Thm. 5.2.	Sec. 5
Hilbert's Nullstellensatz	A polynomial vanishing on $V(I)$ has a power in I.	Sec. 2, 4
Bézout's theorem	Curves without common component meet in finitely many points (at most the product of degrees) [6].	Sec. 2
Malgrange–Ehrenpreis	Every nonzero constant-coefficient operator has a fundamental solution; $P(\partial)$ is onto on C^∞ of a convex set [2, 3].	Sec. 5
Ehrenpreis–Palamodov	Solutions of constant-coefficient systems are integrals of exponential-polynomial solutions over $V(P)$ (discrete sums when the spectrum is finite) [4].	context only
Friedrichs angle	$M + N$ is closed iff the cosine of the angle between $M \ominus (M \cap N)$ and $N \ominus (M \cap N)$ is < 1.	Sec. 5
Miyadera–Voigt	Small relatively bounded perturbations of a semigroup generator generate a semigroup; basis of (5) [7].	Sec. 6
Baker–Campbell–Hausdorff	$e^X e^Y = e^{X+Y+\frac{1}{2}[X,Y]+\cdots}$.	Sec. 6
Wei–Norman	Linear time-dependent systems on a solvable Lie algebra reduce to ODEs for scalar coefficients.	Sec. 6
Cole–Hopf	$u = -2\nu\varphi_x/\varphi$ maps Burgers to the heat equation [9].	Sec. 7
Kružkov	Uniqueness and stability of entropy solutions of scalar conservation laws; defines the flow of $-uu_x$ after the shock [10].	Sec. 7
Inverse scattering (IST)	KdV/NLS are linearized by a nonlinear Fourier transform with trivial time evolution.	Sec. 7

B Primer for readers new to the concepts

This appendix *adds* concrete explanations; nothing in the main text depends on reading it. It covers four notions in this order: ideals and annihilators (§B.1), exponential-polynomials (§B.2), the Chinese remainder theorem (§B.3), and the companion matrix on a numerical example (§B.4).

B.1 Ideals, annihilators and the gcd, starting from integers

Integers. The multiples of 6 form an *ideal* $(6) = \{\dots, -6, 0, 6, 12, \dots\}$: it is closed under addition and under multiplication by any integer. The ideal generated by 4 and 6 is $(4, 6) = (2)$, since $2 = 6 - 4$ and every combination is even: the ideal "remembers" only the gcd.

Operators. Replace integers by operators and "multiple of 6" by "kills u". For $u = e^x$ every operator $Q(\partial_x)$ with $Q(1) = 0$ kills u, i.e. every multiple of $\partial_x - 1$. So

$$\begin{aligned}\operatorname{Ann}(e^x) &= (\partial_x - 1),\\ \operatorname{Ann}(\sin x) &= (\partial_x^2 + 1),\\ \operatorname{Ann}(xe^x) &= \big((\partial_x - 1)^2\big),\\ \operatorname{Ann}(e^x + e^{2x}) &= \big((\partial_x - 1)(\partial_x - 2)\big).\end{aligned}$$

In one variable every annihilator ideal has a single generator (the *minimal annihilator*), exactly as every ideal of integers is (n). In two variables an ideal can need several generators, as $(\tau - \tau_0, \xi - \xi_0)$ does for $u = e^{\tau_0 t + \xi_0 x}$.

The gcd. $\operatorname{Ann}(u) = G_u J_u$ separates the part of the ideal that is a single polynomial multiple (G_u, "infinite part") from the part J_u that has only finitely many common zeros (J_u, "finite debris"). Example: for $u = f(x + t)$ with f generic (not an exponential-polynomial), $\operatorname{Ann}(u) = (\partial_t - \partial_x)$ by Theorem 4.1, so $G_u = \tau - \xi$ and $J_u = \mathcal{R}$; for $u = e^{x+t}$, $\operatorname{Ann}(u) = (\tau - 1, \xi - 1)$, so $G_u = 1$ and $J_u = \operatorname{Ann}(u)$.

B.2 Exponential-polynomials: examples and a test

Function	In EP?	Reason (derivatives of u span ...)
xe^x	yes	$\operatorname{span}\{xe^x, e^x\}$, dimension 2
$x^2e^{3t} + \cos x$	yes	finite sum of polynomial $\times$ exponential ($\cos x = \frac{1}{2}(e^{ix} + e^{-ix})$)
e^{-x^2} (Gaussian)	no	$\frac{d^k}{dx^k}e^{-x^2} = (\text{polynomial of degree } k)\, e^{-x^2}$: independent for all k, infinite dimension
$1/(1 + x^2)$	no	derivatives are rational functions with growing denominators: infinite dimension

This is Lemma 2.5 in action: "$u \in$ EP" means "its derivatives span a finite-dimensional space".

B.3 The Chinese remainder theorem, concretely

Integers. The system $x \equiv 1 \pmod 2$, $x \equiv 2 \pmod 3$ has the unique solution $x \equiv 5 \pmod 6$: $\mathbb{Z}/6 \cong \mathbb{Z}/2 \times \mathbb{Z}/3$. Coprimality of 2 and 3 is what makes the pair of remainders determine x.

One-variable ODE. Let $p(\partial) = (\partial - 1)(\partial - 2)$. Then $\ker p = \ker(\partial - 1) \oplus \ker(\partial - 2) = \operatorname{span}\{e^x, e^{2x}\}$, and the Lagrange-type projectors are explicit: if $u = ae^x + be^{2x}$,

$$\begin{aligned}u_1 &= \frac{\partial - 2}{1 - 2}\, u = \;\; a\, e^x,\\ u_2 &= \frac{\partial - 1}{2 - 1}\, u = \;\; b\, e^{2x}.\end{aligned}$$

This is Eq. (2) with ∂_t replaced by ∂.

Two variables. For the wave equation $Q_1 = \tau - \xi$, $Q_2 = \tau + \xi$ are coprime, but the ideal $(Q_1, Q_2) = (\tau, \xi)$ is not the unit ideal: the two lines meet at the origin. The quotient $\mathcal{R}/(\tau, \xi) = \mathbb{C}$ has dimension 1, which is the constant ambiguity $\delta(P) = 1$ of Example 5.2. Theorem 5.2 says that this finite overlap is the *only* failure of the integer picture.

B.4 The companion matrix on a numerical example

For $u'' - 3u' + 2u = 0$ the companion matrix is $C = \begin{pmatrix} 0 & 1 \\ -2 & 3 \end{pmatrix}$, with characteristic polynomial $\lambda^2 - 3\lambda + 2$ and eigenvalues $1, 2$. The projectors of Eq. (1) are $\Pi_1 = \dfrac{C - 2\mathrm{Id}}{1 - 2}$ and $\Pi_2 = \dfrac{C - \mathrm{Id}}{2 - 1}$, and $e^{xC} = e^x \Pi_1 + e^{2x} \Pi_2$.

C Scope: what this paper contains and what it does not

This paper summarizes the research reported in `REPORT_FINALE.md` (Phases 0–5). The following table compares it with the full project.

Area	In this paper	Not in this paper
Linear identification	criterion, spectral form, limits (Sec. 4)	proofs of the converse direction; details of OP-ALG and variable-coefficient classification
Factor decomposition	Thm. 5.2, holes, Sobolev regularity (Sec. 5)	the explicit $\delta_N(K)$ values (open), the cohomological criterion in detail
Terms and E_{int}	product formula, cost, instability, Dyson, Lie dichotomy (Sec. 6)	boundary asymptotics beyond the stated rate; the balance decomposition
Nonlinear	Burgers asymptotics, dictionary criterion (Sec. 7)	KdV/NLS scattering identification (INT-T1/T2), Hirota N-solitons, analyticity radius, KdV–Burgers remainder
Catalogue of kernels	–	the catalogue of $\sim$60 equation types (C2), the tropical scaling reading
Formalization	Lean statement (Sec. 9)	the Lean sources
Phases 6–11	–	systems and matrix coefficients, variable coefficients and the analytic encoder, convergence, universal dictionary, expanded algorithm: planned and scouted; the proof packages of Phase 8 had not been delivered when this paper was written

The paper therefore reports the results of Phases 0–5 at the level of statements, examples and proof ideas, not the full set of roughly 190 ledger entries, and it must not be read as a report on the extended algorithm of Phases 6–11.

D Notes for the author

These notes collect the points flagged during the editorial revision; each is marked in the text by a red superscript.

E1. the symbols M, d and L in this proposition come from the ledger entry P1-T3 and are not defined in the paper; the author should supply their definitions (presumably M evaluation points, d and L degree and order parameters of the class $\mathfrak{B}$).

E2. Fisher–KPP is announced in the abstract and as a benchmark, but no statement about it appears in the body (the ledger has S4T-P5, an estimate that is not uniform in t). The author may add a one-line statement or drop it from the abstract. A short illustrative example for the exact sequence of the dictionary criterion, derived from the Burgers case, would help the reader; it is not added here because it cannot be derived safely from the existing text.

E3. the research ledger assumes a candidate annihilator P is given. The Gram-matrix procedure above describes one way to obtain it from a dictionary, based on the notation file (the jet Φ and Gram matrix $\mathbb{G}(u)$); the author should confirm that this is the intended Step 1.

E4. The "Ehrenpreis–Palamodov" glossary entry is context only (not used in the body), and Stein is cited in Theorem 5.5 without a bibliography entry.

E5. Overloaded symbols (S, D, J, σ, c) are documented in the caption of Table 2 but not renamed, because renaming would touch many formulas.

E6. Only the easy direction of Theorem 4.1(a) is proved in the text; the converse is deferred to the project's proof package.

E7. The Gaussian example of Section 6.2 and the Lagrange projector formula (2) were derived by the editor, consistent with the ledger (P1-T2, II-T2) but not literally contained in it; please verify.

E8. In Table 3 the parameter d of the threshold $2d-1$ is not defined (it comes from the ledger entry C3-T3); please state what d is.

E9. Proposition 5.1 as originally stated (inclusion in $\ker(L_1, L_2)$) is false: $u = x^2/2 + Dt$ is a counterexample for the heat equation. The corrected statement (inclusion in $\ker(L, L_1L_2)$) and its short proof are the editor's and have *not* been independently reviewed, so the "proved" badge was removed; please re-review.

E10. In Section 6.5, the paper does not say which pair (A_1, B) gives the boundary-induced interaction, in which norm the estimate $\|E_{\mathrm{int}}\| \sim t^{k/2+3/4}$ holds, how the order of contact k is defined, or where the exponent $3/4$ comes from.

E11. The isomorphism $\mathcal{R}u \cap \mathrm{EP} \cong \mathcal{R}/J_u$ in Proposition 4.3 is stated without proof in the text; only the equivalence is derived. Please supply the argument or the ledger reference.

E12. The original sentence about $P = \tau$, $u = e^{x^2}$ (Section 4.2) did not match its claim (τ is irreducible, and $V(\mathrm{Ann}\, u)$ is a line, not a point spectrum); it was replaced by the verified example $P = \tau^2$, $u = e^{x^2}$. Please confirm this is what was meant.

E13. In Section 7.3 the map ψ and the terms "evolutionary dictionary" and "complete dictionary" are not defined. The jet-level obstruction was renamed $\mathcal{J}_u$ to avoid the clash with the cofactor J_u.

References

[1] D. Cox, J. Little, D. O'Shea, *Ideals, Varieties, and Algorithms*, 4th ed., Springer, 2015.

[2] L. Ehrenpreis, Solution of some problems of division I, *Amer. J. Math.* 76 (1954).

[3] B. Malgrange, Existence et approximation des solutions des équations aux dérivées partielles et des équations de convolution, *Ann. Inst. Fourier* 6 (1955–56).

[4] V. P. Palamodov, *Linear Differential Operators with Constant Coefficients*, Springer, 1970.

[5] L. Hörmander, *The Analysis of Linear Partial Differential Operators II*, Springer, 1983.

[6] W. Fulton, *Algebraic Curves*, Benjamin, 1969.

[7] K.-J. Engel, R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Springer, 2000.

[8] F. J. Dyson, The radiation theories of Tomonaga, Schwinger and Feynman, *Phys. Rev.* 75 (1949).

[9] E. Hopf, The partial differential equation $u_t + uu_x = \mu u_{xx}$, *Comm. Pure Appl. Math.* 3 (1950); J. D. Cole, *Quart. Appl. Math.* 9 (1951).

[10] S. N. Kružkov, First order quasilinear equations in several independent variables, *Math. USSR Sb.* 10 (1970).

[11] S. P. Novikov, The periodic problem for the Korteweg–de Vries equation, *Funct. Anal. Appl.* 8 (1974).

[12] Project reports: `REPORT_FINALE.md`, `claims/LEDGER.md`, `PROBLEM.md`, `work/*/PROOF_PACKAGE.md`, `BIBLIOGRAFIA.md` (112 sources with verification state).