

第二章 函数的导数(Derivative)

2.1 导数的定义(The Definition of Derivative)



Multiple Choice Questions


1. Which of the following values is the average rate of change of $f(x) = \sqrt{x+2}$ over the interval $[-1, 2]$?

- (A) -4 (B) $-\frac{1}{3}$ (C) $\frac{1}{3}$ (D) 3

2. Let $f(x) = 4 - 3x$. Which of the following is equal to $f'(0)$?

- (A) 4 (B) -4 (C) -3 (D) 4

3. The graph of f and its tangent line at $x=1$ is shown below, find $f'(1) = (\quad)$.



- (A) 0 (B) $\frac{1}{2}$ (C) 1 (D) 2

4. Let $f(x) = kx^3 - 2x$, ($k \neq 0$). The average rate of $f(x)$ over the interval $[-2, -1]$ is 3, find k .

- (A) $\frac{5}{7}$ (B) $\frac{1}{7}$ (C) $-\frac{1}{7}$ (D) $-\frac{5}{7}$

5. **Calculator** Let $y = \tan e^x - x^2 + 1$. Find $\frac{dy}{dx} \big|_{x=0.5}$.

- (A) -12.056 (B) 0 (C) 271.066 (D) 273.066

6. **Calculator** The graph of $y = \sqrt{x^2+1} - 3x$ crosses the x -axis at one point in the interval $[0, 1]$. What is the slope of the graph at this point?

- (A) 0.354 (B) -0.001 (C) -2.666 (D) 0

7. The population of insects at time t days is modeled by the function P with first derivative $P'(t) = 0.1t^2 + 10t + 120$. Which of the following statements is true?

- (A) At time $t = 10$, the population of insects is 230.
 (B) At time $t = 10$, the population of insects is increasing at a rate of 230 insects per day.
 (C) At time $t = 10$, the population of insects is decreasing at a rate of 230 insects per day.
 (D) At time $t = 10$, the rate of change of the number of insects is increasing at a rate of 230 insects per day.

8. If the line tangent to the graph of a function g at the point $(1, 4)$ passes through the point $(-2, -5)$, what is the slope of this line?

- (A) 3 (B) -3 (C) $\frac{1}{3}$ (D) $-\frac{1}{3}$



Free Response Questions

9. Let $f(x) = x^2 - 2x$

(a) Find the average rate of change of f over $[1, 3]$.

(b) Find $\frac{f(2+h) - f(2)}{h}$.

(c) Find the instantaneous rate of change of f at $x = 2$.

9. a. $\frac{9-6-(1-2)}{3-1} = 2$ c. $\lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h}$

b. $\frac{(2+h)^2 - 2(2+h) - (4-4)}{h} = \lim_{h \rightarrow 0} \frac{h^2 + 4h + 4 - 4 - 4h}{h} = \lim_{h \rightarrow 0} \frac{h^2}{h} = 2$

$= h + 2$

10. Use the definition of derivative to find the derivative of $f(x) = x^3$, then find $f'(2)$.

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^3 + 3xh^2 + 3x^2h + x^3 - x^3}{h} \\ &= \lim_{h \rightarrow 0} (h^2 + 3xh + 3x^2) \\ &= 3x^2 \\ f'(2) &= 3 \times 2^2 = 12. \end{aligned}$$