

Theoretical Bounds on χ_ν in Observer-Patch Holography

Canonical Coherent-Matter Susceptibility and Repair-Charge Coupling

Bernhard Mueller, Alexander Osika, and David Matscheko

August 13, 2026

Paper release: r2019 **Released:** August 13, 2026

Author affiliations: Bernhard Mueller, Pragma Research Inc.; Alexander Osika, Asimov Labs.

Abstract

The coherent-material branch of Observer-Patch Holography (OPH) supplies a canonical scalar source in the same quotient-edge register as the protected collar reserve. On the co-registered presence branch, the dimensionless coupling is $\chi_\nu^{\text{can}} = 1 - P/24$. At the CODATA-derived comparison pixel this is $0.9320429912748350\dots$. Under the weaker mean-count reading it lies between that value and one. This is a channel coefficient, not a force coefficient. A proposed repair-charge condensate action couples the source to a dynamical repair phase. Conditional on that action, repair-current continuity, dilute dust behavior, spherical deep-galaxy scaling, field stress, and a lawful coherent-source force follow from one variational system. A compact neutral coherence contrast has no monopole and cannot support a closed device: no formula proportional only to $g^2 A \chi_\nu \Delta S / (4\pi G)$ is a force theorem for that device class. This paper supplies no dimensional source calibration, ambient repair field, compact-phase completion, or laboratory source receipt; no laboratory force prediction and no numerical lift prediction follow from χ_ν .

Claim Boundary

This paper proves a finite-channel coefficient on a declared OPH branch. It also derives continuity, dilute dust behavior, spherical deep-galaxy scaling, force, and momentum formulae conditional on the repair-charge condensate action in the dark-sector paper [4]. The stated axioms and theorems supply neither that action nor a dimensional source normalization; both enter as premises, and every force statement below is conditional on them.

No laboratory force prediction follows from this paper. The formula $g^2 A \chi_\nu \Delta S / (4\pi G)$ is not a force theorem for a closed compact device, and a static, closed, repair-neutral device cannot lift itself.

The exact scope is:

Object	Scope
Coherent source in the quotient-edge scalar register	finite theorem under the stated source receipts
$\chi_\nu^{\text{can}} = \lambda_{\text{collar}}$	theorem under co-registration and reserve-survival premises
$\chi_\nu^{\text{can}} = 1 - P/24$	exact on the presence branch
Repair-charge action	proposed dynamical completion
Continuity, dust limit, and spherical deep-galaxy scaling	derived conditional on that action and its constitutive premises
Dimensional coupling $q_\star$	undetermined; a named premise of every force statement
Device force	conditional on source charge, external field, and momentum closure
Closed compact neutral device	no force theorem; the formula $g^2 A \chi_\nu \Delta S / (4\pi G)$ is excluded for this class
Laboratory force prediction and numerical lift	none

Contents

1 Canonical Coherent-Matter Source	2
2 Protected-Reserve Coefficient	3
3 Engineering Coordinates	4
4 Repair-Charge Coupling	4
5 Compact-Source Constraint	6
6 Momentum and Energy Closure	6
7 Laboratory Evidence Conditions	7
8 Conclusion	8

1 Canonical Coherent-Matter Source

An OPH technology instantiation is an observer-like self-reading system: a bounded physical or software patch with local state, ports or boundaries, readback, records, feedback or repair moves, and a public evidence bundle. For a nondegenerate coherent material subfederation U , define the

canonical source

$$S_{\text{coh}}^{\text{can}} = \mathbf{1}_{\text{self-read}} \mathbf{R}_U \mathbf{P}_U \mathbf{C}_U. \quad (1.1)$$

The factors state that the patch reads itself, produces a stable record, supports a predictive boundary, and executes a registered repair or control move. If any factor vanishes, the coherent scalar source is zero.

The screen-microphysics branch supplies four structural facts [3]:

- S1. $S_{\text{coh}}^{\text{can}}$ descends to a protected quotient observable.
- S2. A nondegenerate source is nonzero.
- S3. Scalar edge-center exhaustion places it in the unique quotient-local scalar register.
- S4. The protected $\mathbb{Z}_6$ reserve acts on that same register and commutes with scalar activation.

These statements identify the channel. They supply no spacetime action or dimensional gravitational charge.

2 Protected-Reserve Coefficient

Let $\epsilon_{\mathbb{Z}_6}(y)$ be the conditional reserve-presence profile on the co-registered scalar slot, and let $w(y)$ be the normalized scalar weight across the finite collar. The scalar opportunity survives with local factor

$$\lambda_{\text{slot}}(y) = 1 - \epsilon_{\mathbb{Z}_6}(y). \quad (2.1)$$

The finite-thickness collar coefficient is

$$\lambda_{\text{collar}} = \int dy w(y) [1 - \epsilon_{\mathbb{Z}_6}(y)]. \quad (2.2)$$

The protected-reserve receipt gives

$$\int dy w(y) \epsilon_{\mathbb{Z}_6}(y) = \frac{P}{24}. \quad (2.3)$$

The symbolic result below holds for the supplied pixel coordinate. Its decimal evaluation uses the measured-endpoint comparison input $P_C = 1.630968209403959\dots$, obtained from the CO-DATA/NIST fine-structure endpoint. It is not a source-selected OPH value.

Theorem 2.1 (Canonical susceptibility). *Assume the nonzero coherent-source branch, scalar edge-center exhaustion, co-registration with the protected reserve, and the presence reading in Eqs. (2.2)–(2.3). Then*

$$\boxed{\chi_{\nu}^{\text{can}} = \lambda_{\text{collar}} = 1 - \frac{P}{24}}. \quad (2.4)$$

For the comparison input $P = P_C = 1.630968209403959\dots$,

$$\boxed{\chi_{\nu}^{\text{can}} = 0.9320429912748350\dots}. \quad (2.5)$$

Proof. The coherent source enters the scalar opportunity count before the protected reserve acts. Co-registration makes the local surviving fraction $1 - \epsilon_{\mathbb{Z}_6}(y)$. Averaging with $w(y)$ gives $\lambda_{\text{collar}} S_{\text{coh}}^{\text{can}}$. The definition of canonical susceptibility gives $\chi_{\nu}^{\text{can}} S_{\text{coh}}^{\text{can}}$. Nondegeneracy permits cancellation of $S_{\text{coh}}^{\text{can}}$, and Eq. (2.3) gives the value. $\square$

Corollary 2.2 (Mean-count band). *If $\epsilon_{\mathbb{Z}_6}(y)$ is interpreted only as the mean of a nonnegative integer occupancy, then at the same comparison input*

$$\boxed{0.9320429912748350 \dots \leq \chi_{\nu}^{\text{can}} \leq 1}. \quad (2.6)$$

Proof. Markov's inequality gives $1 - \epsilon \leq \Pr[N = 0] \leq 1$. Integration and Eq. (2.3) give the band. $\square$

Remark 2.3 (Depth limit). The iterated fine-grained survival family $(1 - (P/24) 2^{-m})^{2^m}$ tends to $e^{-P/24}$ as the refinement depth m grows. That exponential is the depth limit, unreachable at any finite regulator; the finite one-step survival is the presence value $1 - P/24$, and the presence branch of the theorem uses it.

The theorem excludes a zero scalar-channel coefficient on the stated nondegenerate branch. It does not exclude a zero dimensional gravitational coupling $q_{\star}$, a zero device repair charge, or a zero ambient repair gradient.

3 Engineering Coordinates

An engineering description may absorb an effective coherent capacity into the source coordinate:

$$S_{\text{coh}}^{\text{eng}} = N_{\text{coh}} S_{\text{coh}}^{\text{can}}. \quad (3.1)$$

Invariance of the scalar perturbation requires

$$\chi_{\nu}^{\text{can}} S_{\text{coh}}^{\text{can}} = \chi_{\nu}^{\text{eng}} S_{\text{coh}}^{\text{eng}}, \quad \boxed{\chi_{\nu}^{\text{eng}} = \frac{\chi_{\nu}^{\text{can}}}{N_{\text{coh}}}}. \quad (3.2)$$

Very small values of χ_{ν}^{eng} can therefore represent an order-one canonical coefficient. Equation (3.2) is a chart identity. It does not define $q_{\star}$, measure $S_{\text{coh}}^{\text{can}}$, or predict force.

4 Repair-Charge Coupling

The dark-sector completion takes integer scalar repair occupation and a compact repair phase to a canonical pair (n, θ) . On a fixed real lift of the phase, its relevant action is

$$\begin{aligned} I_{\text{RPC}} = I_{\text{Regge}} + I_{\text{SM}} \\ + \int d\tau \left[- \sum_v V_v n_v D_{\tau} \theta_v - \sum_v V_v \varepsilon(n_v) - \sum_e W_e \mathcal{K}(|(d\theta)_e|) \right. \\ \left. - \sum_v V_v \theta_v \left(\mathcal{Q}_{b,v} + q_{\star} \chi_{\nu}^{\text{can}} S_{\text{coh},v}^{\text{can}} \right) \right]. \end{aligned} \quad (4.1)$$

The source and canonical signs are chosen so phase variation gives the positive-source balance in Eq. (4.3), and translation of a positive localized source gives force $-Q_R \nabla \theta$. The action is a posit: it comes with no globally single-valued compact-phase coupling, no derivation from finite repair dynamics, and no dimensional constants. Each of these three objects is a premise of the conditional consequences below.

The coherent repair-charge density is

$$\mathcal{Q}_{\text{coh}} = q_* \chi_\nu^{\text{can}} S_{\text{coh}}^{\text{can}}. \quad (4.2)$$

The phase equation is

$$D_\tau n + d^\dagger j_R = \mathcal{Q}_b + q_* \chi_\nu^{\text{can}} S_{\text{coh}}^{\text{can}}. \quad (4.3)$$

For a finite region Ω ,

$$\frac{d}{d\tau} \sum_{v \in \Omega} V_v n_v + \sum_{e \in \partial \Omega} W_e j_{R,e} = \sum_{v \in \Omega} V_v \left(\mathcal{Q}_{b,v} + q_* \chi_\nu^{\text{can}} S_{\text{coh},v}^{\text{can}} \right). \quad (4.4)$$

The boundary flux is part of the physical source ledger.

Proposition 4.1 (Conditional rotor consequences). *Assume Eq. (4.1), a positive branch $n > 0$, and the infrared constitutive functions*

$$\varepsilon(n) = m_R n + \frac{u_R}{3} n^3, \quad \mathcal{K}(y) = \frac{\kappa_R}{3} y^3, \quad m_R, u_R, \kappa_R > 0. \quad (4.5)$$

Then:

(i) *phase variation gives the exact finite repair-current balance Eq. (4.4);*

(ii) *the homogeneous dilute phase has*

$$\rho_R = m_R n + \frac{u_R}{3} n^3, \quad p_R = \frac{2u_R}{3} n^3, \quad (4.6)$$

so source-free expansion gives $n \propto a^{-3}$, $\rho_R \propto a^{-3}$, and $w_R \simeq 0$ when $u_R n^2 \ll m_R$;

(iii) *if $\mathcal{Q}_b = \beta_b \rho_b$ and $\mathbf{a}_R = -\beta_b \nabla \theta$, the static spherical exterior branch gives*

$$a_R = \sqrt{a_b a_0}, \quad v^4 = G M_b a_0, \quad a_0 = \frac{\beta_b^3}{4\pi G \kappa_R}; \quad (4.7)$$

(iv) *translation of the coherent source gives Eq. (4.9) and the momentum ledger Eq. (6.2).*

These statements are consequences of the proposed action, not derivations of that action from the finite OPH theorems.

Proof. Item (i) follows by varying θ and summing the local equation over a finite region. For item (ii), the first-order rotor Hamiltonian density is $\varepsilon(n)$, and its barotropic pressure is $p_R = n\varepsilon'(n) - \varepsilon(n)$. Source-free homogeneous continuity then gives $na^3 = \text{constant}$. For item (iii), spherical integration of $\nabla \cdot (\kappa_R |\nabla \theta| \nabla \theta) = \beta_b \rho_b$ gives $4\pi r^2 \kappa_R |\theta'| \theta' = \beta_b M_b$; the stated acceleration and definition of a_0 give the result. Item (iv) follows by translating the source term in Eq. (4.1) and applying translation invariance to matter plus repair field. $\square$

Define the integrated device repair charge

$$Q_R^{\text{dev}} = q_\star \chi_\nu^{\text{can}} \int_\Omega S_{\text{coh}}^{\text{can}}(x) \, \text{d}^3x. \quad (4.8)$$

For a prescribed external repair phase, translation of the source gives

$$F_i^{\text{coh}} = -q_\star \chi_\nu^{\text{can}} \int_\Omega S_{\text{coh}}^{\text{can}}(x) \, \partial_i \theta_{\text{ext}}(x) \, \text{d}^3x. \quad (4.9)$$

If the external gradient varies slowly across the device,

$$F_i^{\text{dev}} \simeq -Q_R^{\text{dev}} \partial_i \theta_{\text{ext}}. \quad (4.10)$$

The sign of Q_R^{dev} may select attractive or repulsive response. Positive repair energy is compatible with signed charge when $\varepsilon(n) = \varepsilon(|n|)$.

Remark 4.2 (Switchable-sign boundary). The gated source in Eq. (1.1) is nonnegative under its displayed definition. With fixed $q_\star$ and positive χ_ν^{can} , exchanging top and bottom source profiles does not reverse the monopole Q_R^{dev} . A switchable force sign therefore requires either a signed orientation factor derived as part of the coherent source map or a controlled reversal of $\nabla \theta_{\text{ext}}$. The finite channel theorem supplies neither operation.

5 Compact-Source Constraint

Proposition 5.1 (No monopole from a neutral compact source). *Suppose the coherent repair source is compactly supported and satisfies*

$$\int_\Omega \mathcal{Q}_{\text{coh}} \, \text{d}^3x = 0. \quad (5.1)$$

Then it has no repair monopole. Its exterior field begins at dipole order or higher. In a uniform external gradient, the leading forces on its positive and negative charges cancel.

Proof. Integrating Eq. (4.3) in a static state makes the exterior flux equal the integrated source. Equation (5.1) sets the monopole flux to zero. The multipole expansion therefore begins beyond the monopole. A uniform external gradient couples only to total charge at leading order, which vanishes. $\square$

A bottom-to-top contrast

$$\Delta S = S_{\text{bottom}} - S_{\text{top}} \quad (5.2)$$

can shape a dipole, select the sign of injected repair charge, couple to a nonuniform external field, or drive an alternating repair current. It does not by itself define a nonzero Q_R^{dev} . No formula proportional only to $g^2 A \chi_\nu \Delta S / (4\pi G)$ is a force theorem for a closed compact device.

6 Momentum and Energy Closure

For the cubic condensed branch, the repair-field stress is

$$T_{ij}^R = \kappa_R |\nabla \theta| \partial_i \theta \partial_j \theta - \delta_{ij} \frac{\kappa_R}{3} |\nabla \theta|^3 + \dots. \quad (6.1)$$

The exact force ledger is

$$F_i^{\text{matter}} + \frac{dP_i^R}{dt} + \oint_{\partial\Omega} T_{ij}^R n^j dA = 0. \quad (6.2)$$

A local repulsive response requires either nonzero device repair charge with a field tail or explicit momentum transfer through Eq. (6.2). A static, closed, repair-neutral device cannot lift itself.

Any switchable force also requires a toggle-energy ledger. Let $E(q, s)$ be the energy at position q and internal state s , and define

$$\text{Toggle}(q) = E(q, \text{on}) - E(q, \text{off}). \quad (6.3)$$

For an on/off spatial cycle between q_1 and q_2 , conservation gives

$$W_{\text{cycle}} = \text{Toggle}(q_1) - \text{Toggle}(q_2). \quad (6.4)$$

A claimed position-dependent switchable force with no corresponding toggle or field-energy difference violates the closed energy ledger.

7 Laboratory Evidence Conditions

A force experiment has physical standing only if it supplies:

1. an instrument-independent measurement of $S_{\text{coh}}^{\text{can}}$ or its declared engineering coordinate;
2. the dimensional calibration $q_\star$;
3. a measured ambient repair gradient or a controlled emitted repair field;
4. a nonzero integrated repair charge, or a measured outgoing repair-current and momentum flux;
5. a complete toggle, maintenance-power, and field-energy ledger;
6. detuned, dummy, reversed-sign, thermal, acoustic, electromagnetic, mechanical, and center-of-mass controls;
7. a force result that follows the repair-field geometry and sign law in Eq. (4.9).
8. for any claimed switchable reversal, a derived signed source factor or an independently measured reversal of the external repair gradient.

Without these receipts, a null result does not measure χ_ν^{can} , and a nonzero balance signal does not establish repair charge. A test of χ_ν^{can} requires the action, source calibration, and operational source observable listed above.

8 Conclusion

The canonical coherent-matter susceptibility is an order-one channel coefficient fixed by protected-reserve survival on the stated OPH branch. The repair-charge condensate completion gives that coefficient a lawful dynamical role: it weights the coherent source in the repair-current equation. The same proposed action yields exact finite continuity, a dilute dust limit, the spherical deep-galaxy scaling, repair-field stress, and a device-force law. A device force depends on the dimensional calibration, integrated repair charge, external field, and field-momentum ledger. Repulsive response is conditionally allowed. Reactionless lift from a compact neutral contrast is excluded, the closed-compact-device formula $g^2 A \chi_\nu \Delta S / (4\pi G)$ is not a force theorem, and no laboratory force prediction follows from this paper.

References

- [1] B. Mueller et al., *From Observer Consensus to Standard Physics*. https://wkaxfdgxoqmgwhgshymt.supabase.co/storage/v1/object/public/papers/from_observer_consensus_to_standard_physics.pdf
- [2] B. Mueller et al., *Deriving Standard Model Gauge Structure from Observer Overlap Consistency*. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/deriving_standard_model_gauge_structure_from_observer_overlap_consistency.pdf
- [3] B. Mueller et al., *Federated Echoshedral Screen Microphysics: Patch Hardware, Records, and Observer Synchronization in OPH*. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/screen_microphysics_and_observer_synchronization.pdf
- [4] B. Mueller and D. Matscheko, *Observer-Patch Holography and the Dark Matter Phenomenon: A Repair-Charge Condensate Completion*. https://github.com/FloatingPragma/observer-patch-holography/blob/main/cosmology/oph_dark_matter_paper.pdf