

Boltzmann Transport from Observer-Patch Consistency: a Theorem-Conditional Reduction

OPH cosmology program

Release r1564, July 20, 2026

Abstract

This note shows that standard general-relativistic photon–baryon Boltzmann transport follows on the Observer-Patch Holography (OPH) branch when its geometry, electromagnetic action, physical photon, electron mass, and recombination history are supplied. It locates the repair-exchange and collar-stress interfaces where an OPH-specific effect could enter. A physical cosmic microwave background (CMB) prediction requires finite source receipts, transfer functions, and likelihood closure.

1 Claim boundary

Everything below is conditional on the branch theorems it cites. The derivation upgrades the status of the *transport form* used by the diagnostic CAMB lane from “imported wholesale” to “theorem-conditional reduction with declared imports.” It does not certify the initial spectrum (amplitude or tilt), the recombination history, the absolute Thomson normalization, or any likelihood statement. Those remain the gates tracked in issues #330, #522, #374, and #363 respectively. No statement in this note reads Planck data.

2 Branch inputs

Assumption 2.1 (A-GEOM: Einstein branch). *The observer-facing normal form carries a Lorentzian metric g_{ab} satisfying $G_{ab} + \Lambda g_{ab} = 8\pi G_{\text{geom}} \langle T_{ab} \rangle$, with $G_{\text{geom}} = \ell_\star^2$ from the edge-entropy/area-law theorem and $\Lambda \ell_\star^2 = 3\pi/N_\star$ from capacity closure. $N_\star$ is the SL-4 estimate read from measured Λ with the scale bridge of the strange-loop principle register; Λ is an input to this assumption, not an OPH output. Promotion of $N_\star$ beyond the SL-4 estimate runs through generator G2 of the project consistency stack: constructing the readback map F and its contraction certificate. Provenance: branch theorem (paper stack A22/A5); not re-proved here.*

Assumption 2.2 (A-QED: electromagnetic action and photon-pole receipt). *The realized branch carries the Standard Model quotient and the Maxwell kinetic action with a U(1) vertex of strength $\alpha = \alpha_{\text{root}}$ (source-side). In addition, this transport note assumes a positive-energy physical QED quantization with two transverse states and a positive-residue isolated pole at $k^2 = 0$ on the ordinary Lorentz-vacuum, unbroken electromagnetic phase. This extra photon-pole receipt is not implied by the reconstructed abstract group. Charged-lepton ratios are branch outputs; the absolute electron mass m_e is empirically anchored. Provenance: branch theorem (A20/A21), the explicit quantum-pole assumption just stated, and one declared import (m_e).*

Assumption 2.3 (A-BG: background). *The homogeneous limit of the branch is spatially flat FRW, $ds^2 = a^2(\tau) [-d\tau^2 + \delta_{ij}dx^i dx^j]$, with Friedmann dynamics from Assumption 2.1. Scalar perturbations are written in the conformal Newtonian gauge,*

$$ds^2 = a^2(\tau) \left[-(1 + 2\psi) d\tau^2 + (1 - 2\phi) \delta_{ij} dx^i dx^j \right],$$

with conventions of Ma–Bertschinger (1995). Overdots are ∂_τ .

Assumption 2.4 (A-REC: recombination import). *The free-electron history $n_e(\tau)$ (equivalently $x_e(\tau)$) is computed from standard atomic physics (Saha/Peebles-class) using imported atomic data. This is the same import class as the cesium-clock wall in the gravity stack: it contains no CMB-target information, so it is a permissible declared input under the anti-fitting rule, but it is an import, not a branch output.*

Assumption 2.5 (A-INIT: initial conditions (gated)). *Initial curvature perturbations $\zeta(\mathbf{k})$ are Gaussian with spectrum $\Delta_\zeta^2(k) = A_s (k/k_*)^{n_s-1}$. The Gaussian and statistically isotropic character is the MaxEnt screen-covariance theorem; the finite tilt certificate is issue #522. The radial theorem packet in issue #330 proves that one-shell inversion is nonunique and supplies physical source-dilation and cross-covariance tomography as conditional uniqueness routes. Finite source instantiation in issue #579 is work in progress. This note consumes A_s and n_s as gated inputs and derives no initial spectrum.*

3 Theorem B1: conditional refinement Liouville transport

Definition 3.1 (One-photon record density). *Conditional on the physical pole in Assumption 2.2, let $\mathcal{E}_\gamma$ be its positive-energy photon sector. Between record commits, a single excitation is a record continued across patch overlaps. Coarse-graining patch position and overlap momentum labels defines the phase-space record density $f(x^i, q, \hat{n}, \tau)$ on the forward mass shell, where q is comoving momentum magnitude and $\hat{n}$ the direction.*

Theorem 3.2 (Refinement Liouville). *On a branch with the Maxwell action, quantum photon-pole receipt, and overlap Lorentz kinematics (A-QED, A-GEOM), let P_r be the quotient-lumpable free-flow transition kernel at step h_r . For every frozen smooth compactly supported test function ϕ , set*

$$L_r \phi = \frac{P_r \phi - \phi}{h_r}, \quad \Gamma_r(\phi) = \frac{P_r(\phi^2) - (P_r \phi)^2}{h_r}.$$

Receipt BOLTZ-R1 consists of compact containment/tightness, convergence of initial laws, uniform-on-compact generator convergence $L_r \phi \rightarrow X_H \phi$ to the null-geodesic Hamiltonian vector field, vanishing quadratic variation $\Gamma_r(\phi) \rightarrow 0$, uniqueness of the Hamiltonian ODE, refinement and schedule naturality, and vanishing boundary and phase-space-measure defects. Under this receipt, the between-commit record flow satisfies, in the refinement limit,

$$\frac{Df}{d\tau} \equiv \frac{\partial f}{\partial \tau} + \frac{dx^i}{d\tau} \frac{\partial f}{\partial x^i} + \frac{dq}{d\tau} \frac{\partial f}{\partial q} + \frac{d\hat{n}}{d\tau} \cdot \frac{\partial f}{\partial \hat{n}} = C[f],$$

where the characteristics are the null geodesics of g_{ab} and $C[f]$ is supported on record commits only. In particular the collisionless flow preserves the Liouville measure.

Proof sketch, with the open receipt named. For every test function, the discrete process minus the time integral of $L_r \phi$ is a martingale whose quadratic variation is controlled by $\Gamma_r(\phi)$. Tightness gives

subsequential limits and vanishing quadratic variation makes every limiting martingale deterministic. Generator convergence says that each limit solves the martingale problem for X_H , equivalently the null-geodesic Hamiltonian ODE. Uniqueness identifies the whole limit, independently of the accepted repair schedule. Thus $dx/d\tau$, $dq/d\tau$, and $d\hat{n}/d\tau$ are the geodesic-flow coefficients. On the background of Assumption 2.3, $dq/d\tau = q[\dot{\phi} - \dots]$ has the form used in §5.

Measure preservation. The identified limit is Hamiltonian flow on the mass shell, so it preserves the canonical Liouville measure. The finite measure-defect clause checks that the quotient measures converge to that canonical measure rather than to a different coarse volume form. Accepted repair confluence only removes schedule dependence of committed endpoints. It supplies neither an inverse between-commit dynamics nor a phase-space Jacobian. The orientation doubling of a finite label register is bookkeeping and does not produce either property.

Receipt status. Photon-sector convergence of the finite patch flow to the displayed generator and zero-quadratic-variation limits is work in progress. The abstract Lean layer proves the corresponding schedule-independence and refinement-naturality statement in its setting. The photon receipt is named BOLTZ-R1, and §7 gives its simulator test. Theorem 3.2 is conditional on this receipt. $\square$

Corollary 3.3 (Collision and diffusion alternatives). *If the finite generator decomposes as $L_r = L_r^{\text{free}} + C_r$ and $C_r\phi \rightarrow C\phi$, then the limiting density obeys*

$$\partial_\tau f + \{f, H\} = C^* f.$$

A nonzero limit of Γ_r produces a diffusion/Fokker–Planck term and does not satisfy the deterministic Liouville conclusion.

Remark 3.4. The transport model places the collision functional at record commits and uses the independently certified measure-preserving flow between them. Claims about entropy production therefore refer to the commit terms counted by $C[f]$. The accepted repair relation and the reversible equilibrium or between-commit dynamics remain separate structures.

4 Theorem B2: record commits give the Thomson collision functional

Theorem 4.1 (Scalar-intensity commit-collision functional). *Let commits between photon-class and electron-class records occur with the soft-limit matrix element of the recovered U(1) vertex (A-QED). Then, to first order in perturbations and at low occupancy, the collision functional in Theorem 3.2 is the Thomson functional: in the electron rest frame,*

$$C[f](q, \hat{n}) = a n_e \sigma_T \left[\frac{3}{16\pi} \int d\Omega' (1 + (\hat{n} \cdot \hat{n}')^2) f(q, \hat{n}') - f(q, \hat{n}) \right], \quad \sigma_T = \frac{8\pi}{3} \frac{\alpha^2}{m_e^2},$$

boosted to the frame where baryons move with velocity v_b . The source-derived reversible proposal kernel is assumed to obey detailed balance on the angular state space. Write/check orientation labels alone do not imply that property, and the accepted strict-descent repair relation is not the equilibrium kernel. For every fixed photon energy q , isotropic angular distributions are stationary under this Thomson operator. This displayed scalar-intensity kernel does not by itself derive the Stokes-space Thomson matrix needed for polarization.

Proof sketch. At low occupancy the commit rate between momentum cells is bilinear in the occupancies with kernel given by the squared vertex; the soft (photon energy $\ll m_e$) limit of the recovered $U(1)$ vertex is classical Thomson scattering with the stated total cross-section and $1 + \cos^2$ angular kernel; this step is standard QED evaluated on the branch couplings and is not repeated here. The detailed-balance receipt equates the stationary weights of each reversible proposal pair. The resulting symmetric rest-frame kernel damps angular anisotropy at each fixed q . It does not mix frequencies, so its stationary kernel contains every isotropic spectrum $f(q)$ rather than a unique Bose or Planck occupancy. Momentum conservation per commit, summed over commits, gives equal and opposite momentum transfer densities to the electron–baryon fluid, with coefficient fixed by $\bar{\rho}_\gamma/\bar{\rho}_b$ bookkeeping; this is the Compton drag used in §5. $\square$

Remark 4.2 (Polarization provenance). The polarization hierarchy used below imports the standard polarized Thomson scattering matrix acting on Stokes brightness. The scalar $1 + \cos^2 \theta$ kernel displayed in Theorem 4.1 is its intensity projection, not a derivation of the Q/U collision terms. A source-level OPH derivation of the polarized commit kernel from the recovered vertex is work in progress. The polarization sector and the associated 16/15 diffusion factor are textbook imports rather than consequences of the displayed operator.

Remark 4.3 (Blackbody provenance). The Planck background spectrum is an initial-condition and thermal-history import in this note, not a consequence of the displayed Thomson functional. Even recoil-inclusive Comptonization conserves photon number and generically approaches a Bose–Einstein spectrum with a chemical potential. Selecting the zero-chemical-potential Planck spectrum requires the appropriate early thermalization history, including photon-number-changing channels such as double Compton scattering and bremsstrahlung, or an independently justified blackbody initial condition. An OPH derivation of the blackbody spectrum requires those channels and their rates rather than inferring spectral uniqueness from angular detailed balance.

Remark 4.4 (Provenance of the normalization). $\alpha = \alpha_{\text{root}}$ is source-side. m_e is an import (Assumption 2.2): OPH supplies no source-derived charged-lepton absolute mass, so σ_T ’s normalization is import-tagged (BOLTZ-N1 in the ledger). $n_e(\tau)$ is import-tagged by Assumption 2.4 (BOLTZ-X1).

Remark 4.5 (OPH-specific channels). Two and only two places exist where OPH kernels can modify this functional at linear order: (i) a number-changing repair-exchange channel $\Gamma_{\text{rec}}(k, a)$ adds a monopole source and a spectral-distortion term; it enters $C[f]$ additively and is gated by the #374 receipts (active fiber, physical clock, common-parent response); (ii) a collar-packet stress $\rho_A, B_A(k, a)$ enters only the gravitational source side (the Einstein constraints in §5), never $C[f]$. With both gates closed at zero, the transport below is exactly standard.

5 Theorem B3: the linear multipole hierarchy

Theorem 5.1 (Reduction to the standard hierarchy). *Linearize Theorems 3.2–4.1 on the background of Assumption 2.3 with scalar perturbations, expand the temperature brightness $F_\gamma(\mathbf{k}, \hat{n}, \tau)$ and polarization brightness G_γ in Legendre multipoles with $\mu = \hat{k} \cdot \hat{n}$. Writing $\delta_\gamma = F_{\gamma 0}$, $\theta_\gamma = \frac{3}{4}kF_{\gamma 1}$,*

$\sigma_\gamma = \frac{1}{2}F_{\gamma 2}$, and $\dot{\kappa} \equiv an_e\sigma_T$, the system is (Ma–Bertschinger 1995, eq. 63):

$$\begin{aligned}\dot{\delta}_\gamma &= -\frac{4}{3}\theta_\gamma + 4\dot{\phi}, \\ \dot{\theta}_\gamma &= k^2\left(\frac{1}{4}\delta_\gamma - \sigma_\gamma\right) + k^2\psi + \dot{\kappa}(\theta_b - \theta_\gamma), \\ 2\dot{\sigma}_\gamma &= \frac{8}{15}\theta_\gamma - \frac{3}{5}kF_{\gamma 3} - \frac{9}{5}\dot{\kappa}\sigma_\gamma + \frac{1}{10}\dot{\kappa}(G_{\gamma 0} + G_{\gamma 2}), \\ \dot{F}_{\gamma\ell} &= \frac{k}{2\ell+1}\left[\ell F_{\gamma(\ell-1)} - (\ell+1)F_{\gamma(\ell+1)}\right] - \dot{\kappa}F_{\gamma\ell}, \quad \ell \geq 3, \\ \dot{G}_{\gamma\ell} &= \frac{k}{2\ell+1}\left[\ell G_{\gamma(\ell-1)} - (\ell+1)G_{\gamma(\ell+1)}\right] + \dot{\kappa}\left[-G_{\gamma\ell} + \frac{1}{2}(F_{\gamma 2} + G_{\gamma 0} + G_{\gamma 2})\left(\delta_{\ell 0} + \frac{\delta_{\ell 2}}{5}\right)\right],\end{aligned}$$

coupled to baryons,

$$\dot{\delta}_b = -\theta_b + 3\dot{\phi}, \quad \dot{\theta}_b = -\frac{\dot{a}}{a}\theta_b + c_b^2k^2\delta_b + k^2\psi + \frac{4\bar{\rho}_\gamma}{3\bar{\rho}_b}\dot{\kappa}(\theta_\gamma - \theta_b),$$

and to the Einstein constraints of Assumption 2.1,

$$k^2\phi + 3\frac{\dot{a}}{a}\left(\dot{\phi} + \frac{\dot{a}}{a}\psi\right) = -4\pi Ga^2\delta\rho_{\text{tot}}, \quad k^2(\phi - \psi) = 12\pi Ga^2(\bar{\rho} + \bar{p})\sigma_{\text{tot}},$$

where $\delta\rho_{\text{tot}}$ and σ_{tot} include the gated ρ_A, B_A contributions of Remark 4.5. These contributions are zero on the receipt-failing branch. With $\Gamma_{\text{rec}} = B_A = 0$ this is verbatim the standard Einstein–Boltzmann system.

Proof. For scalar intensity, insert the geodesic characteristics of Theorem 3.2 on the perturbed metric, the collision functional of Theorem 4.1, keep first order, and project on Legendre multipoles. The $G_{\gamma\ell}$ equations additionally use the imported polarized Thomson matrix stated above. The remaining steps are the textbook computation (Ma–Bertschinger 1995; Dodelson, *Modern Cosmology*); no OPH-specific term survives except the two gated channels, which we have displayed at their unique insertion points. $\square$

6 Acoustic normal form, damping, and the line of sight

Proposition 6.1 (Acoustic oscillations). *In the tight-coupling regime $\dot{\kappa} \gg k, \frac{\dot{a}}{a}$, define $R \equiv 3\bar{\rho}_b/4\bar{\rho}_\gamma$ and $\Theta_0 = \delta_\gamma/4$. To leading order in $1/\dot{\kappa}$,*

$$\frac{d}{d\tau}\left[(1+R)\dot{\Theta}_0\right] + \frac{k^2}{3}\Theta_0 = -\frac{k^2}{3}(1+R)\psi + \frac{d}{d\tau}\left[(1+R)\dot{\phi}\right],$$

a driven oscillator with sound speed $c_s^2 = 1/[3(1+R)]$. The WKB solution gives, at last scattering τ_ ,*

$$[\Theta_0 + \psi](\tau_*) \propto \cos(kr_s(\tau_*)), \quad r_s(\tau_*) = \int_0^{\tau_*} c_s d\tau,$$

so extrema sit at $k_n r_s(\tau_) = n\pi$: compressions (odd n) are baryon-enhanced, rarefactions (even n) suppressed, and the dark-matter driving of the potentials boosts the third peak: every feature of the measured TT curve is produced by this system and by no OPH-specific term.*

Proposition 6.2 (Silk damping). *At next order in $1/\dot{\kappa}$ the oscillator acquires the diffusion envelope e^{-k^2/k_D^2} with*

$$k_D^{-2} = \int_0^\tau \frac{d\tau'}{6(1+R)\dot{\kappa}} \left[\frac{R^2}{1+R} + \frac{16}{15} \right],$$

the 16/15 arising from the polarization terms of Theorem 5.1; photon diffusion between commits is the physical mechanism, and its OPH reading is transparent: on scales crossed by a random walk of commit-free transport legs, the record field cannot maintain contrast.

Proposition 6.3 (Line-of-sight solution). *With visibility $g(\tau) = \dot{\kappa} e^{-\kappa(\tau, \tau_0)}$, the temperature multipoles today are*

$$\Theta_\ell(k, \tau_0) = \int_0^{\tau_0} d\tau \left\{ g \left[\Theta_0 + \psi + \frac{\Pi}{4} \right] j_\ell(k(\tau_0 - \tau)) + \left(\frac{g v_b}{k} \right) \dot{j}_\ell + \frac{3}{4k^2} (g\Pi)'' j_\ell + e^{-\kappa} (\dot{\psi} + \dot{\phi}) j_\ell \right\},$$

($\Pi = F_{\gamma 2} + G_{\gamma 0} + G_{\gamma 2}$; every unlabelled j_ℓ has argument $k(\tau_0 - \tau)$), and

$$C_\ell^{TT} = 4\pi \int \frac{dk}{k} \Delta_\zeta^2(k) \left| \frac{\Theta_\ell(k, \tau_0)}{\zeta(k)} \right|^2.$$

Given Assumption 2.5's gated (A_s, n_s) and the imports of Assumptions 2.2–2.4, this reproduces the curve computed by the diagnostic CAMB lane; agreement of an independent implementation with CAMB at the sub-percent level is receipt BOLTZ-V1.

7 Receipts and simulator tests

Receipt	Statement	Status
BOLTZ-R1	Finite-lattice refinement Liouville: coarse-grained photon-class record transport has quotient lumpability, tightness, convergent initial laws and first conditional moments, vanishing scaled second conditional moments, generator convergence on a frozen test basis, unique geodesic characteristics, and vanishing schedule, refinement, boundary, mass-shell, and measure defects. This receipt is independent of repair confluence and of the orientation-doubled label count.	open; simulator test below
BOLTZ-N1	σ_T normalization: absolute m_e is a declared import; the charged-family absolute-scale theorem is work in progress.	declared import
BOLTZ-X1	$x_e(\tau)$ recombination history from imported atomic data; contains no CMB-target information.	declared import
BOLTZ-V1	Independent hierarchy implementation matches CAMB TT/TE/EE at stated tolerances with identical inputs.	open; engineering
Tilt certificate Amplitude + lift	Theorem-grade n_s (clock normalization). A_s and radial screen-to- k map.	issue #522 conditional radial mathematics #330; finite source receipt #579 is work in progress
Kernels	$\Gamma_{\text{rec}}, \rho_A, B_A$ physical receipts.	issue #374

Simulator test for Boltz-R1. Freeze a smooth compactly supported test basis. At each regulator estimate the first conditional moment $\hat{L}_r\phi$ and scaled conditional variance $\hat{\Gamma}_r(\phi)$, together with quotient-measure, mass-shell, schedule, refinement, boundary-leakage, and collision/free-channel residuals. Pass requires $\hat{L}_r\phi \rightarrow X_H\phi$ and $\hat{\Gamma}_r(\phi) \rightarrow 0$ on the frozen basis with the preregistered uniform compact envelopes. Random schedule packet tests are supplemental. A nonzero second-moment limit selects the diffusion alternative rather than the Liouville theorem.

8 What this note does and does not close

It records, conditionally on the cited branch theorems, the *scalar angular form* of the transport and the locations where the two gated OPH kernels would enter. The polarization hierarchy, blackbody background, absolute normalizations, and recombination history remain imported. Thus the physical-CMB prediction gate contains a source-level polarized collision bridge, blackbody/thermalization provenance, BOLTZ-R1, and BOLTZ-V1, in addition to the pre-existing gates (#330, #522, #374, #363). It does not certify initial conditions, normalizations, recombination, or likelihoods, and nothing here reads Planck data. Under the anti-fitting rule, the acoustic peak structure remains attributable to standard plasma physics; OPH's falsifiable exposure stays concentrated in the primordial inputs, exactly where issues #330 and #522 put it.

References

- C.-P. Ma and E. Bertschinger, *Cosmological Perturbation Theory in the Synchronous and Conformal Newtonian Gauges*, ApJ 455, 7 (1995).
- S. Dodelson and F. Schmidt, *Modern Cosmology*, 2nd ed., Academic Press (2020).
- U. Seljak and M. Zaldarriaga, *A Line-of-Sight Integration Approach to Cosmic Microwave Background Anisotropies*, ApJ 469, 437 (1996).
- W. Hu and M. White, *CMB Anisotropies: Total Angular Momentum Method*, Phys. Rev. D 56, 596 (1997).
- OPH paper stack: *Observers Are All You Need*; *Reality as a Consensus Protocol*; *Recovering Relativity and the Standard Model from Observer Overlap Consistency*; *OPH Cosmology Finite-Source CMB Program* (this repository).