

Observer-Patch Holography Cosmology Data and Likelihood Contracts

Provenance, Comparability, and Claim Boundaries

B. Müller

August 30, 2026

Paper release: r2033 **Released:** August 30, 2026

Author affiliation: Bernhard Mueller, Pragma Research Inc.

Abstract

This paper separates the conditional Observer-Patch Holography (OPH) source theorem from a physical cosmology result. A curve overlay or reduced residual is a diagnostic. A physical result requires one finite source instantiation, compatible transfer, declared data and covariance, and complete nuisance treatment. The source, transfer, and likelihood package is not supplied here.

1 Comparison Classes

Every reported number belongs to one of four classes:

1. **source-derived:** all physical inputs follow from the declared OPH construction without using the comparison data;
2. **imported:** established background or microphysical inputs are declared explicitly;
3. **fit:** one or more parameters are inferred from the same data;
4. **diagnostic:** the calculation tests shape, scale, code behavior, or an interface without supporting a physical OPH claim.

The cosmology comparisons in scope are diagnostic or imported. No source-derived joint cosmology likelihood exists.

The exact finite source result computes canonical source height from authenticated direct-parent edges and proves strict increase along generated ancestry. Independently, a real axis and the rank-three source quotient define a one-plus-three ambient Lorentzian precursor. An explicit positive **timeScale** on source height enters only through event placement. This does not determine intrinsic poset dimension or supply a physical clock or continuum, background solution, conserved cosmological stress, initial conditions, transfer functions, or an observable likelihood. Those objects remain subject to the model-information and no-data-use rules below.

2 Required Model Information

A physical comparison must state:

1. the covariant action and source map;
2. the background solution and species content;
3. the gauge-invariant initial conditions;
4. the perturbation, collision, and exchange equations;
5. the observable projection and numerical tolerances;
6. every external input and fitted parameter.

For the repair-charge dark-sector proposal, the dilute dust-like equation of state is a conditional action consequence. The abundance, relativistic stress, initial perturbations, lensing map, clusters, and nonlinear structure equations are not supplied here.

3 Required Data Information

A comparison must identify the data release, selection, masks, covariance, calibration model, nuisance parameters, priors, and combination rule. Dataset overlap and shared calibration must be included. A diagonal sum is not a joint likelihood when the covariance is non-diagonal.

The model and data layers are separate. Measured values that select a source formula, normalization, branch, or applicability domain are inputs to that comparison rather than OPH outputs.

4 Observable Contracts

Observable family	Required theory output	Required comparison object
Cosmic microwave background (CMB)	C_ℓ^{TT} , C_ℓ^{TE} , C_ℓ^{EE} , lensing, foreground model	map or band-power likelihood with covariance and nuisance model
Baryon acoustic oscillations (BAO) and supernovae	$H(z)$, $D_A(z)$, luminosity distance, sound horizon	survey likelihood with calibration covariance
Weak lensing	metric potentials, nonlinear matter power, intrinsic alignments	tomographic likelihood with masks and covariance
Growth and redshift-space distortions (RSD)	$P_m(k, z)$ and $f\sigma_8(z)$	windowed survey likelihood with cross-bin covariance
Clusters	mass function, selection, observable–mass map, lensing calibration	count and calibration likelihood
Galaxies	disk dynamics, gas and stellar nuisances, external environment	galaxy-level likelihood with selection and covariance

5 Diagnostic Results

The reported CMB overlays, compressed background comparisons, and galaxy scaling checks are diagnostic. Their formulas or inputs are selected with knowledge of the comparison data, omit required covariance or nuisance structure, or lack a physical source map. They test numerical paths and conditional formulas and do not establish a cosmological prediction.

6 No-Data-Use Boundary

A source-derived calculation must expose its dependency graph. The source side may use mathematical constants, declared OPH finite artifacts, and explicitly imported physical inputs. It may not use the observed quantity being claimed as an output, a fit to that quantity, or a proxy calibrated from the same dataset.

This provenance rule is enforced through the declared dependency graph and data-use classification.

7 Theorem scope

The screen-spectrum and radial-lift mathematics are conditional theorems. One-shell inversion is nonunique; physical source dilation and radial cross-covariance tomography are the two uniqueness routes. Finite source instantiation, repair-charge cosmological source, Boltzmann bridge, and joint likelihood are not supplied. All cosmological data products are diagnostic. No cosmological result carries an OPH falsification verdict.

AI Assistance Disclosure

This research project used research-grade commercial models, including Anthropic’s Fable and OpenAI’s GPT-5.6-Sol, for research support, software development, editing, and synthesis. The authors are responsible for the paper’s claims, methods, and final text.