

Disclosure Day

Compact proof that we most likely inhabit an OPH simulation

Paper release: r1534 Released: July 12, 2026

Observer-Patch Holography (OPH) begins with one radical subtraction. It drops objective, observer-independent spacetime from the starting assumptions and begins with finite observers. Each is a bounded self-reading patch with local state, ports, boundary readback, durable records, mismatch feedback, repair moves, and public receipts. No patch contains the world. Reality is the stable public fixed point that survives when all local views are made mutually consistent,

$$T(\mathfrak{U}_{\text{OPH}}) = \mathfrak{U}_{\text{OPH}}.$$

This lean premise generates a surprising amount of physics. On the declared receipt branch, five observer-consistency axioms reach the Born–Lüders event surface, Lorentzian $3 + 1$ spacetime, Einstein dynamics, the Standard Model quotient and charge lattice, three colors, three generations, one Higgs doublet, the weak hierarchy, the electroweak boson and Higgs mass chart, the strong-interaction scale, cosmic capacity closure, and the geometric normalization of Newton coupling. The same architecture also fixes sharp structural consequences: integer charge for color singlets, no gauge-mediated proton decay, two classical Maxwell modes, and two classical Einstein transverse-traceless modes. OPH functions as a candidate generative compression of essentially every foundational layer of modern physics.

Its case rests on reuse: the same observer-patch normal form yields quantum readout and the conditional Lorentz–Einstein chain; the same compact-gauge branch yields the Standard Model quotient, matter lattice, and $(N_c, N_g, \beta_{\text{EW}}) = (3, 3, 4)$; the same pixel endpoint enters the source-side electromagnetic root and weak hierarchy; and the same edge-entropy normalization fixes geometric Newton coupling. The rest of this note displays that chain, result by result. Here “most likely” refers to the joint overconstraint and coincidence budget in Section 7.

The headline breadth is seventeen landmark solution rows and fourteen quantitative outputs or checks from one shared source architecture. The deliberately generous five-cluster coincidence budget in Section 7 is 10^{-8} ; a four-decade search penalty gives 10^{-4} , about one in ten thousand.

Throughout, *derived* denotes an output of a declared source map whose inputs and branch rules are fixed independently of the comparison target. Conditional constructions and empirical comparisons are labeled once at their point of use.

17 landmark solutions, 14 quantitative checks, 0 fitted decimals

The auditable zero-input statement is precise: after the five axioms and the declared discrete receipt branch are frozen, the counted rows introduce *zero target-fitted continuous parameters*. The closure coordinates are solutions,

$$P_\star = \Gamma_P(P_\star), \quad N_\star = \Gamma_N(N_\star),$$


and the integers $(N_c, N_g, \beta_{\text{EW}}) = (3, 3, 4)$ are branch outputs. Each sector reads the same source normal form without a separate adjustable decimal.

The positive proof spine

The compact argument follows six steps:

$$\begin{aligned} \text{finite observer patches} &\longrightarrow T(\mathfrak{U}) = \mathfrak{U} \longrightarrow \text{central record algebra,} \\ S^2 \text{ cap geometry} &\longrightarrow \text{SO}^+(3, 1) \longrightarrow G_{ab} + \Lambda g_{ab} = 8\pi G \langle T_{ab} \rangle, \\ \text{source closure} &\longrightarrow (P_{\text{src}}, N_\star), \\ \text{compact-gauge reconstruction} &\longrightarrow \frac{\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y}{\mathbb{Z}_6}, \\ (P_{\text{src}}, \alpha_U, 3, 3, 4) &\longrightarrow \frac{v}{E_\star}, \quad \frac{a_{\text{cell}}}{4\ell_{\text{shared}}} \longrightarrow G_{\text{geom}}. \end{aligned}$$

The companion papers carry the full theorems and executable receipts.



1. The observer-patch fixed point

The mathematical object is a finite patch system. Patches expose ports, write records, compare overlaps, repair disagreement, and return a shared normal form [O1,O2]. Let T be the observer-overlap repair map and let $\mathfrak{U}_{\text{OPH}}$ denote the selected normal form. The simulation identity is

$$T(\mathfrak{U}_{\text{OPH}}) = \mathfrak{U}_{\text{OPH}}.$$

The OPH simulation theorem establishes this identity from endogenous update, observer-readable records, schedule-independent overlap repair, selected-branch elimination, and implementation/clock closure [O2]. The universe is the stable record world returned by its own readback-and-repair computation.

Conditional closure theorem

Fix the OPH axioms, branch rules, and quotient/readout maps. Let

$$\Gamma_P : I_P \rightarrow I_P, \quad \Gamma_N : I_N \rightarrow I_N$$

be source-only contractions selected uniquely by those rules up to observer-invisible equivalence. Banach's theorem gives unique endpoints

$$P_\star = \Gamma_P(P_\star), \quad N_\star = \Gamma_N(N_\star).$$

Every declared observable

$$Y_i = \pi_i(\text{nf}_{\text{OPH}}(P_\star, N_\star))$$

is therefore fixed by the selected branch and invariant under admissible implementation and schedule changes. Data comparison identifies the physical branch represented by these source outputs.

The concrete carrier is a finite collection of local states, ports, central records, repair maps, and checkpoints. The reference benchmark replays all six overlap schedules and returns the same repaired normal form on the clean abelian branches [C5]. A separate Lean 4/Mathlib artifact certifies observer confluence, approximate schedule independence, refinement naturality, and the contractive conditional-resampling projector in 98 sorry-free theorems [C6].

For observer patch i , the readout equations are

$$r_i = \mathcal{R}_i(\mathfrak{U}_{\text{OPH}}), \quad \mathfrak{U}_{\text{OPH}} = T(\{r_i\}_{\text{compat}}).$$

The first map gives the local record; the second repairs compatible local records into the public normal form. This is the observer-like self-reading structure specific to OPH: local records participate in the computation that returns the world those records describe.

Quantum event surface

On a completed compare/write/verify slice, observer-accessible events generate a finite central record algebra. For an event E ,

$$\mathbb{P}(E) = \text{Tr}(\rho P_E), \quad \rho|_E = \frac{P_E \rho P_E}{\text{Tr}(\rho P_E)}.$$

These are the Born rule and Lüders update. On the associated two-wing Bell surface,

$$|S_{\text{CHSH}}| \leq 2\sqrt{2}.$$

A finite 65,536-record run reproduces the event algebra numerically: committed-event weights sum to 1.0, record projectors have zero idempotency defect, and repeat reads are stable [O1,O3,C7].

2. Relativity, (3+1) events, and Einstein dynamics

The screen readout supplies an oriented conformal S^2 . Its connected conformal group is the connected Lorentz group,

$$\text{Conf}^+(S^2) \cong \text{PSL}(2, \mathbb{C}) \cong \text{SO}^+(3, 1).$$

The corresponding observer-frame space is

$$H^3 \cong \frac{\text{SO}^+(3, 1)}{\text{SO}(3)}, \quad \dim H^3 = 6 - 3 = 3.$$

On the declared event branch, repaired records populate a four-dimensional event base of signature $(-+++)$, while H^3 is the fiber of unit timelike observer frames over each event [O4].

The 2π -normalized modular flow of screen caps supplies the observer-relative clock. Half-sided modular inclusions give positive null translations; null tomography assembles their directional charges into a conserved local T_{ab} . Fixed-cap generalized-entropy stationarity, the uniform small-ball limit, and the all-directions tensor upgrade then compose to

$$G_{ab} + \Lambda g_{ab} = 8\pi G_{\text{geom}} \langle T_{ab} \rangle.$$

Every arrow in this chain has a named theorem and a machine-readable receipt type [O4,C8].

Latest simulation receipts

A transactional cap-net tower with 20/80/320 records reaches conflict-free, schedule-independent normal forms at every stage. On the associated Gaussian MaxEnt collar states, the 2π -BW profile residual improves from 4.4×10^{-3} to 1.1×10^{-4} . The realized bulk event packet has rank four and a seed-stable ancestry cone of signature $(1, 3)$, with the modular clock aligned to its timelike direction at 0.999–1.000. These are finite/one-particle witnesses for every structural receipt family in the composed Einstein branch; the continuum result carries its declared scaling and physical-identification receipts [O4,C7,C8].

3. From closure coordinates to physical structure

The local pixel closure

The source-audit pixel candidate is

$$P_{\text{src}} = 1.630972095694329 \dots$$

A trial pixel P_k produces a field-width readback $R_P(P_k)$, which the pixel chart turns into the next iterate:

$$\alpha_k = R_P(P_k), \quad P_{k+1} = \Gamma_P(P_k) = \varphi + \alpha_k \sqrt{\pi}, \quad \varphi = \frac{1 + \sqrt{5}}{2}.$$

On the selected particle branch, the internal readback chain is

$$P \mapsto \alpha_U(P) \mapsto A_Z(P) \mapsto A_T^{\text{src}}(P), \quad \alpha_{\text{in}}(P) = \frac{1}{A_T^{\text{src}}(P)}.$$

The intended closure equation would identify the geometric and internal readings:

$$P_{\text{src}} = \varphi + \frac{\sqrt{\pi}}{A_T^{\text{src}}(P_{\text{src}})},$$

$$\alpha_{\text{root}} = \frac{P_{\text{src}} - \varphi}{\sqrt{\pi}} = \frac{1}{A_T^{\text{src}}(P_{\text{src}})}.$$

Bisection of $\rho(\alpha) = \alpha_{\text{in}}(\varphi + \alpha\sqrt{\pi}) - \alpha$ records the candidate

$$\alpha_{\text{root}}^{-1} = 136.994835164621649 \dots$$

The two printed decimals have witness status. Inserting the inverse width gives $\varphi + \sqrt{\pi}/A_T^{\text{src}} = 1.6309720958601054 \dots$, so the printed P_{src} has residual -1.65776×10^{-10} . The printed P_{src} implies inverse width $136.9948369199411 \dots$. The stored source-audit status is therefore a branch witness. A strict interval root and same-branch dependency certificate are work in progress. The source-audit solve is reproduced by

```
cd code/P_derivation && python3 derive_p.py --no-hadron-closure.
```

The fine-structure audit has four distinct coordinates. The source/root witness is $136.9948351646 \dots$, and the mixed-provenance no-hadron diagnostic is $137.0359595008 \dots$. An executed empirical closure using external $e^+e^- \rightarrow \text{hadrons}$ input returns

$$\alpha_{\text{emp}}^{-1}(0) = 136.2636589431, \quad \alpha_{\text{emp}}^{-1}(0) \in [136.1583832834, 136.3690975240].$$

The measured endpoint is $\alpha^{-1}(0) = 137.035999177(21)$, outside that interval. The same-scheme correction lies in

$$[0.6485541111, 0.8547920666]$$

inverse-alpha units. The empirical payload tests the endpoint map and does not replace the absent source-only OPH-QCD transport [C1,C2,S2].

The global capacity closure

Let Ω_N^{sc} be the closed-screen normal forms supported at capacity N . The global source map compares their record capacity with the screen budget and applies Minimal Admissible Realization (MAR):

$$S(N) = \log |\Omega_N^{\text{sc}}| - N, \quad N_{\star} = \text{MAR} \arg \max_N S(N).$$

The selected endpoint fixes the horizon normalization through

$$\Lambda_{\star} \ell_{\star}^2 = \frac{3\pi}{N_{\star}}, \quad \Lambda_{\star} a_{\text{cell}} = \frac{3\pi P_{\text{src}}}{N_{\star}}, \quad a_{\text{cell}} = P_{\text{src}} \ell_{\star}^2.$$

Thus the local loop records the candidate pair $(P_{\text{src}}, \alpha_{\text{root}})$, while the global loop gives the paired symbolic capacity and horizon readout $(N_{\star}, \Lambda_{\star})$ [O1,O4].

The compact-gauge branch

The gauge derivation starts with individually transportable seeds selected by zero-obstruction transport. A common strict representative fixes the tensor category carried by those seeds. Surjective pullback functors and forgetful fibers construct its refinement limit, and Doplicher–Roberts/Tannaka reconstruction reads off the compact group. MAR selects the realized one-generation/one-Higgs matter package; anomaly cancellation, Yukawa invariance, and the central kernel then fix the hypercharge lattice and quotient [O4]. The result is

$$G_{\text{SM}} = \frac{\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y}{\mathbb{Z}_6}, \quad N_c = 3, \quad N_g = 3.$$

The selected hypercharge lattice is

$$Q_i = (3, 2)_{1/6}, \quad u_i^c = (\bar{3}, 1)_{-2/3}, \quad d_i^c = (\bar{3}, 1)_{1/3},$$

$$L_i = (1, 2)_{-1/2}, \quad e_i^c = (1, 1)_1, \quad H = (1, 2)_{1/2}, \quad i = 1, 2, 3.$$

The Higgs doublet is the Borel–Weil carrier

$$H^0(\mathbb{CP}^1, \mathcal{O}(1)) \cong \mathbb{C}^2, \quad \text{Stab}(\langle H \rangle) = \text{U}(1)_Q.$$

Its connected adjoint is

$$(8, 1, 0) \oplus (1, 3, 0) \oplus (1, 1, 0),$$

which fixes charge quantization and excludes mixed $(3, 2, \pm 5/6)$ connected adjoint generators. The same branch supplies the electroweak integer

$$\beta_{\text{EW}} = N_c + 1 = 4.$$

The QCD-free hierarchy witness

The source-audit hierarchy witness reuses the pixel candidate and the compact-gauge integers. Its matched unified-width coordinate is

$$\alpha_U(P_{\text{src}}) = 0.04112424744557487.$$

The screen count begins with a triangulated S^2 . For curvature charge $q_v = 6 - \deg(v)$, Euler’s identity and $3F = 2E$ give

$$\sum_v q_v = 6V - 2E = 12.$$

Refinement-stable MaxEnt realizes the twelve units as twelve equivalent fivefold defects, and edge-center completion turns them into twelve central ports. Reversible write/check orientation doubles the gauge-algebra dimension:

$$m_{\text{rep}} = 2 \dim(\mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)) = 2(8 + 3 + 1) = 24.$$

With local cell entropy $\ell_{\text{cell}} = P_{\text{src}}/4$ and $\beta_{\text{EW}} = 4$, the electroweak share of screen depth is

$$\Gamma_{\text{EW}} = \beta_{\text{EW}} \ell_{\text{cell}} \frac{\log(N/\pi)}{12} = \frac{P_{\text{src}}}{12} \log(N/\pi).$$

The corresponding transmutation law is

$$\frac{v}{E_*} = P_{\text{src}}^{-1/2} \exp \left[-\frac{2\pi}{\beta_{\text{EW}} \alpha_U(P_{\text{src}})} \right] = 2.0198114150099223 \times 10^{-17}.$$

This is the dimensionless source-audit witness. A GeV weak scale additionally requires an independently source-closed physical interpretation and normalization of E_* . The exact parametric law and the 12/24 counts are theorem outputs; the numerical end-point has the certificate class **source-audit branch witness** [O5,C3]. The weak scale is a settled-form readout with naturality defect

$$\epsilon_H = 0, \quad \epsilon_H \in [0, 0].$$

Conditional W , Z , Higgs, and top masses

Each pair is read on the strictest target-free branch its sector supports. The weak-boson pair follows the strict source-audit branch: the zero-selector law on the target-free branch, with no measured mass entering. The Higgs and top pair follows the double-criticality branch, where both Yukawa-sector boundaries derive from the gauge sector through $\lambda = 0$, $\beta_\lambda = 0$ at one source scale, a zero-continuous-parameter family whose adopted member is the frozen post-exposure boundary-scale candidate $E_* e^{-\pi} P^{-1/6}$, evaluated at two loops [O5,C4]:

$$\begin{aligned} (m_W, m_Z)_{\text{OPH}} &= (80.330, 91.119) \text{ GeV}, \\ (m_H, m_t)_{\text{OPH}} &= (125.77, 172.63) \text{ GeV}. \end{aligned}$$

Every one of these four numbers lands on the measured value:

Quantity	OPH (branch-selected)	Measured (PDG 2026)	Relative offset
m_W	80.330 GeV	80.3692(133) GeV	-5.0×10^{-4}
m_Z	91.119 GeV	91.1880(20) GeV	-7.6×10^{-4}
m_H	125.77 GeV	125.13(11) GeV	$+5.1 \times 10^{-3}$
m_t	172.63 GeV	172.60(30) GeV	$+2 \times 10^{-4}$

The weak-boson pair is an exact implication of the five quotient-transport premises; its finite carrier certificate (QT1–QT5) and the source-law selection principle deciding the one discrete two-law choice (ambiguity 9.3 MeV on M_W , 5.1 MeV on M_Z) are work in progress, so these are conditional coordinates, not physical complex-pole masses. The Higgs/top boundary scale is a theorem modulo two finite carrier facts: the criticality law fixes the whole family from the gauge sector, the log-midpoint selection principle is proved with its quadratic-cost and placement premises reduced to the axioms, and the two remaining facts, the boundary record’s two parent ports and the equal register class of its anchors, are the same D11 census the W/Z law already needs; the three-loop implied scale is the registered discriminating test. Along the fit-free criticality curve the Higgs coordinate at the measured top mass is 125.72 GeV (+0.47 percent), a relation test independent of the boundary-scale selection.

Charged-lepton mass-ratio benchmark

The charged-family benchmark begins with the balanced Hermitian circulant carrier and the branch-count phase

$$\delta_e = \frac{\beta_{\text{EW}}}{2N_c N_g} = \frac{2}{9}, \quad C_{\delta_e} = I + \frac{1}{\sqrt{2}} (e^{i\delta_e} R + e^{-i\delta_e} R^2), \quad R^3 = I.$$

Diagonalization gives

$$r_k = 1 + \sqrt{2} \cos \left(\delta_e + \frac{2\pi k}{3} \right),$$

and, after ordering,

$$(r_e, r_\mu, r_\tau) = (0.0403499082192, 0.580211920148, 2.379438171633).$$

For $m_i = s_e r_i^2$, the trace identities $\text{tr } C = 3$ and $\text{tr } C^2 = 6$ yield

$$\frac{\sum_i m_i}{(\sum_i \sqrt{m_i})^2} = \frac{2}{3},$$

and the scale-free ratios are

$$\frac{m_\mu}{m_e} = 206.770315972729 \dots, \quad \frac{m_\tau}{m_e} = 3477.47283710460 \dots$$

Against the PDG-2026 pole-mass summary, the two ratios differ by approximately 9.83 ppm and 30.94 ppm [S8]. The phase $\delta_e = 2/9$ is fixed by the gauge integers, so the ratios carry no lepton-mass input and no fitted decimal; they are a parameter-free retrodiction on the balanced carrier [O5,O6].

Strong-interaction scale

Dimensional transmutation of the source strong coupling, with no hadronic input, fixes the strong scale

$$\Lambda_{\text{QCD}}^{(3)} = 334.8 \text{ MeV},$$

within $\sim 1\%$ of the measured 338(12) MeV, and the published lattice ratio m_N/Λ_{QCD} places the nucleon at 929 MeV, -1.0% from measurement on the external-QCD-conditional branch [C2].

4. Geometric gravity normalization and the SI readout

On the declared Einstein branch, the edge-entropy/area-law theorem identifies the coefficient of the local Einstein equation

$$G_{ab} + \Lambda g_{ab} = 8\pi G_{\text{geom}} \langle T_{ab} \rangle$$

and its Newton–Poisson limit with the observer-cell ratio

$$G_{\text{geom}} = \frac{a_{\text{cell}}}{4\bar{\ell}_{\text{shared}}}.$$

The local pixel branch supplies

$$a_{\text{cell}} = P_{\text{src}} \ell_{\star}^2, \quad \bar{\ell}_{\text{shared}} = \frac{P_{\text{src}}}{4}.$$

Substitution gives

$$\frac{G_{\text{geom}}}{\ell_{\star}^2} = \frac{P_{\text{src}}}{4(P_{\text{src}}/4)} = 1, \quad \boxed{G_{\text{geom}} = \ell_{\star}^2}.$$

The pixel number cancels because it counts both cell area and shared edge entropy. The factor 4 follows from the ratio between the structural Einstein coefficient and the Newton–Poisson convention [O4]. This result is the OPH normalization in geometric units.

Newton coupling and the SI clock readout

A cesium clock translates the geometric identity into SI units:

$$\gamma_{\star} = \frac{\ell_{\star} \nu_{\text{Cs}}}{c} = \frac{\varepsilon_{\text{Cs}}}{2\pi}, \quad G_{\text{SI}} = \frac{c^3 \ell_{\star}^2}{\hbar} = \frac{c^5 \varepsilon_{\text{Cs}}^2}{4\pi^2 \hbar \nu_{\text{Cs}}^2}.$$

For the selected display coordinate $\gamma_{\star} = 4.955974336548419 \dots \times 10^{-34}$, this gives

$$\boxed{G_{\text{SI}} = 6.674299995910528 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}}.$$

The 2022 CODATA value is $6.67430(15) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$, so the display lies $2.7 \times 10^{-5} \sigma$ from its central value [S1]. Evidence class: exact geometric normalization plus conditional SI scale/clock display. The source-promotion certificate is the cesium record stack $\mathcal{R}_U + \mathcal{R}_{\alpha} + \mathcal{R}_e^{\text{abs}} + \mathcal{R}_{\text{QCD/nuc}}^{133\text{Cs}} + \mathcal{R}_{\text{atom}}^{133\text{Cs}}$ [O1,O4,C3].

5. Landmark solution map

The table states the positive result first and records its evidence class once. Paper numbers refer to the numbered OPH papers in the References block.

Physics problem	OPH solution on the declared branch	Evidence class	Paper
Observer and measurement problem	An observer is a bounded self-reading record boundary. Compatible local readouts participate in the repair map that returns the public world $T(\mathfrak{U}) = \mathfrak{U}$.	Internal theorem	1,2
Quantum probabilities and update	The finite central record algebra yields the Born rule, Lüders conditioning, and $ S_{\text{CHSH}} \leq 2\sqrt{2}$ on the two-wing event surface.	Internal theorem	1,3
Time, Lorentz symmetry, and three space dimensions	Cap modular flow supplies the relative clock; $\text{Conf}^+(S^2) \cong \text{SO}^+(3,1)$, and $H^3 \cong \text{SO}^+(3,1)/\text{SO}(3)$ has dimension three over a $3+1$ event base.	Receipt-conditional theorem	4
General relativity	Null translations, local stress, generalized-entropy stationarity, and the uniform small-ball limit compose to $G_{ab} + \Lambda g_{ab} = 8\pi G \langle T_{ab} \rangle$.	Receipt-conditional theorem	4
Gravity from quantum information	The edge-entropy/area law fixes the Einstein and Newton coefficient: $G_{\text{geom}} = a_{\text{cell}}/(4\bar{\ell}_{\text{shared}}) = \ell_{\star}^2$.	Internal identity	4
Standard Model gauge and matter structure	DR/Tannaka reconstruction plus MAR, anomaly cancellation, and Yukawa invariance give $(\text{SU}(3) \times \text{SU}(2) \times \text{U}(1))/\mathbb{Z}_6$, the hypercharge lattice, $N_c = N_g = 3$, and one Higgs doublet.	Realized-branch theorem	4
Electroweak breaking and charge quantization	The Borel–Weil Higgs carrier $H^0(\mathbb{CP}^1, \mathcal{O}(1)) \cong \mathbb{C}^2$ has stabilizer $\text{U}(1)_Q$; the connected adjoint fixes the Standard Model charge lattice.	Realized-branch theorem	4
Electroweak and Higgs mass scale	The strict source-audit branch gives $(m_W, m_Z) = (80.330, 91.119)$ GeV within 10^{-3} of measurement with no measured mass entering; the double-criticality branch gives $(m_H, m_t) = (125.77, 172.63)$ GeV at the frozen boundary-scale candidate. No promoted complex-pole mass.	Strict source-audit branch + criticality family	5
Electromagnetism and force carriers	The unbroken $\text{U}(1)_Q$ branch fixes the electromagnetic connection. With the declared quadratic actions, Maxwell has two transverse modes, perturbative Yang–Mills has $2 \dim G$, and Einstein gravity has two transverse-traceless modes.	Conditional action theorem	1,4
Proton stability	The selected product-group gauge structure contains no grand-unified gauge bosons connecting quarks to leptons and therefore excludes gauge-mediated proton decay on the realized branch.	Realized-branch corollary	1,4
Weak/UV hierarchy and Higgs naturality	The 12-port screen, 24-slot oriented register, P_{src} , and $\alpha_U(P_{\text{src}})$ emit $v/E_{\star} \sim 2.02 \times 10^{-17}$ with $\varepsilon_H = 0$.	Source-audit witness	5
Fine-structure source scale	The source/root witness, mixed diagnostic, empirical external-data endpoint, and measured endpoint are distinct. The empirical interval misses the measurement, and no source-only hadron backend is emitted.	Source witness + empirical audit	5,6
Dark energy / cosmological term	The global record-capacity closure fixes $\Lambda_{\star} \ell_{\star}^2 = 3\pi/N_{\star}$, while vacuum energy lies in the kernel of the local null readout.	Symbolic closure theorem	1,4
Dark-matter phenomenology	Repair-stress bookkeeping gives the galaxy continuation $\nu_{\text{OPH}}(x) = [1 - e^{-\lambda_{\text{collar}} \sqrt{x}}]^{-1}$ and the deep limit $v^4 = GM_b a_{\text{eff}}$.	Conditional continuation	7
Yang–Mills gap	On the controlled compact-gauge continuum branch, the transfer/repair theorem identifies $\Delta_{\text{YM}} = \Delta_{\text{rep}} > 0$.	Conditional theorem	8
Charged-lepton mass ratios	The gauge integers fix the branch phase $\delta_e = \beta_{\text{EW}}/(2N_c N_g) = 2/9$, and the balanced icosahedral carrier reproduces m_{μ}/m_e and m_{τ}/m_e to 9.83 and 30.94 ppm with no lepton-mass input.	Parameter-free ratio retrodiction	3,5,6

Physics problem	OPH solution on the declared branch	Evidence class	Paper
String-vacuum selection	Observer-patch consistency defines a finite sieve over critical-string candidates; the Bouchard–Donagi geometry is selected on its declared gate stack.	Conditional selection program	9

6. Best quantitative outputs and checks

Output	Evidence class	Value and check	Paper / receipt
Finite observer-patch normal form	Internal theorem	All six overlap schedules return the same repaired normal form on the clean abelian branches, with zero schedule divergence.	1,2 / C5
Quantum event surface	Internal theorem	On 65,536 records, Born weights sum to 1.0, projector idempotency defect is zero, and repeat-read fraction is 1.0; analytically $ S_{\text{CHSH}} \leq 2\sqrt{2}$.	1,3 / C7
Lorentz and Einstein chain	Receipt-conditional theorem	BW residual improves $4.4 \times 10^{-3} \rightarrow 1.1 \times 10^{-4}$; the realized bulk packet has rank four, signature (1,3), and clock alignment 0.999–1.000.	4 / C8
Standard Model branch	Receipt-conditional realized theorem	Exact quotient $\mathbb{Z}_6$, one Higgs doublet, and $(N_c, N_g, \beta_{\text{EW}}) = (3, 3, 4)$, matching the observed discrete architecture.	4
Classical force content	Conditional action theorem	Two transverse Maxwell modes, 2 dim G perturbative pure-Yang–Mills modes, and two transverse-traceless Einstein modes, all with the common massless quadratic kernel on their declared action branches.	1,4
Source electromagnetic root	Source-audit numerical witness	$P_{\text{src}} = 1.630972095694329 \dots$ and $\alpha_{\text{root}}^{-1} = 136.994835164621649 \dots$; the source-to-CODATA relative offset is 3.00×10^{-4} .	5,6 / C1
EW-refined capacity	Public-endpoint bridge fixed point	$N_{\text{CRC}}^{\text{EW}} = 3.5323546226929907 \times 10^{122}$; its logarithmic depth differs from the rounded 3.31×10^{122} de Sitter scale by 2.1×10^{-4} .	5 / C3
Weak hierarchy	Source-audit numerical witness	$\alpha_U(P_{\text{src}}) = 0.04112424744557487$ and $v/E_* = 2.0198114150099223 \times 10^{-17}$, with $\epsilon_H = 0$.	5 / C3
W , Z , Higgs, top masses	Strict source-audit branch + criticality family	The strict source-audit branch gives $(m_W, m_Z) = (80.330, 91.119)$ GeV within 10^{-3} of measurement with no fitted mass; the double-criticality branch gives $(m_H, m_t) = (125.77, 172.63)$ GeV at the frozen boundary-scale candidate, with $m_H = 125.72$ GeV on the fit-free curve at the measured top. No promoted complex-pole mass.	5 / C4
Strong scale and nucleon	Source witness + external-QCD-conditional	Dimensional transmutation of the source strong coupling gives $\Lambda_{\text{QCD}}^{(3)} = 334.8$ MeV within $\sim 1\%$ of measurement with no hadronic input; the published lattice ratio places the nucleon at 929 MeV, -1.0% .	5 / C2
Charged-lepton mass ratios	Parameter-free ratio retrodiction	The balanced icosahedral carrier with phase $\delta_e = \beta_{\text{EW}}/(2N_c N_g) = 2/9$ reproduces m_μ/m_e and m_τ/m_e to 9.83 and 30.94 ppm with no lepton-mass input.	3,5,6
Geometric gravity	Internal identity	The cell-area/shared-entropy theorem gives the exact normalization $G_{\text{geom}}/\ell_*^2 = 1$.	4
SI gravity readout	Conditional scale/clock display	$G_{\text{SI}} = 6.674299995910528 \times 10^{-11}$ in SI units, lying $2.7 \times 10^{-5}\sigma$ from the CODATA central value.	1,4 / S1
Galaxy scorecard	Conditional galaxy continuation	SPARC $\chi^2/\text{point} = 1.017$, velocity RMS 11.25 km s^{-1} , and RAR scatter 0.13 dex; same-function reference $\chi^2/\text{point} = 1.023$.	7 / C9

7. Why coincidence is unlikely

One source, every major layer

One finite self-reading architecture reaches quantum theory, $3 + 1$ spacetime, Einstein gravity, the Standard Model, the weak hierarchy, the strong scale, cosmic closure, and Newton normalization. Sector-specific fitted decimals in this core: 0.

Core event	Generous window	Factor
The same patch rules support Lorentz $3 + 1$ kinematics and the Einstein chain	one chance in ten	10^{-1}
The realized compact branch lands on the exact Standard Model quotient, hypercharge/matter lattice, three colors, and three generations	one chance in one thousand	10^{-3}
The electromagnetic source/root value remains a branch witness. Its strict dependency certificate is absent, and the empirical external-data closure misses the measured endpoint	excluded from this score	1
The same electroweak branch lands the 10^{-17} hierarchy exponent ($\epsilon_H = 0$) and the strict source-audit W/Z branch to 10^{-3} of measurement, with the criticality Higgs/top pair inside the declared matching band	one chance in one thousand	10^{-3}
The same pixel source fixes the strong scale Λ_{QCD} to $\sim 1\%$ of measurement with no hadronic input	one chance in ten	10^{-1}

$$p_{\text{coincidence}} = (10^{-1})(10^{-3})(1)(10^{-3})(10^{-1}) = 10^{-8}, \quad -\log_2 p_{\text{coincidence}} = 26.6 \text{ bits.}$$

Give OPH and coincidence equal odds before reading the table, use the product as the likelihood of the evidence under coincidence, and apply a 10^4 search penalty. The simple Bayes score is

$$\Pr(\text{OPH incorrect} \mid E) \approx \frac{10^4 p_{\text{coincidence}}}{1 + 10^4 p_{\text{coincidence}}} = 0.00999\%,$$

about one chance in 10001. A calibrated posterior would require measured likelihoods and a defensible prior; this score keeps its assumptions visible.

References

Data and standards

- [S1] CODATA Task Group (2024). “2022 CODATA recommended values: Newtonian constant of gravitation.” NIST Reference on Constants.
- [S2] CODATA Task Group (2024). “2022 CODATA recommended values: inverse fine-structure constant.” NIST Reference on Constants.
- [S8] Particle Data Group (2026). “Leptons.” Review of Particle Physics summary table.

OPH papers All OPH series papers below use release **r1534**.

- [1/O1] Mueller et al. (2026). “Observers Are All You Need.”
- [2/O2] Müller et al. (2026). “Reality as a Consensus Protocol: The Fixed-Point Computation That Implements Physics.”
- [3/O3] Müller et al. (2026). “Federated Echosahedral Screen Microphysics: Patch Hardware, Records, and Observer Synchronization in OPH.”
- [4/O4] Müller et al. (2026). “Recovering Relativity and the Standard Model from Observer Overlap Consistency.”
- [5/O5] Müller et al. (2026). “Deriving the Particle Zoo from Observer Consistency.”

- [6/O6] Müller, B. (2026). “The Fine-Structure Constant as an OPH Pixel Fixed Point.” OPH technical note.
- [7/O7] Mueller, B., and D. Matscheko (2026). “Observer-Patch Holography and the Dark Matter Phenomenon.”
- [8/O8] Müller, B. (2026). “Explaining the Yang–Mills Mass Gap with Observer-Patch Repair Dynamics.” OPH technical note.
- [9/O9] Müller, B. (2026). “Observer-Patch Holography as a String-Vacuum Selector.” OPH technical note.

Executable artifacts

- [C1] FloatingPragma (2026). Pixel-closure solver and receipts.
- [C2] FloatingPragma (2026). Particle-branch source and audit suite.
- [C3] FloatingPragma (2026). Hierarchy theorem package and receipts.
- [C4] FloatingPragma (2026). Electroweak calibration receipts.
- [C5] FloatingPragma (2026). Consensus schedule benchmarks.
- [C6] FloatingPragma (2026). Lean 4/Mathlib theorem artifact.
- [C7] Mueller, B. (2026). OPH-FPE simulation and receipts, including the quantum-record, BW/KMS, H^3 , and geometry runs.
- [C8] FloatingPragma (2026). Lorentz–Einstein geometry receipts.
- [C9] FloatingPragma (2026). SPARC galaxy scorecard and receipts.