# A Finite Black-Hole Information Ledger for Observer-Patch Holography

The OPH collaboration

November 2, 2026

**Paper release:** `r2058` **Released:** November 2, 2026

**Collaboration affiliation:** Pragma Research Inc.

**Abstract**

On the layered source family of the spacetime construction, with specified population and read law, the number of unordered read pairs crossing a surface per layer obeys an area law with exact limiting coefficient $(\pi/4)n^2a^4$ per unit area for site density $n$ and read radius $a$, and declaring one nat of cut entropy per crossing read fixes the Planck-area dictionary $n^2a^4 = 1/(\pi\ell_P^2)$. This declared count is distinct from native regional entropy, which on the two-dimensional source screen carries no nonzero coefficient times boundary-seam count. Around it we formulate a finite information ledger for observer-patch models using machine-checked dynamics and record channels. A continuous one-parameter star-automorphism group on a finite-dimensional private algebra acts by blockwise unitary conjugation; for a supplied projective partition, the public record channel is the exact uniform mixture of independent sign unitaries; classical relative-entropy contraction and finite fluctuation identities supply complementary thermodynamic inputs; and a declared rich-fibre adapter constructs noncommutative regional algebras with commuting disjoint blocks and unique restriction gluing. The support-aware entropy identity for pinching is a known theorem that the library has not formalized; we state its proposed thermality and information-relocation consequences as finite conjectures. Together these results and conjectures make the black-hole information question precise within the supplied finite model.

**One theory and its executable hypothesis.** OPH proposes closure: one complete history is both a timeless self-referential structure and its own internal computation, and observers inside it recover and realize its specification. The hypothesis that makes this proposal concrete and testable is executable.

> *A simple, fully specified local repair and memory law on the proposed carrier architecture admits continuing observers whose compatible records and interventions have the observed* $3+1$*-dimensional quantum-field description and a common classical-gravity limit. The preferred matter mechanism is localized topological defects of native overlap transport, with rest energy derived from their excess maintenance dynamics in the same physical clock and energy dictionary.*

One source process must meet every part of this statement: records, clocks, matter, energy and responsive geometry come from the same law, state and refinement. Coherence, protected codes, radial directions, fields, clocks and couplings may be specified as part of its design; their implementation and observable consequences must be established on that process. A failed specified law rejects that law and its tested interpretation; it rejects the wider proposal only when the rejected choice is proved necessary.

**The tower.** The papers form one tower over the executable model. Each layer states a hypothesis and proves finite theorems under displayed premises. A conclusion passes to the next layer only after that premise is proved on the same model process; until then it is a conditional implication.

0. **Closure and agreement.** One history reads itself into existence, and agreement on shared records defines the public world. This is the interpretation and acceptance criterion; no theorem selects a unique universe.

1. **Repair mathematics.** Local agreement repair reaches global consistency, and nonzero cycle holonomy is the residue that local repair cannot remove. Fixed-point, normal-form, frustration and finite event-algebra theorems hold on their stated domains.

2. **Carrier and repair law.** Federated twelve-port icosahedral carriers with the native C/E seam repair are the substrate. The seam rule follows from four stated requirements, the quantum record laws are classified, and the uniform native repair selects a rank-three slow-response band.

3. **Executable observer.** Observers are continuing native processes that prepare, read, retain and act on records. Finite complete-output executions, overlapping-observer correspondence, event clocks and checkpoint continuation are verified; one continuing observer with every service and resource accounted is the open join.

4. **Clocks and fields.** Modular flow of observer algebras supplies candidate local clocks, and native repair supplies a positive continuum generator of spectral dimension two on the screen. A declared radial extension gives a conditional smooth $1+3$ scalar description. Complete finite source factors join spatial CAR dynamics, particle creation, repair and a common calibrated clock. Physical validation of that clock and energy normalization, and an interacting chiral $3+1$ limit, remain open.

5. **Gauge structure and matter.** Gauge structure is the transport of overlap data, and stable massive matter is a localized defect whose boundary transport carries a holonomy that admitted repairs cannot erase. The compact Lie type $\mathfrak{u}(1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(3)$ follows under complete-response premises, and fixed nonzero holonomy forces positive residual mismatch. A specified coherent ordering memory protects native exchange-history sectors, and a colored extension implements their fermionic exchange. A localized physical defect, its maintenance-to-mass law and a predicted measured spectrum remain unestablished; the earlier double-load candidate is transient.

6. **Gravity.** Einstein dynamics is the coarse response of the same process. The null-balance and tensor-completion implication is proved under its premises; realizing those premises on one refinement tower is open, and the static radial product fails null balance.

7. **Cosmology and tests.** The same process must fix the vacuum, dark sector, primordial response and black-hole records, and pass independent prospective tests. The cosmology papers and the fine-structure, Koide, de Sitter and Yang–Mills supplements carry conditional branches, diagnostics and retrospective comparisons at this layer.

**Completion and stopping.** The aim is a consistent and complete theory: every layer realized on one process, or a proved obstruction that removes OPH or a stated branch of it. Results that no join needs leave the active chain with their provenance. A universal obstruction over a frozen admissible class, an unavoidable excluded prediction, or a contradiction among indispensable premises stops that class. A failed candidate, an unconstructed map or a resource stop has a narrower meaning.

The ordered joins, retained foundations and stopping rules are recorded with the executable model in the public repository.[1]

**Fixed physical boundaries.** Repair counts, host runtime and entropy production are distinct from energy; the maintenance-to-energy map must be derived with one common signal speed and no per-species conversion. The modular parameter is dimensionless until a specified and implemented clock calibration relates it to duration. Stationary repair or record entropy is distinct from the fixed-volume generalized-entropy variation that the Einstein construction consumes. A massless photon makes the infinite-volume theory gapless, so sector gaps and thresholds name their state and limit; confinement precludes free massless gluons as observable particles; a quantum graviton lies outside the classical Einstein theorem.

## This paper's hypothesis and place in the tower

This paper sits at layer seven and develops the black-hole branch as a finite record ledger. Its hypothesis is that black-hole information is relocated, never destroyed: the ambient evolution of a private algebra is unitary, the thermal-looking public record is produced by an explicit unital channel, and the information missing from the public record is the relative entropy $D(\rho \,\|\, \mathcal{E}(\rho))$, carried by coherences that no observer access cut exposes.

Statements labeled *Theorem* are machine-checked in the Lean library at the cited modules. Statements labeled *Known theorem* are standard finite-dimensional results that the library has not formalized. Statements labeled *Conjecture* are unproved, and nothing here claims them. The crossing-read area law holds on a supplied layered population with a declared one-nat-per-read count. The phrase "black hole" names the shape of the question: the model has no continuum horizon, gravitational collapse, Hawking flux or evaporation endpoint, and no observable prediction follows from the finite ledger.

For the branch to apply, the executable model must supply on one process a faithful continuum realization of the finite source-causal carrier, a calibrated identification of native event clocks with proper duration [8], Einstein-branch geometry and stress, and a record channel derived from its native record laws; any observable additionally needs a prospectively fixed measurement contract under immutable custody. The native construction constrains this join. Its record-discarded repair channel attenuates coherence and is not unitary on the carrier, while each finite observation history becomes unitary once its record environment is kept (Section 3); its executed record law is not idempotent, so the ledger's pinching is not derived from it [12]; and its regional entropy carries no boundary-seam coefficient on the two-dimensional source screen [13], so the area law is a declared count rather than a native entropy law. Its continuum operator is a positive two-dimensional spatial operator with no radial domain or quantum vacuum [14, 15], and its causal statement is a bound of one seam per ordered repair event with no laboratory calibration [16]. The branch is refuted by an access cut of the specified net that exposes the off-block coherences, by a failure of the generalized-second-law-shaped monotone with a unital record channel, or by a proof that the native record laws admit no channel whose missing information is relocated rather than destroyed.

[1] https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/docs/COMPLETION_PROGRAM.md

# 1 The question in finite form

The black-hole information problem sets two commitments against each other: microscopic dynamics that preserves information, and a macroscopic record that looks exactly thermal. In the observer-patch model both commitments are formal objects, and the question becomes finite: given a unitary ambient dynamics on a finite private algebra, and given that what any observer can hold is a public record produced by a specific channel, account exactly for the information that the record does not carry.

The model supplies sharper versions of both sides than a generic quantum system would.

First, within the supplied pointwise-continuous star-automorphism class, unitarity is not an additional assumption. The Stone-converse packet proves that every pointwise-continuous one-parameter star-automorphism group of a unital finite-dimensional private star subalgebra is blockwise unitary conjugation generated by one self-adjoint Hamiltonian per block. The theorem makes no claim about dynamics outside that class.

Second, the record channel is not a generic decoherence map. For every projective partition, the publicization pinching is exactly the uniform average of the $2^k$ independently signed block reflections. The channel that produces the public record is therefore an explicit uniform mixture of unitaries, with the mixing unitaries named. It is unital; entropy nondecrease for this channel is part of Known theorem 3.1. This pinching is the publicization of a supplied partition. The native record laws of the executable observer form one convex family classified by five coherence multipliers, in which exactly two members are idempotent and the executed law C is not; the ledger's channel is not derived from that family [12].

The conjectured ledger combines these two facts with the exact thermodynamic theorems. It would identify the information absent from the public record with observer-inaccessible coherences of the ambient state through the formula in Known theorem 3.1. The executable observer computes this missing information exactly for its own records. Its complete classical tape and final load register recover every initial load population but not an arbitrary unknown quantum state; the missing information is coherence between different committed histories. The optimal entanglement fidelity over all decoders is $D^{-2}\sum_h w_h \operatorname{rank}(P_h)^2$, which equals 1301/2028 for the full-carrier two-event C observer and 3329/6084 for E [11].

# 2 Machine-checked inputs

Each input below is a machine-checked result in the repository, cited by module. The first two are standard finite-dimensional facts: the first is the Skolem–Noether theorem combined with Stone's theorem on each Wedderburn block, and the second is character orthogonality on the group of diagonal signs. They enter as checked inputs, not as new results. Their joint consequence and the precise conjectures are stated in Section 3.

**Theorem 2.1** (Finite Stone packet)**.** *Every continuous one-parameter group of star automorphisms of a unital, finite-dimensional private star subalgebra containing the ambient identity acts by blockwise unitary conjugation. On each Wedderburn block it is generated by a fixed self-adjoint Hamiltonian, unique up to a real scalar.*

**Theorem 2.2** (Publicization is a uniform mixture of sign unitaries)**.** *For every supplied $k$-block projective partition, the pinching map is the uniform average of the $2^k$ diagonal sign unitaries (Lean module `EventAlgebra.RecordMajorization` [18]); averaging only the two global signs does not pinch, so the independence of the signs is load-bearing.*

**Theorem 2.3** (Exact data processing and second law)**.** *Every stochastic kernel preserving a faithful stationary reference contracts Kullback–Leibler divergence to that reference (`kl_push_le`), the conditional heat bath obeys the second law (`heatBath_secondLaw`). For finite real-valued distributions and kernels, the integral fluctuation identity holds under its positivity, normalization and stationary-reference hypotheses; the pointwise Crooks identity additionally assumes detailed balance. The proofs are in the Lean modules `Thermodynamics.FiniteConditionalRepair` and `Thermodynamics.FluctuationTheorems` [18].*

**Theorem 2.4** (Support boundary for entropy architecture)**.** *The totalized continuous-functional-calculus logarithm sends every projection to zero, so the raw finite trace formula for relative entropy assigns the value zero to an orthogonal pair of density projections whose support inclusion fails (`EventAlgebra.SpectralEntropyBoundary` [18]). A support-aware extended-value layer is therefore mandatory, not stylistic.*

**Theorem 2.5** (Conditional rich-fibre regional packet)**.** *The locally pinned fresh-seed source run supplies split-fibre labels at four pairwise-disjoint observer windows. A declared finite adapter constructs a genuinely noncommutative block algebra with a nonunital two-by-two designated-fibre matrix corner at every observer region, elementwise commutation across distinct constructed blocks, computed four-window coverage of the declared top algebra, and unique restriction gluing with its reconstruction corollary (`QFT.RichFibreWitness`, `QFT.RichFibreRegionalNet` [18]). That source run does not produce the matrix units, state, regional restrictions or repair of this adapter, and the corner is neither a tensor factor nor a `TensorSplitReceipt`. A source-attached regional-factor theorem requires an operator algebra and a justified factor or local-channel adapter.*

**Theorem 2.6** (Finite KMS identity, degenerate inheritance boundary)**.** *The finite algebraic KMS identity holds for Gibbs states of support-graded flows, and the truncation-8 nets inherit KMS structure only degenerately (`QFT.StructuralInheritance` [18]). The nondegenerate continuation on the rich-fibre net is Conjecture 3.6.*

Because the native full configuration reference has a tracial limit at fixed modular parameter, its limiting modular automorphism is the identity, and it supplies no nondegenerate KMS state of this kind [15].

## 3 Finite conjectures

Throughout, $\rho$ is a finite density matrix on the ambient algebra, $\mathcal{E}$ the pinching of a supplied projective partition, $S$ the support-aware von Neumann entropy, and $D(\cdot \,\|\, \cdot)$ the support-aware relative entropy. The statements of this section add finite hypotheses to the preceding theorems and do not follow from them. Two statements of this section are known theorems that the library has not formalized; they are labeled accordingly.

**Known theorem 3.1** (Support-aware entropy core)**.** *For the pinching $\mathcal{E}$ of a projective partition,*

$$S(\mathcal{E}(\rho)) - S(\rho) = D(\rho \,\|\, \mathcal{E}(\rho)) \geq 0,$$

*with equality exactly at the public fixed states, where ($\mathcal{E}(\rho) = \rho$), and the sign-unitary mixture is entropy-nondecreasing. This core is the subset of the spectral-information chain required by the ledger; the more general majorization, maximum-entropy, and H-theorem scope is not used here.*

*Standard proof.* The pinching is the uniform average of $2^k$ sign conjugations, so $\mathcal{E}(\rho) \geq 2^{-k}\rho$ and the support of $\rho$ lies in that of $\mathcal{E}(\rho)$; the relative entropy is finite. The operator $\log \mathcal{E}(\rho)$ is block diagonal, and $\mathcal{E}$ is a trace-preserving conditional expectation onto block-diagonal operators, so $\operatorname{Tr} \rho \log \mathcal{E}(\rho) = \operatorname{Tr} \mathcal{E}(\rho) \log \mathcal{E}(\rho)$. Hence $S(\mathcal{E}(\rho)) - S(\rho) = \operatorname{Tr} \rho \log \rho - \operatorname{Tr} \rho \log \mathcal{E}(\rho) = D(\rho \,\|\, \mathcal{E}(\rho))$, which is nonnegative with equality exactly when $\rho = \mathcal{E}(\rho)$ by Klein's inequality. What remains open is the support-aware extended-value formalization in the library, which the support boundary theorem above makes mandatory. □

**Known theorem 3.2** (Finite no-loss statement)**.** *Let the specified ambient dynamics be a unitary dilation of repair that retains its record environment. The joint state of system and records evolves by the forced unitary conjugation of the Stone packet, and ordinary entropy of that joint state is preserved; when both arguments co-evolve under the same unitary, each explicitly named jointly unitarily invariant quantity is also preserved (including support-aware relative entropy and trace distance when those quantities are defined). The statement quantifies over the specified dynamics class; no toy evolution is substituted and no arbitrary coordinate-dependent divergence is covered.*

This is unitary invariance of the von Neumann entropy and of every jointly unitarily invariant quantity; its content for the ledger lies in the premise that the specified dynamics is a unitary dilation that retains its records.

The record-discarded native repair channel is not of this kind: it strictly attenuates a coherent mode, which rules out unitary conjugation on the carrier, and under independent seam attempts the unread binary fixed-number state converges to its tracial entropy maximum. Each finite native observation history has an exact unitary representation once its record environment is kept, with a dimensionless Hermitian action $A_0$ whose exponential $\exp(-iA_0)$ is that unitary representation [8, 9, 10].

**Conjecture 3.3** (Ledger identity and relocation)**.** *For the specified record channel, the public entropy gain equals $D(\rho \,\|\, \mathcal{E}(\rho))$; the quantity is nonnegative, vanishes exactly at the public fixed states, and is carried entirely by the off-block coherences that no access cut of the specified net exposes. Information is relocated, never destroyed, and the relocation is exactly quantified.*

The equality and its sign are Known theorem 3.1; the open part is the statement that no access cut of the specified net exposes the off-block coherences.

**Conjecture 3.4** (Generalized-second-law-shaped monotone)**.** *The composition of the cap first law with Kullback–Leibler contraction yields one exact monotone combining record entropy and the capacity ledger under repair, with finite constants and a negative control exhibiting failure of the monotone when unitality of the record channel is dropped.*

A classical sector-level instance follows from the joint counting identity $S_{\mathrm{count}}(p, d) = H(p) + \sum_i p_i \log d_i = \log M - D(p \,\|\, d/M)$: a repair kernel on sectors that preserves $d/M$ contracts the divergence by `kl_push_le`, so $S_{\mathrm{count}}$ does not decrease, while a reset kernel with stationary law $(1/4, 3/4)$ on two equal sectors does not preserve $d/M$ and lowers $S_{\mathrm{count}}$ from its maximum [13]. The conjecture concerns the quantum record channel and the cap first law.

**Conjecture 3.5** (Complementarity receipts)**.** *There is a composition of the generic operational access-cut interface, the supplied rich-fibre data, and a justified operator/local-channel adapter that forms a two-observer ledger in which each accessible algebra determines its own record entropy, the declared block-commutation theorem supplies an algebraic locality receipt, and a glued top observable is uniquely reconstructed within the declared restriction system. No physical no-signalling or refinement-natural interpretation follows before that attachment.*

**Conjecture 3.6** (Nondegenerate finite KMS receipt)**.** *The modular data of a restricted state on one genuinely noncommutative rich-fibre region satisfies the finite KMS identity at a capacity-linked inverse temperature, nondegenerately. This receipt supplies a finite KMS input but does not establish structural QFT thermality.*

**Conjecture 3.7** (Finite Page-crossover inequality)**.** *For a bipartition of the specified finite net, the exact $2^k$ sign-unitary average obeys a finite inequality that locates the crossover of accessible-record information without Haar integration.*

*Remark* 3.8 (Conditional finite implication). Known theorems 3.1 and 3.2 with Conjectures 3.3–3.6 would imply simultaneous ambient unitarity and a thermal public record, with the inaccessible information equal to the computable functional $D(\rho \,\|\, \mathcal{E}(\rho))$ and localized in observer-inaccessible coherences. The implication supplies no physical black-hole attachment. A source-selected order embedding with calibrated density (a causal-set faithful embedding), manifoldlikeness, and a calibrated identification of native event clocks with proper time are additional hypotheses. Any smooth Einstein reading also requires same-family tensor-curvature convergence or the independent continuum small-ball/null-balance identification.

## 4 Crossing-read area law on the source family

The layered read law of the conservative source family supplies a finite count that scales with area. Fix a population of constant density $n$, sites per unit volume, in a bounded convex window $\Omega \subset \mathbb{R}^3$, a read radius $a$, and the complete-neighbour read law: an event at layer $j+1$ reads every event at layer $j$ within distance $a$, the same site included. For a plane section $\Sigma = \Omega \cap \{x_1 = P\}$, a crossing read of one layer is an unordered site pair $\{s, t\}$ with $s \neq t$, $\|t - s\| \leq a$, and $s_1 < P < t_1$. Each such pair is realized in every layer by one read in each direction across $\Sigma$, from the lower layer to the upper layer; the ordered read count is therefore twice the unordered count, and the unordered count per layer is the primary quantity here. A same-site read never crosses a plane, and a site with $s_1 = P$ belongs to neither side.

**Proposition 4.1** (Crossing reads per unit area)**.** *Let $S_k \subset \Omega$ be finite populations with a partition of $\Omega$ into cells of common volume $1/n_k$, each cell contained in the ball of radius $H_k$ about its site, with read radii $a_k \to 0$ and $H_k/a_k \to 0$. Let $\Sigma = \Omega \cap \{x_1 = P\}$ have positive area $|\Sigma|$. The number $X_k$ of unordered crossing reads of $\Sigma$ per layer satisfies*

$$X_k = n_k^2 \,|\Sigma| \int_0^{a_k} \frac{\pi (a_k - u)^2 (2a_k + u)}{3} \,\mathrm{d}u \,(1 + o(1)) = \frac{\pi}{4}\, n_k^2 a_k^4 \,|\Sigma| \,(1 + o(1)). \tag{1}$$

*The relative error is bounded by a constant multiple of $H_k/a_k + a_k$, the constant depending on $\Omega$ and $\Sigma$.*

*Proof.* Write $u(s) = P - s_1$ for the depth of a site below $\Sigma$ and $C(s) = \{t \in \Omega :\ t_1 > P,\ \|t - s\| \leq a_k\}$, so that $X_k = \sum_{0 < u(s) \leq a_k} |S_k \cap C(s)|$. The cell of a site $t$ lies in $B(t, H_k)$; a cell that meets both $C(s)$ and its complement lies in the $2H_k$-collar of $\partial C(s)$. Hence $|S_k \cap C(s)|$ differs from $n_k \operatorname{vol} C(s)$ by at most $n_k$ times the volume of that collar. The boundary $\partial C(s)$ consists of a spherical cap, a planar disc and pieces of $\partial\Omega$, of total area $O(a_k^2)$, so the error is $O(n_k H_k a_k^2)$ per site. The slab $0 < u(s) \leq a_k$ contains $O(n_k a_k |\Sigma|)$ sites, which gives a total error $O(n_k^2 H_k a_k^3 |\Sigma|)$, of relative order $H_k/a_k$.

The function $f(s) = n_k \operatorname{vol} C(s)$ on the slab is bounded by $\frac{2\pi}{3} n_k a_k^3$ and is Lipschitz with constant $O(n_k a_k^2)$: its depth derivative is $-\pi n_k (a_k^2 - u^2)$ and a lateral displacement changes $C(s)$ through

the faces of $\Omega$ by a set of measure at most $\pi a_k^2$ per unit displacement. The same cell argument, applied to the slab, bounds $|\sum_{0<u(s)\le a_k} f(s) - n_k \int_{0<u\le a_k} f \,\mathrm{d}^3 s|$ by $n_k$ times the Lipschitz constant times $H_k$ times the slab volume, plus the contribution of the sites within $H_k$ of the two slab faces; both are $O(n_k^2 H_k a_k^3 |\Sigma|)$, again of relative order $H_k/a_k$.

Finally, for $s$ at lateral distance more than $a_k$ from the relative boundary of $\Sigma$ in the plane, $C(s)$ is the untruncated spherical cap of $B(s, a_k)$ beyond height $u$, with volume

$$\int_u^{a_k} \pi(a_k^2 - z^2)\,\mathrm{d}z = \frac{\pi(a_k - u)^2(2a_k + u)}{3}.$$

The truncated sites occupy a lateral collar of area $O(a_k)$ relative to $|\Sigma|$, so the integral over the slab equals $|\Sigma| \int_0^{a_k} \pi(a_k - u)^2(2a_k + u)/3\,\mathrm{d}u$ up to a relative error $O(a_k)$. The depth integral is $\pi a_k^4/4$. □

**Corollary 4.2** (Golden-family value)**.** *For the golden source population $S_q$ of $q^3$ sites in $\Omega = [0, L]^3$, $L = 2/\sqrt{\varphi + 2}$, with $n_q = q^3/L^3$, $a_q = L/\sqrt{q}$ and cell assignment error $H_q = 2\sqrt{3}L/q$, one has $n_q^2 a_q^4 = q^4/L^2$, and the unordered crossing count per layer through a full cross-section $\Sigma = \Omega \cap \{x_i = P\}$ is*

$$X_q = \frac{\pi q^4}{4}(1 + O(q^{-1/2})). \tag{2}$$

*For the mid-plane $P = L/2$ the continuum count with the caps truncated by the cube is*

$$n_q^2 \int_{s\in\Omega,\, s_1<L/2} \mathrm{vol}\{t \in \Omega :\ t_1 > L/2,\ \|t - s\| \le a_q\}\,\mathrm{d}^3 s = q^4 \left(\frac{\pi}{4} - \frac{8}{15\sqrt{q}} + \frac{1}{12\,q}\right). \tag{3}$$

*Proof.* Here $H_q/a_q = 2\sqrt{3}\,q^{-1/2}$ and $a_q/L = q^{-1/2}$, so Proposition 4.1 gives (2). For (3) substitute $d = t - s$. For $0 < d_1 \le a_q \le L/2$ and $|d_2|, |d_3| \le L$, the set of $s \in \Omega$ with $s + d \in \Omega$ and $s_1 < L/2 < s_1 + d_1$ has volume $d_1(L - |d_2|)(L - |d_3|)$. Over the half ball $d_1 > 0$, $\|d\| \le a_q$, the moments are $\int d_1 = \pi a_q^4/4$, $\int d_1|d_2| = 4a_q^5/15$ and $\int d_1|d_2||d_3| = a_q^6/12$. Hence the count equals $n_q^2 a_q^4(\pi L^2/4 - 8La_q/15 + a_q^2/12)$, which is (3) after $n_q^2 a_q^4 = q^4/L^2$ and $a_q = L/\sqrt{q}$. □

**Proposition 4.3** (Area law and the Planck-area dictionary)**.** *Declare that each crossing read carries one nat of cut entropy across $\Sigma$ per layer. In the limit of Proposition 4.1 the cut entropy is the area law*

$$S_{\mathrm{cut}}(\Sigma) = \frac{\pi}{4}\, n^2 a^4\, |\Sigma|. \tag{4}$$

*Identifying (4) with $|\Sigma|/(4\ell_P^2)$ fixes*

$$n^2 a^4 = \frac{1}{\pi \ell_P^2}, \qquad \ell_P = \frac{1}{\sqrt{\pi}\, n\, a^2}, \tag{5}$$

*and on the golden family $\ell_P = L/(\sqrt{\pi}\, q^2)$.*

*Proof.* Equation (4) is (1) with one nat per read. Equating $(\pi/4)n^2a^4$ with $1/(4\ell_P^2)$ and solving for $\ell_P^2$ gives (5); with $n^2a^4 = q^4/L^2$ this reads $\ell_P^2 = L^2/(\pi q^4)$. □

*Remark* 4.4 (What is not claimed). Equation (5) is a calibration of the read scale by the area coefficient, in the same way that the layer duration $a/c$ is a calibration of the clock. No horizon is constructed: $\Sigma$ is a plane section of the source window, with the same read law on both sides. The nat-per-read reading is a declared dictionary, without a derivation from the record channel of the finite ledger. No temperature, no Hawking flux and no physical value of $\ell_P$ follow; $L$ is the side

of the source window, and the family fixes only the ratio $\ell_P/L = 1/(\sqrt{\pi}\,q^2)$. Dou and Sorkin [7] count causal links across a horizon in a Poisson-sprinkled causal set and obtain an area law whose coefficient depends on the sprinkling; here the population is deterministic, the count is exact at every finite $q$, and the coefficient $\pi/4$ in (1) is the exact limit.

The depth integral in (1), the value $(\pi/4)n^2a^4$, the golden identity $n_q^2 a_q^4 = q^4/L^2$ with the cross-section value $\pi q^4/4$, and the dictionary (5) with its golden value $\ell_P^2 = L^2/(\pi q^4)$ are machine-checked in the Lean module `CrossingReadAreaLaw` (`capVolume_integral`, `crossingDensity_eq`, `golden_crossing_count`, `planck_area_of_area_law`, `golden_planck_length_sq`) with the standard axioms only. The limit statement of Proposition 4.1 is analytic and is proved above. The crossing-read count is a declared one-nat-per-read quantity on a supplied three-dimensional population. On the two-dimensional native source screens, regional reduced entropy carries no nonzero coefficient times the boundary-seam count, so this law is distinct from a native entropy-area law; a whole-screen reading as the boundary of a three-dimensional region needs its own region map.

**Finite receipt.** The exact crossing counts of the golden family at $q = 5, 8, 13, 21, 34, 55$ are recorded with an independent verifier. The population and the neighbour relation are rebuilt from the definitions, the relation agrees edge for edge with the causal-poset evidence package at every level, every side decision is exact in $\mathbb{Q}(\varphi)$, and no site lies on any of the five planes $x_1 = L/2$, $x_1 = L/4$, $x_1 = 3L/4$, $x_2 = L/2$, $x_3 = L/2$. Table 1 lists the mid-plane counts. The counts across $x_2 = L/2$ and $x_3 = L/2$ equal the $x_1 = L/2$ count at every level, as the product population is symmetric under coordinate permutations. At $q = 34$ the mid-plane count is 907 080 against the box-corrected value 930 602.0, ratio 0.975, and against the infinite-plane value 1 049 555.8, ratio 0.864; at $q = 55$ the count is 6 496 004 against 6 542 684.6, ratio 0.993, and against 7 186 884.1, ratio 0.904. The ratio to the box-corrected value is closer to one than the ratio to the infinite-plane value at every level; it lies within 0.13 of one at every level and within 0.026 of one for $q \geq 21$, while the ratio to the infinite-plane value rises from 0.658 to 0.904. The residual at fixed $q$ is the finite population itself: the two sides of the mid-plane hold 84 700 and 81 675 sites at $q = 55$, and the mean number of neighbours of a site, 1459.3 at $q = 55$, compares with the cube mean 1461.1 of $n_q \,\mathrm{vol}(B(s, a_q) \cap \Omega)$ and with the untruncated ball value 1708.6.

| $q$ | count | $\pi q^4/4$ | ratio | box-corrected | ratio |
|---|---|---|---|---|---|
| 5 | 323 | 490.9 | 0.658 | 352.2 | 0.917 |
| 8 | 2 164 | 3 217.0 | 0.673 | 2 487.3 | 0.870 |
| 13 | 17 754 | 22 431.8 | 0.791 | 18 390.1 | 0.965 |
| 21 | 129 830 | 152 745.0 | 0.850 | 130 882.5 | 0.992 |
| 34 | 907 080 | 1 049 555.8 | 0.864 | 930 602.0 | 0.975 |
| 55 | 6 496 004 | 7 186 884.1 | 0.904 | 6 542 684.6 | 0.993 |

Table 1: Unordered crossing reads per layer across the mid-plane $x_1 = L/2$ of the golden family, the infinite-plane value $\pi q^4/4$, and the box-corrected value (3), with the ratios of the count to each.

**Declared cut entropy and native regional entropy.** The population, its read radius and the layer read law of the preceding section are specified inputs, and the cut entropy is a declared count on read links; the seam-count wording of that section refers to these read links, not to native repair seams. It is distinct from native regional entropy. On a source screen of $N = 240 \cdot 4^l$ ports, a complete refining base-face patch with $12 \cdot 4^l$ ports has $3 \cdot 2^l$ crossing seams; its stationary

reduced entropy grows in proportion to $4^l$, while the return reference's mutual information is at most $\log(12 \cdot 4^l + 1) + 2025/(64N)$, which per crossing seam tends to zero. Neither native regional quantity carries a nonzero universal coefficient times boundary-seam count on that two-dimensional source [13]. A reading of a whole screen's cell count as the area of a three-dimensional region's boundary is not excluded, and that region map is a premise. The Lean statements of the preceding section are in the module `Geometry.CrossingReadAreaLaw` [18].

## 5 Relation to standard approaches

The Page analysis, the Hayden–Preskill protocol, and island computations concern continuum or holographic systems with gravitational dynamics. The finite ledger formulates their information question using separately proved unitary dynamics and a specified record channel. Its proposed entropy and thermality composition is conditional on Known theorems 3.1 and 3.2 with Conjectures 3.3–3.6. Within the supplied automorphism class, the Stone converse gives unitary implementation; the public channel is the explicit sign-unitary mixture. A physical black-hole interpretation additionally requires the spacetime, horizon and stress attachments stated above.

## 6 Observables discipline

The finite ledger implies no observable black-hole signature. A proposed observable claim derived from the ledger, such as saturation-regime relaxation structure or record-capacity signatures in compact-object transients, requires a prospectively fixed clock, frame, scale, and readout contract under immutable custody before any data comparison.

A conditional instrument template is defined for the integer-$k$ ringdown transition comb. It is not preregistered or frozen and therefore supports no observable claim; no ringdown value is frozen in the prediction register [17]. The transition rule is an imported continuation premise, not a source-derived consequence of the finite ledger. For an emission it assumes positive integer record dimensions satisfying $d_{\text{before}} = k d_{\text{after}}$, so $k$ must divide the initial dimension. The signed black-hole entropy change is $\ln d_{\text{after}} - \ln d_{\text{before}} = -\ln k$; the positive entropy loss entering the emitted-energy formula is $\ln k$. For positive remnant mass and subextremal spin $|\chi| < 1$, conditional on that rule, candidate features above the rotation line sit at $x_k = \ln k/(8\pi)$, with $x = (GM/(c^3 g(\chi)))(\omega - m\Omega_H)$ and $g(\chi) = 2\sqrt{1-\chi^2}/(1+\sqrt{1-\chi^2})$. Their offset-subtracted ratios equal $\ln k_a / \ln k_b$ under a common scale and offset for a denominator tooth $k_b \geq 2$, whose offset-subtracted value is proved nonzero. At $|\chi| = 1$, $g(\chi) = 0$: the candidate teeth collapse to the rotation line, and the rescaled coordinate and these ratios are undefined. Observed frequencies must use the detector-frame remnant mass $M_{\text{det}} = (1+z)M_{\text{source}}$ in both terms; the ratio is redshift-free only with this consistent frame convention. The tooth relation uses the leading small-transition Kerr first law; finite-step background corrections are not derived by this template.

The declared linewidth-to-spacing ratio is $64\pi^2 p_0/(a g(\chi)^2 \ln k)$, which is mass-independent but spin-dependent; a constant Page coefficient is not controlled near extremal spin. The declared $(k-1)/k$ KMS factor is a within-line net-response factor, not a cross-$k$ transition probability or prior. The algebraic invariance, divisibility and entropy-sign boundary, ladder bracketings, producer, independent byte-identical verifier, and tests are target-blind. No event likelihood has been evaluated. Published posterior samples alone are neither a strain likelihood nor a model evidence. This template is not anchorable until a source-derived transition, common detector likelihood or sufficient likelihood product, detector-frame nuisance model, normalized integer prior or explicit finite submodel, event selection, trials, greybody normalization, no-comb reference, and

convergence rule are fixed. The computed $2 \le k \le 12$ ladder is only a finite candidate submodel; excluding it would not reject the unrestricted integer family.

## 7 Verification surface

The proved theorems cited in Section 2 carry `#print axioms` receipts with only the standard axioms, no admissions, and no native-evaluation shortcut, in the modules named inline. The support-aware entropy statements use the general spectral-information definitions. Known theorems 3.1 and 3.2 with Conjectures 3.3–3.6 are not proved by those receipts.

## Credited prior work

The results of this paper build on the following established work. Each entry names an established result as it is used here and cites its sources.

- Bekenstein–Mukhanov area quantization, Kerr ringdown spectroscopy, and quasinormal-mode templates [19, 21, 20].
- Dou and Sorkin (2003), black-hole entropy as causal links crossing the horizon of a Poisson-sprinkled causal set [7].

## References


[1] S. W. Hawking, *Particle creation by black holes*, Commun. Math. Phys. 43 (1975) 199–220.

[2] S. W. Hawking, *Breakdown of predictability in gravitational collapse*, Phys. Rev. D 14 (1976) 2460–2473.

[3] D. N. Page, *Average entropy of a subsystem*, Phys. Rev. Lett. 71 (1993) 1291–1294.

[4] P. Hayden and J. Preskill, *Black holes as mirrors: quantum information in random subsystems*, JHEP 09 (2007) 120.

[5] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian, and A. Tajdini, *The entropy of Hawking radiation*, Rev. Mod. Phys. 93 (2021) 035002.

[6] G. Lindblad, *Completely positive maps and entropy inequalities*, Commun. Math. Phys. 40 (1975) 147–151.

[7] D. Dou and R. D. Sorkin, *Black-hole entropy as causal links*, Found. Phys. 33 (2003) 279–296.

[8] FloatingPragma (2026). The runnable OPH model: observer implementation, tests, receipts and construction guide. https://github.com/FloatingPragma/observer-patch-holography/tree/r2058/model.

[9] FloatingPragma (2026). A local action for native quantum observations. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/model/native_action/README.md.

[10] FloatingPragma (2026). Native quantum mixing and the full-carrier memory boundary. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/model/observer/QUANTUM_MIXING.md.

[11] FloatingPragma (2026). What complete native records can recover. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/model/observer/RECORD_RECOVERY.md.

[12] FloatingPragma (2026). Admissible record laws for native repair. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/model/record_laws/README.md.

[13] FloatingPragma (2026). Tangent first laws and finite entropy changes: native reference entropy and region boundaries. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/code/geometry/ENTROPY_FIRST_LAW_AUDIT.md.

[14] FloatingPragma (2026). A spatial fluctuation field from one native history: the common spatial generator, its locality and spectral dimension, and the boundary-trace test. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/model/physical_space_time/OCCUPATION_FIELD.md.

[15] FloatingPragma (2026). Joint time and field laws on matched source screens. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/model/observer_time_fields/REFINEMENT.md.

[16] FloatingPragma (2026). Exact native causal cones. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/model/common_source/CAUSALITY.md.

[17] FloatingPragma (2026). Frozen prediction ladder. https://github.com/FloatingPragma/observer-patch-holography/blob/r2058/docs/FROZEN_PREDICTION_LADDER.md.

[18] FloatingPragma (2026). Observer Patch Holography Lean formalization. https://github.com/FloatingPragma/observer-patch-holography/tree/r2058/Lean.

[19] Bekenstein, J. D., and Mukhanov, V. F. (1995). "Spectroscopy of the quantum black hole." *Physics Letters B* **360**, 7–12. https://doi.org/10.1016/0370-2693(95)01148-J

[20] Berti, E., Cardoso, V., and Starinets, A. O. (2009). "Quasinormal modes of black holes and black branes." *Classical and Quantum Gravity* **26**, 163001. https://doi.org/10.1088/0264-9381/26/16/163001

[21] Dreyer, O., Kelly, B., Krishnan, B., Finn, L. S., Garrison, D., and Lopez-Aleman, R. (2004). "Black-hole spectroscopy: testing general relativity through gravitational-wave observations." *Classical and Quantum Gravity* **21**, 787–803. https://doi.org/10.1088/0264-9381/21/4/003

## AI Assistance Disclosure

This research project used research-grade commercial models, including Anthropic's Fable and OpenAI's GPT-6 Astra, for research support, software development, editing, and synthesis. The authors are responsible for the paper's claims, methods, and final text.