

OPH Cosmology as a Finite-Source Prediction Program: Observer-Screen Synchronization, Dark Stress, and the Physical CMB Gate

B. Müller

June 24, 2026

Paper release: r1502 **Released:** June 24, 2026

Abstract

OPH cosmology asks whether finite observer-screen synchronization can do the jobs usually assigned to inflation and dark initial conditions. The program ties flatness to spatial holonomy, horizon coherence to same-boundary repair, near scale invariance to the screen spectrum, hot initial data to MaxEnt release, and acoustic peaks to ordinary Boltzmann transfer from finite OPH source artifacts. A 256k-patch simulator run gives a strong diagnostic CMB curve, with shape correlation 0.9951542364, normalized RMSE 0.0984455548, and zero mean peak-location error against the local binned TT table. The physical claim remains conditional. A CMB prediction needs finite source provenance, physical scale calibration, a covariant collar-packet stress parent, physical $\rho_A(a)$, $B_A(k, a)$, and $\Gamma_{\text{rec}}(k, a)$ kernels, Boltzmann transfer, and frozen likelihood receipts.

What This Paper Contributes

The OPH core papers recover local observer-facing geometry, records, modular flow, gauge/matter structure, and finite screen microphysics. The cosmology work asks a different question: which large-scale data can be predicted once the finite observer-screen state is released into an Einstein-Boltzmann cosmology?

The program paper has four jobs.

1. It consolidates the inflation-free OPH cosmology branch into the core repository’s cosmology directory, while keeping it outside the release build until the theorem gates close.
2. It separates diagnostic simulator outputs from physical predictions. A good visual or measurement-facing CMB overlay is not a physical CMB prediction unless its inputs are source-only finite OPH artifacts fixed before likelihood comparison.
3. It names the finite objects required for physical promotion of simulation outputs in CMB, vacuum, or cosmological uses of the dark sector: source provenance, physical mode calibration, freezeout, covariant stress, anomaly kernels, transfer, likelihood, and quotient ensemble receipts. The dark-sector derivations themselves are owned by the released cosmology paper `cosmology/oph_dark_matter_paper.tex`.
4. It gives the staged cosmology branch one theorem surface for the finite-source and likelihood claim gates.

Contents

1	Claim Boundary	3
2	Finite Quotient Ensembles and Claim Tiers	3
3	Finite Screen Spectrum Theorem Package	8
4	Physical Scale Bridge and Mode Calibration	11
5	Compatibility With the Core Stack	15
6	Source Corpus	16
7	Imported Theorem Map	17
8	Extended Cosmology Agenda	18
9	Finite Cosmological State	19
10	Inflation-Free Branch	20
11	Physical CMB Finite-Source Contract	21
12	Physical Scale Bridge	23
13	Dark/Anomaly Interface Imported from the Dark-Matter Paper	23
14	Physical Kernel Interface	25
15	Boltzmann Transfer and Frozen Likelihood	26
16	Vacuum and Quantum-Fluctuation Boundary	28
17	Claim-Gate Summary	28
18	Physical Promotion Rule	28
19	Conclusion	29

1 Claim Boundary

This is a continuation paper, not a release checkpoint and not a published OPH paper bundle member. It is part of the public core repository as the shared cosmology contract for paper and simulator work.

The branch claim is:

OPH can replace the logical jobs of inflation on a finite-source branch:
 flatness by direct/CMH spatial-holonomy selection, horizon coherence by same-boundary repair,
 near scale invariance by the geometric screen spectrum, hot initial data by MaxEnt release,
 and acoustic peaks by Boltzmann transfer from source-only OPH artifacts.

Here “can” means “can on the declared Phase III finite-source continuation branch if the listed source, stress, lift, transfer, and likelihood gates close.” It does not mean that inflation replacement, CMB spectra, or dark-sector cosmology are part of the recovered Phase I SM/GR core.

The branch has no physical CMB prediction, completed Planck/ACT likelihood run, OPH-native quantum vacuum, full cosmological dark-matter likelihood, or production particle-formation simulation. The 256k simulator run is measurement-facing diagnostic evidence; its promotion receipts keep the physical claim closed.

Remark 1.1 (No simulator shortcut). The paper-side theorem gates are not bureaucratic checks. They are the distinction between a simulation that happens to resemble CMB data and a prediction whose input artifacts were fixed without using CMB data. The simulator should not resume physical CMB promotion until those gates are mathematically specified here and then implemented as deterministic receipts.

2 Finite Quotient Ensembles and Claim Tiers

Finite quotient ensemble theorem surface. Fix a finite regulator r . The physical presentation space is Σ_r , the presentation redundancy groupoid is Γ_r , and the finite physical quotient is

$$Q_r = \Sigma_r / \Gamma_r, \quad \pi_r : \Sigma_r \rightarrow Q_r.$$

The quotient removes nonphysical presentation data: gauge representatives, port relabelings, mesh labels, shard or worker identifiers, queue order, repair schedule identifiers, retry counters, timestamps unless declared semantic, hidden carrier coordinates, and inert ancillary labels. If only settled configurations carry probability, the probability space is the normal-form subset

$$N_r = n_r(Q_r).$$

The map n_r is a normal-form map, not a probability law.

Observable algebras and reference states. In the finite classical case the quotient observable algebra is

$$\mathcal{O}_r = \ell^\infty(Q_r).$$

In the finite quantum case the physical algebra is a declared quotient algebra $\mathcal{A}_r^{\text{phys}}$ with state

$$\omega_r(A) = \text{Tr}(\rho_r A).$$

When the reference object is obtained from a finite lifted carrier, the load-bearing data are not an abstract groupoid cardinality alone. They are a tracially pointed quotient

$$\left(\mathcal{A}_{r,b}^{\text{phys}}, \tau_{r,b}^0\right), \quad \mathcal{A}_{r,b}^{\text{phys}} = z_{r,b} B(\tilde{\mathcal{H}}_r)^{G_r} z_{r,b},$$

where $U_r : G_r \rightarrow U(\tilde{\mathcal{H}}_r)$ is the compact gauge action and $z_{r,b}$ is the central projection for the declared boundary or superselection sector. The reference trace is

$$\tau_{r,b}^0(A) = \frac{\text{Tr}_{\tilde{\mathcal{H}}_r}(A)}{\text{Tr}_{\tilde{\mathcal{H}}_r}(z_{r,b})}.$$

If

$$z_{r,b} \tilde{\mathcal{H}}_r \cong \bigoplus_{\alpha} V_{\alpha} \otimes M_{\alpha},$$

with $d_{\alpha} = \dim V_{\alpha}$ and $m_{\alpha} = \dim M_{\alpha}$, then

$$\mathcal{A}_{r,b}^{\text{phys}} \cong \bigoplus_{\alpha} I_{V_{\alpha}} \otimes B(M_{\alpha}), \quad p_{r,\alpha} = \frac{d_{\alpha} m_{\alpha}}{\sum_{\beta} d_{\beta} m_{\beta}}.$$

These are the induced central-sector weights only after the carrier representation and boundary sector have been fixed.

OPH quotient ensemble. An OPH quotient ensemble is specified by a quotient-intrinsic base weight and action

$$m_r : Q_r \rightarrow \mathbb{R}_{>0}, \quad S_r : Q_r \rightarrow \mathbb{R} \cup \{+\infty\},$$

and

$$w_r(q) = m_r(q) e^{-S_r(q)}, \quad Z_r = \sum_{q \in Q_r} w_r(q), \quad \mu_r(q) = Z_r^{-1} w_r(q).$$

Equivalently, one may state an intrinsic projective prior ν_r on Q_r and set $\mu_r = (n_r)_{\#} \nu_r$. Uniform quotient counting, uniform representative counting pushed to the quotient, groupoid weights, and tracial central-sector weights are different physical claims. The paper must declare which one is being used.

Normal-form projector non-selection. For any map $N : Q \rightarrow Q_{\text{nf}}$, the induced map on laws

$$\mathcal{C}_Q(\mu) = N_{\#} \mu$$

is idempotent:

$$\mathcal{C}_Q^2 = \mathcal{C}_Q.$$

Every law supported on Q_{nf} is fixed. Therefore settlement or canonicalization never selects a unique physical probability law by itself.

Finite MaxEnt quotient ensemble. For finite Q , positive m , and quotient observables $F_1, \dots, F_k$, maximizing

$$\mathcal{H}_m(\nu) = - \sum_q \nu(q) \log \frac{\nu(q)}{m(q)}$$

subject to

$$\sum_q \nu(q) = 1, \quad \sum_q \nu(q) F_a(q) = c_a$$

has the full-support solution, when the feasible full-support surface is nonempty,

$$\mu(q) = \frac{m(q) \exp[-\sum_a \theta_a F_a(q)]}{Z(\theta)}.$$

Boundary optima obey the same formula after restricting to their support. On a finite noncommutative quotient algebra with faithful reference state σ_r ,

$$\rho_r = \frac{\exp(\log \sigma_r - \sum_a \theta_a F_{r,a})}{\text{Tr} \exp(\log \sigma_r - \sum_a \theta_a F_{r,a})}.$$

The finite constraint ledger must name every $F_{r,a}$, its units and support, the target expectation and source, sector or zero-mode treatment, refinement transformation, and proof that no run output or observational output entered the source definition.

Refinement compatibility and RG closure. For $s \succeq r$, let $c_{sr} : Q_s \rightarrow Q_r$ be the physical coarse map. Exact compatibility of weighted ensembles is equivalent to the fiber-sum identity

$$\sum_{q' : c_{sr}(q')=q} m_s(q') e^{-S_s(q')} = \alpha_{sr} m_r(q) e^{-S_r(q)}$$

for a constant $\alpha_{sr} > 0$ independent of q . Then

$$(c_{sr})_{\#} \mu_s = \mu_r.$$

If the one-step defects are

$$\delta_{k+1,k} = \|(c_{k+1,k})_{\#} \mu_{k+1} - \mu_k\|_{\text{TV}},$$

then

$$\|(c_{nr})_{\#} \mu_n - \mu_r\|_{\text{TV}} \leq \sum_{k=r}^{n-1} \delta_{k+1,k}.$$

For exponential-family refinement, exact closure requires the fine conditional free energy

$$G_{sr,\theta}(q_r) = -\log \mathbb{E}_{m_s^0} [\exp[-\theta \cdot F_s(Q_s)] \mid c_{sr}(Q_s) = q_r]$$

to equal $\kappa_{sr}(\theta) + R_{sr}(\theta) \cdot F_r(q_r)$. If the residual is uniformly bounded by ε , the induced total-variation defect is bounded by $\tanh \varepsilon$.

Implementation invariance and representative lifting. If implementations A, B have quotient bijections $h_r : Q_r^A \rightarrow Q_r^B$ satisfying

$$m_r^B(h_r q) = m_r^A(q), \quad S_r^B(h_r q) = S_r^A(q), \quad h_r \circ c_{sr}^A = c_{sr}^B \circ h_s,$$

then

$$(h_r)_{\#} \mu_r^A = \mu_r^B.$$

For tracially pointed quantum quotients the corresponding equivalence is a trace-preserving quotient equivalence. It is invariant under unitary intertwiners preserving the gauge action and sector, and under inert trivial ancillas $A \mapsto A \otimes I_{\text{anc}}$. It is not invariant under arbitrary changes of gauge-representation multiplicities.

If an implementation stores representatives, a representative-level law must be a conditional lift

$$\tilde{\mu}_r(x) = \mu_r(\pi_r x) \kappa_r(x \mid \pi_r x), \quad \sum_{x : \pi_r(x)=q} \kappa_r(x \mid q) = 1.$$

Then $(\pi_r)_{\#} \tilde{\mu}_r = \mu_r$. Uniform representative sampling yields orbit-size weights and is physical only if representative counting is the declared base measure.

Quotient-lumpable kernels and sampler correctness. A representative kernel $\tilde{P}(x, y)$ descends to Q_r only when

$$P_Q(q, q') = \sum_{y: \pi(y)=q'} \tilde{P}(x, y)$$

is independent of the chosen representative $x \in \pi^{-1}(q)$. For $w(q) = m(q)e^{-S(q)}$ and proposal $R(q, q')$ with reciprocal support, the Metropolis–Hastings acceptance rule

$$a(q, q') = \min \left\{ 1, \frac{w(q')R(q', q)}{w(q)R(q, q')} \right\}$$

gives detailed balance

$$\mu(q)R(q, q')a(q, q') = \mu(q')R(q', q)a(q', q).$$

Repair-informed proposals must include the Hastings asymmetry term; otherwise the stationary law is generically changed.

Repair generators are not selectors. A repair generator of the form

$$L_{\text{rep}} = \sum_C c_C (I - E_C)$$

is a relaxation or sampling object after a law has been selected. Conditional expectations E_C are defined on $L^2(X_r, \pi_r)$, so the reference law π_r is input. On overlapping collars the expectations need not commute. The correct finite gap certificate is the Poincare constant

$$\kappa_r = \inf_{f \perp 1} \frac{\sum_C \|(I - E_C)f\|^2}{\|f\|^2}.$$

If local fiber rates have a positive lower bound γ_* , then

$$L_{\text{rep}} \geq \gamma_* \kappa_r (I - P_0).$$

Finite repair completeness gives $\kappa_r > 0$ at fixed regulator. A uniform refinement lower bound $\inf_r \kappa_r > 0$ is a separate theorem or receipt.

Finite evidence accuracy. For bounded coarse observables O , if

$$\|\hat{\mu}_s - \mu_s\|_{\text{TV}} \leq \epsilon_{\text{samp}}$$

and the refinement defects sum to ϵ_{ref} , then

$$\left| \mathbb{E}_{\hat{\mu}_s}[O \circ c_{sr}] - \mathbb{E}_{\mu_r}[O] \right| \leq 2\|O\|_{\infty}(\epsilon_{\text{samp}} + \epsilon_{\text{ref}}).$$

Continuum-facing observables require a realization map and correlation Cauchy bound, not just a finite histogram.

Vacuum promotion gate. A stationary sampler is not a physical vacuum. For any faithful target law one can build a positive transfer operator with that law as ground state, so positivity alone is not a selector. Vacuum promotion requires source Euclidean slab data

$$\mathfrak{S}_r^E = (Q_r, m_r^0, J_r, V_r, a_{t,r})$$

whose conductance $J_r(q, q') = J_r(q', q) \geq 0$, local potential V_r , and slab thickness $a_{t,r}$ are derived without using the target law or sampler output. With connected event graph,

$$(H_r^E f)(q) = \frac{1}{m_r^0(q)} \sum_{q'} J_r(q, q') (f(q) - f(q')) + V_r(q) f(q)$$

is self-adjoint and bounded below on $L^2(Q_r, m_r^0)$; its finite Feynman–Kac semigroup is positivity improving. Perron–Frobenius gives a unique positive normalized ground state Ω_r , and the finite vacuum law is

$$\mu_r^{\text{vac}}(q) = |\Omega_r(q)|^2 m_r^0(q).$$

For $T_r = e^{-a_{t,r}(H_r^E - E_{0,r})}$, the Doob kernel is stochastic and detailed-balanced with μ_r^{vac} . Continuum promotion additionally requires reflection positivity or equivalent reconstruction plus refinement compatibility of the transfer family.

Primordial and cosmological prediction firewall. A screen covariance is not automatically a primordial curvature spectrum. The map

$$C_\ell = 4\pi \int_0^\infty \frac{dk}{k} \Delta_\zeta^2(k) j_\ell^2(k\chi_\star)$$

has a null space unless a radial prior, finite parametric family, or source-stress bridge is declared. OPH primordial promotion requires the source-only stress, single-clock, entropy-repair, curvature-evolution, adiabatic-mode, phase-coherence, screen-to-radial-lift, radial-null-space, and forward-projection receipts. Observable CMB promotion further requires frozen source, solver, likelihood, no-data-use, pooled-reducer, and falsification-rule receipts.

Claim tiers and required receipts. Every ensemble-facing run records immutable ensemble id, claim tier, regulator id, representative schema hash, gauge action hash, canonicalizer hash, base measure, action coefficients, coarse maps, zero-mode projector, amplitude convention, sampler hash, smoothing policy, source provenance, and explicit nonclaims. The seed belongs to the run receipt, not to the ensemble definition. The claim tiers are

- $E0$: seed noise, proposal noise, repair jitter,
- $E1$: conventional reference ensemble,
- $E2$: OPH-native quotient ensemble,
- $E3$: OPH vacuum,
- $E4$: OPH primordial field,
- $E5$: observable cosmological prediction.

The evidence bundle must keep separate receipts for stationary-law schedule invariance, detailed balance of the aggregate kernel, and pathwise partition invariance. Deterministic replay of semantic random streams or a canonical serial chain is useful, but it is not pathwise partition invariance. Smoothing must preserve raw coefficients, raw spectra, smoothing kernels, smoothed coefficients, smoothed spectra, and hashes of each stage; it is not part of S_r unless explicitly declared.

3 Finite Screen Spectrum Theorem Package

This fragment owns the screen-level spectrum theorem used by the staged cosmology branch. It does not own TT, TE, EE, lensing, likelihoods, or physical CMB promotion. Those remain downstream Boltzmann and data-contract gates.

Definition 3.1 (finite screen regulator). *For regulator r , a finite screen regulator is*

$$\mathfrak{S}_r = (\mathcal{T}_r, M_r, \Gamma_r, \mathcal{Q}_r, C_{sr}, J_{rs}),$$

where $\mathcal{T}_r$ is a finite cellulation of the screen, $M_r \succ 0$ is the area/quadrature mass matrix, Γ_r is the hidden-presentation groupoid, $\mathcal{Q}_r = \Sigma_r/\Gamma_r$ is the physical quotient, and C_{sr} , J_{rs} are the coarse-graining and interpolation maps. Shape regularity and quadrature convergence require

$$f^\top M_r g \rightarrow \int_{S^2} f g d\Omega$$

with an explicit band-limited error bound

$$\left| f^\top M_r g - \int f g d\Omega \right| \leq \varepsilon_{M,r}(L) \|f\|_{H^s} \|g\|_{H^s}.$$

Patch count alone is therefore not an angular-resolution certificate.

Definition 3.2 (geometric screen scalar). *Let the quotient-visible collar-volume readout be*

$$J_{X,r}(x) = \lambda_r(x) \sqrt{\det \sigma_{AB,r}(x)} > 0,$$

and let $\bar{J}_{X,r}$ be emitted by an independently defined homogeneous or frozen-background branch, not by a CMB fit. Define

$$q_{0,r} = \frac{1}{3} \log \frac{J_{X,r}}{\bar{J}_{X,r}}.$$

On a certified spherical chart, with $B_r = [1, n_x, n_y, n_z]$,

$$\Pi_{<2,r} = B_r (B_r^\top M_r B_r)^{-1} B_r^\top M_r, \quad q_r = (I - \Pi_{<2,r}) q_{0,r}.$$

On an irregular screen, B_r is replaced by the certified generalized eigenprojector onto the finite $\ell = 0, 1$ subspace. Hard-coded feature z -scores or observer-summary weights are diagnostic proxies, not q_r .

Proposition 3.3 (quotient invariance). *If a recharting U preserves the screen inner product and geometric readout,*

$$U^\top M'_r U = M_r, \quad q'_{0,r} = U q_{0,r}, \quad B'_r = U B_r,$$

then $\Pi'_{\geq 2,r} U = U \Pi_{\geq 2,r}$, and $q'_r = U q_r$.

Proof. Substitution gives

$$\begin{aligned} \Pi'_{<2,r} U &= U B_r (B_r^\top U^\top M'_r U B_r)^{-1} B_r^\top U^\top M'_r U \\ &= U B_r (B_r^\top M_r B_r)^{-1} B_r^\top M_r = U \Pi_{<2,r}. \end{aligned}$$

Subtracting from U gives the high-mode identity. □

Definition 3.4 (physical scalar precision and action). *The scalar precision K_r must have a physical origin on $V_r = \text{im } \Pi_{\geq 2,r}$. Allowed branches are:*

$$K_r^{\text{Hess}} = \nabla^2 \Phi_r \Big|_{q=0},$$

for a quotient-visible mismatch or release free energy; a repair Dirichlet form

$$K_r^{\text{diss}} = H_r - T_r^\top H_r T_r, \quad T_r^\top H_r T_r \preceq H_r;$$

or the reversible generator form

$$K_r^{\text{Dir}} = -\frac{1}{2}(L_r + L_r^{\dagger H}).$$

A raw nonsymmetric repair matrix or caller-supplied κ is not a precision operator. The absolute normalization must be fixed independently, for example by

$$K_0 Y_{\ell m} = \ell(\ell + 1) Y_{\ell m}$$

on the $\theta = 0$ branch. Once normalized,

$$S_{\text{scr},r}[q] = \frac{1}{2A_{q,r}} \langle q, K_r q \rangle_r = \frac{1}{2A_{q,r}} q^\top M_r K_r q.$$

Theorem 3.5 (finite MaxEnt screen covariance). *Let $K_r e_i = \kappa_i e_i$ with an M_r -orthonormal basis of V_r , $\kappa_i > 0$, and $d_r = \dim V_r$. Among continuous densities in coefficients $q = \sum_i a_i e_i$ with $\mathbb{E}[a_i] = 0$ and*

$$\frac{1}{2} \mathbb{E} \langle q, K_r q \rangle_r = E_{q,r}^{\text{src}},$$

the unique entropy maximizer is

$$d\mu_r(q) = Z_r^{-1} \exp \left[-\frac{1}{2A_{q,r}} \langle q, K_r q \rangle_r \right] dq, \quad A_{q,r} = \frac{2E_{q,r}^{\text{src}}}{d_r},$$

and

$$\mathbb{E}[a_i a_j] = \delta_{ij} \frac{A_{q,r}}{\kappa_i}.$$

Proof. The Euler–Lagrange equation for entropy with normalization and quadratic-energy constraints gives a Gaussian density with inverse temperature β_r . The expected energy is

$$E_{q,r}^{\text{src}} = \frac{1}{2} \sum_i \kappa_i \frac{1}{\beta_r \kappa_i} = \frac{d_r}{2\beta_r}.$$

Thus $A_{q,r} = \beta_r^{-1} = 2E_{q,r}^{\text{src}}/d_r$. Strict concavity of entropy gives uniqueness. $\square$

Remark 3.6 (discrete quotient branch). For a discrete finite quotient with intrinsic base weight $m_r(q)$, the exact Gibbs law is

$$\mu_r(q) = Z_r^{-1} m_r(q) e^{-\beta_r E_r(q)}.$$

A Gaussian statement on that branch requires a separate Laplace or central-limit theorem with controlled higher-order terms.

Definition 3.7 (source release energy). *The amplitude source is an independently selected quotient ensemble ν_r^{src} on frozen or released collar normal forms:*

$$E_{q,r}^{\text{src}} = \frac{1}{2} \int \langle q_r(x), K_r q_r(x) \rangle_r d\nu_r^{\text{src}}(x).$$

The receipt must expose ν_r^{src} , the base measure, K_r , d_r , $E_{q,r}^{\text{src}}$, $A_{q,r}$, no-observation ancestry, and the same operator normalization used in the action. MaxEnt alone does not determine amplitude.

Theorem 3.8 (rotational screen spectrum). *If the continuum precision K_θ commutes with the $SO(3)$ action on scalar functions, then by Schur's lemma each $\mathcal{H}_\ell$ is an eigenspace. For the exact conformal-shell family,*

$$\Lambda_\ell(\theta) = \frac{\Gamma(\ell + 2 + \theta/2)}{\Gamma(\ell - \theta/2)}, \quad \ell \geq 2.$$

Then $\Lambda_\ell(0) = \ell(\ell + 1)$, and the MaxEnt covariance gives

$$C_\ell^q = A_q \frac{\Gamma(\ell - \theta/2)}{\Gamma(\ell + 2 + \theta/2)}.$$

Consequently $\mathcal{D}_\ell^q = \ell(\ell + 1)C_\ell^q/(2\pi) \sim (A_q/2\pi)\ell^{-\theta}$.

Theorem 3.9 (finite refinement error). *Suppose that for $2 \leq \ell \leq L$*

$$\left| \frac{\lambda_{\ell m, r}}{\Lambda_\ell(\theta)} - 1 \right| \leq \varepsilon_{K, r}(L), \quad \left| \frac{A_{q, r}}{A_q} - 1 \right| \leq \varepsilon_{A, r}, \quad \varepsilon_{K, r} < 1.$$

Then $C_{\ell m, r}^q = A_{q, r}/\lambda_{\ell m, r}$ satisfies

$$\left| \frac{C_{\ell m, r}^q}{A_q/\Lambda_\ell(\theta)} - 1 \right| \leq \frac{\varepsilon_{A, r} + \varepsilon_{K, r}}{1 - \varepsilon_{K, r}}.$$

Proof. Write $A_{q, r} = A_q(1 + a_r)$ and $\lambda_{\ell m, r} = \Lambda_\ell(1 + b_{\ell m, r})$. Then

$$\frac{C_{\ell m, r}^q}{A_q/\Lambda_\ell} = \frac{1 + a_r}{1 + b_{\ell m, r}},$$

and the displayed bound follows from $|a_r| \leq \varepsilon_A$, $|b_{\ell m, r}| \leq \varepsilon_K$. □

Theorem 3.10 (refinement-semigroup tilt). *Let R_b be the scalar refinement map for scale ratio $b > 1$. If one isolated scalar covariance mode has positive eigenvalue $\lambda(b)$, $\lambda(b_1 b_2) = \lambda(b_1)\lambda(b_2)$, and λ is continuous, then there is a unique real θ with*

$$\lambda(b) = b^{-\theta}, \quad \theta = -\frac{\log \lambda(b)}{\log b}.$$

Proof. Set $g(t) = -\log \lambda(e^t)$. The semigroup law gives $g(t + s) = g(t) + g(s)$. Continuity makes $g(t) = \theta t$, hence the result. □

Target 3.11 (edge-center $P_\star/48$ tilt). *The identity $\theta = P_\star/48$ is a theorem-grade statement only after three lemmas pass: an oriented edge-center reserve $\rho_{\text{full}} = P_\star/24$, a scalar half-collar projector $\rho_q = P_\star/48$, and a reserve-to-RG Ward identity equating this reserve with $-d \log \mathcal{D}_\ell^q / d \log \ell$. Until then $P_\star/48$ and $e(P_\star - \varphi)$ are distinct diagnostic hypotheses.*

Theorem 3.12 (thin-shell power-law lift). Assume $q(\hat{n}) = Z_\star \Pi_{\ell \geq 2} \zeta_\star(R_\star \hat{n})$ and

$$\Delta_\zeta^2(k) = A_\zeta (k/k_\star)^{-\theta}.$$

Then

$$C_\ell^q = 4\pi Z_\star^2 \int d \ln k \Delta_\zeta^2(k) j_\ell^2(k R_\star),$$

and, for $-2 < \theta < 4$,

$$A_q = \pi^{3/2} Z_\star^2 A_\zeta (k_\star R_\star)^\theta \frac{\Gamma(1 + \theta/2)}{\Gamma(3/2 + \theta/2)}.$$

Thus

$$A_\zeta = \frac{A_q}{\pi^{3/2} Z_\star^2 (k_\star R_\star)^\theta} \frac{\Gamma(3/2 + \theta/2)}{\Gamma(1 + \theta/2)}.$$

Proposition 3.13 (finite-window bound). Let $\Psi_\ell(k) = \int dr W(r) j_\ell(kr)$, $W \geq 0$, $\int W dr = 1$, with mean $R_\star$ and variance σ_R^2 . If

$$\delta_\ell(k) = \frac{k^2 \sigma_R^2}{2} \sup_{r \in \text{supp } W} |j_\ell''(kr)|,$$

then

$$|C_{\ell,W}^q - C_{\ell,\text{shell}}^q| \leq 4\pi Z_\star^2 \int d \ln k \Delta_\zeta^2(k) \left[2\delta_\ell(k) + \delta_\ell(k)^2 \right].$$

Proposition 3.14 (radial non-identifiability). For a finite radial basis, $C = Ap$. If $A \in \mathbb{R}^{N_\ell \times N_k}$ has rank r , then $\dim \ker A = N_k - r$, and every $p + v$ with $v \in \ker A$ gives the same screen spectrum. Radial promotion therefore requires either a source theorem restricting p , or a declared prior with a published null basis, resolution kernels, positivity checks, and prior-sensitivity report.

Target 3.15 (screen-spectrum receipt set). The theorem-grade screen spectrum requires the following receipt families before any primordial or CMB promotion:

1. geometry, scalar quotient, low-mode projector, scalar precision, and operator normalization;
2. quotient ensemble selection, scalar release energy, and MaxEnt screen covariance;
3. scalar refinement tilt and angular screen spectrum with a finite error budget;
4. radial window, radial null-space, forward residual, and transfer firewall.

4 Physical Scale Bridge and Mode Calibration

This fragment owns the physical scale bridge used by the staged cosmology branch. It separates finite labels that are useful diagnostic proxies from labels that are physically calibrated for a Boltzmann solver.

Definition 4.1 (cosmology claim tiers). Every cosmology artifact has one of three claim tiers:

$$\text{DIAGNOSTIC_PROXY}, \quad \text{CONDITIONAL_PHYSICAL}, \quad \text{OPH_NATIVE_PHYSICAL}.$$

DIAGNOSTIC_PROXY covers cap angles, graph mode numbers, repair cycles, and other finite labels with no physical calibration. *CONDITIONAL_PHYSICAL* imports a frozen external FLRW geometry packet and then proves the downstream spectral, clock, freezeout, and transfer contracts relative to that packet. *OPH_NATIVE_PHYSICAL* requires the same packet to be derived from the OPH quotient carrier.

Definition 4.2 (notation firewall). *The branch uses separate symbols for separate objects:*

$$K_{\text{FLRW}} \in \{-1, 0, +1\}, \quad k, \quad \ell_{\text{src}}, \quad L_{\text{CMB}}, \quad \ell_{\star}, \quad L_{\text{IR}}.$$

K_{FLRW} is spatial curvature, k is comoving wavenumber, ℓ_{src} is an angular sector on the primordial source shell, L_{CMB} is the observed-sky transfer output multipole, $\ell_{\star}$ is the OPH scale-certificate length, and L_{IR} is an infrared domain size. The canonical internal unit for comoving k is Mpc^{-1} ; $h \text{Mpc}^{-1}$ is only a display unit after h is frozen.

Definition 4.3 (finite cosmological geometry packet). *At regulator r , a finite cosmological geometry packet is*

$$\mathfrak{G}_r = (Q_r, \{\Sigma_{r,n}\}, \{\mathcal{K}_{r,n}\}, \{h_{r,n}\}, \{\mu_{r,n}\}, \{N_{r,n}\}, \{\beta_{r,n}\}, \{u_{r,n}\}, \mathcal{L}_r, \mathcal{B}_r, \{\iota_{rs}\}, \mathcal{R}_{\ell_{\star}}).$$

Here Q_r is the physical quotient normal form, $\Sigma_{r,n}$ are common-clock spatial slices, $\mathcal{K}_{r,n}$ finite cell complexes, $h_{r,n}$ positive spatial metric or Dirichlet-form data, $\mu_{r,n}$ physical volume measures, $N_{r,n} > 0$ lapse, $\beta_{r,n}$ shift, $u_{r,n}$ the cosmological observer congruence, $\mathcal{L}_r$ lineage, $\mathcal{B}_r$ topology and boundary conditions, ι_{rs} refinement maps, and $\mathcal{R}_{\ell_{\star}}$ the independent dimensional-scale receipt. The readout

$$\text{CosmoGeomRead}_r : \text{nf}(Q_r) \rightarrow \mathfrak{G}_r$$

is OPH-native only if it factors through the quotient and is invariant under hidden carrier coordinates, port labels, schedules, worker partitions, and equivalent checkpoint presentations. For the conditional branch the same packet may be imported, but must record `geometry_origin=IMPORTED_FLRW`.

Definition 4.4 (source-screen embedding packet). *A screen mode has physical radial meaning only after an embedding packet*

$$\mathfrak{E}_r = (S_r, j_r : S_r \hookrightarrow \Sigma_{r,n_{\star}}, O_r, \chi_r, W_r(\chi), J_r)$$

has been supplied. S_r is the source screen, j_r embeds it in the freezeout slice, O_r is the observer or radial origin, χ_r is physical comoving geodesic distance, W_r is the radial release window, and J_r is the area/volume Jacobian. The window must obey

$$W_r(\chi) \geq 0, \quad \int W_r(\chi) d\chi = 1.$$

Definition 4.5 (FLRW reduction and physical mode packets). *An FLRW reduction of $\mathfrak{G}_r$ is*

$$\text{FLRWReduce}_r(\mathfrak{G}_r) = (a_r(\tau), \gamma_{r,ij}, K_{\text{FLRW}}, \bar{u}_r, \varepsilon_{\text{FLRW},r})$$

with

$$h_{r,ij}^{\text{phys}}(\tau) = a_r(\tau)^2 \gamma_{r,ij} + \delta h_{r,ij}.$$

The residual packet bounds shear, vorticity, acceleration, density-gradient, spatial-curvature, Hamiltonian, momentum, and domain-backreaction errors. A scalar mode packet supplies mass and stiffness forms

$$M_{r,ij}^{(0)} = \int_{\Sigma_r} \phi_i \phi_j d\mu_{\gamma_r}, \quad K_{r,ij}^{(0)} = \int_{\Sigma_r} \gamma_r^{ab} \nabla_a \phi_i \nabla_b \phi_j d\mu_{\gamma_r},$$

the generalized problem $K_r^{(0)} v_{rj} = \lambda_{rj} M_r^{(0)} v_{rj}$, spectral projectors, mode lineage, interval eigenvalues, matrix hashes, topology and boundary hashes, and a safe resolved band. Vector, tensor, and spin-2 sectors require their own compatible operators when stress or polarization variables are used.

Definition 4.6 (proper time, scale factor, and freezeout packet). *Proper time is read from lapse along the selected congruence after shift effects are accounted for. For a lineage-tracked comoving domain D_r ,*

$$a_{V,r}(\tau) = \left[\frac{V_r(\tau)}{V_r(\tau_0)} \right]^{1/3}, \quad \log \frac{a_{\Theta,r}(\tau)}{a_{\Theta,r}(\tau_0)} = \int_{\tau_0}^{\tau} \frac{\bar{\Theta}_r(\tau')}{3} d\tau',$$

and the receipt records

$$\varepsilon_{a,r} = \sup_{\tau} |\log a_{V,r}(\tau) - \log a_{\Theta,r}(\tau)|.$$

The convention $a(\tau_0) = 1$ is fixed before reporting $1+z = 1/a$; outside the certified FLRW/geodesic branch redshift is

$$1+z = \frac{(u_{\mu}p^{\mu})_{\text{emit}}}{(u_{\mu}p^{\mu})_{\text{obs}}}.$$

Conformal time is $\eta(\tau) - \eta(\tau_0) = \int_{\tau_0}^{\tau} d\tau'/a(\tau')$. Mode-wise freezeout requires persistent bounds on curvature, nonadiabatic, isocurvature, decaying-mode, repair-forcing, phase, and constraint residuals. If the mode release times do not share a common surface within tolerance, the output is a `MODE_DEPENDENT_FREEZEOUT_MAP`, not a single freezeout surface.

Proposition 4.7 (dimensional rescaling no-go). *If the finite data used by a proposed calibration are unchanged under $h \mapsto s^2h$, then $\Delta_{s^2h} = s^{-2}\Delta_h$ and physical wavenumber transforms as $k \mapsto k/s$. Therefore cap angle, graph incidence, patch count, and normalized covariance cannot determine nonzero physical k without a dimensional metric receipt.*

Proof. If $-\Delta_h v = \lambda v$, then $-\Delta_{s^2h} v = s^{-2}\lambda v$. The finite input has not changed while the physical wavenumber has. No function of the unchanged input can select the changed answer. $\square$

Proposition 4.8 (angular projection cannot determine radial k). *At finite angular cutoff the map $T : f(k) \mapsto \{\int d\log k \Psi_{\ell}(k) f(k)\}_{\ell \leq L}$ has finite-dimensional codomain and infinite-dimensional domain. Hence it has nontrivial kernel: angular screen modes do not uniquely determine a radial spectrum or physical k -label.*

Proposition 4.9 (rank-one clock and scale-factor no-goes). *A rank-one source family $\mathcal{F} = \hat{\mathcal{F}} \circ \chi$ fixes a common ordering, not proper elapsed time, because $f \circ \chi$ gives the same fibers for any strictly increasing f . Likewise $h = a^2\gamma$ is invariant under $a \mapsto ca$, $\gamma \mapsto c^{-2}\gamma$. A lapse or operational clock receipt and the convention $a_0 = 1$ are therefore separate physical inputs.*

Proposition 4.10 (cap modes and unit strings do not calibrate). *Cap eigenmodes depend on a self-adjoint boundary or extension choice, so they are not full-sky multipoles until a full-sky extension or mode-coupling operator is certified. Changing metadata from `inverse_cap_opening_angle_proxy` to `Mpc`⁻¹ changes no numerical object and therefore proves no calibration.*

Theorem 4.11 (finite operator and angular multipole consistency). *For conforming shape-regular finite spaces with consistent mass and stiffness forms and fixed boundary sector, $M_r = M_r^{\dagger} > 0$, $K_r = K_r^{\dagger} \geq 0$, and the generalized Rayleigh quotient converges to the Laplace–Beltrami quotient on each resolved band. For a closed source screen with area radius $R_{A,r}$, the normalized operator $\bar{\Delta}_{\Omega,r} = R_{A,r}^2 \Delta_{S_r}$ calibrates ℓ_{src} when the spectral projector onto $I_{\ell} = (\ell^2, (\ell+1)^2)$ has rank $2\ell+1$, bounded residual from $\ell(\ell+1)$, and converges under refinement.*

Proof. Positivity follows from the integral forms. Min–max convergence of the quadratic forms gives eigenvalue-cluster convergence; degenerate sectors are identified by projectors. The round-sphere eigenvalue $\ell(\ell+1)$ is isolated with multiplicity $2\ell+1$, so norm-resolvent convergence gives the Riesz projector. $\square$

Theorem 4.12 (physical comoving k , scale factor, and freezeout). *Assume a valid physical geometry packet, FLRW reduction, $a_0 = 1$, dimensional ruler, topology and boundary sector, and finite spatial operator. On the flat branch,*

$$-\Delta_{\gamma_r} v_{rj} = k_{rj}^2 v_{rj}.$$

If the finite eigenvalue uses h_r^{phys} , then

$$k_{rj} = a_r \sqrt{\lambda_{rj}^{\text{phys}}}, \quad k_{rj} = \frac{a_r}{\ell_\star} \sqrt{\widehat{\lambda}_{rj}^{\text{phys}}}$$

for the dimensionless eigenvalue $\widehat{\lambda}_{rj}^{\text{phys}} = \ell_\star^2 \lambda_{rj}^{\text{phys}}$. For $K_{\text{FLRW}} \neq 0$ the branch must define the solver map $k_{\text{solver}} = \kappa_{K_{\text{FLRW}}}(\lambda)$ and use the corresponding hyperspherical kernels. If the FLRW residual and volume/expansion consistency errors are below tolerance, a_D , z , and η are calibrated. A single freezeout surface is promoted only when the persistent mode-wise residuals and common-surface bound pass.

Proof. For $h^{\text{phys}} = a^2 \gamma$, homogeneity makes $\Delta_{h^{\text{phys}}} = a^{-2} \Delta_\gamma$, so the eigenvalue relation follows. The volume equation $\mathcal{L}_u dV = \Theta dV$ gives $d \log a_D / d\tau = \langle \Theta \rangle_D / 3$. Curvature conservation follows by integrating $|\zeta'| \leq \varepsilon_\star \mathcal{H} |\zeta|$ over the persistent interval, while the other residuals certify adiabatic growing-mode dominance and constraint consistency. $\square$

Theorem 4.13 (finite physical scale bridge). *Assume valid receipts for geometry or imported conditional geometry, source-screen embedding, FLRW background reduction, dimensional ruler, angular and spatial mode operators, mode normalization, proper time and scale-factor history, freezeout, multi-limit convergence, and no post-hoc calibration. Then the finite bridge*

$$\text{SOPH}_{r,\star} : (P_{r,j}, P_{r,\ell_{\text{src}}}, \chi_{r,n}) \longmapsto (k_{r,j}, \ell_{\text{src}}, a_r(\chi_{r,n}), z_r(\chi_{r,n}), \eta_r(\chi_{r,n}), \Sigma_{\star,r})$$

is invariant under hidden carrier presentation, unique within the declared topology and boundary sector, and convergent on each resolved physical band. The observed CMB multipole remains the transfer-solver output index:

$$C_{L_{\text{CMB}}}^{XY} = 4\pi \int d \log k \mathcal{P}_\zeta(k) \Delta_{L_{\text{CMB}}}^X(k) \Delta_{L_{\text{CMB}}}^Y(k).$$

Target 4.14 (exact physical-scale receipt conjunctions). *The code and papers use structured receipts, not bare booleans:*

$$\begin{aligned} \text{PHYSICAL_K_RECEIPT} &= R_{\text{geometry}} \wedge R_{\text{embedding}} \wedge R_{\text{scale}} \wedge R_{\text{FLRW}} \wedge R_{\text{operator}} \wedge R_{\text{normalization}} \wedge R_{\text{boundary}} \\ \text{SOURCE_ANGULAR_MODE_RECEIPT} &= R_{\text{sphere/extension}} \wedge R_{\text{angular operator}} \wedge R_{\text{cluster}} \wedge R_{\text{refinement}}, \\ \text{CALIBRATED_A_EVOLUTION_RECEIPT} &= R_{\text{proper time}} \wedge R_{\text{lineage}} \wedge R_{\text{FLRW}} \wedge R_{\text{volume}} \wedge R_\Theta \wedge R_{\text{consistency}} \wedge R_{\text{refinement}} \\ \text{PHYSICAL_FREEZEOUT_SURFACE_RECEIPT} &= R_{\text{spacelike}} \wedge R_{\text{persistence}} \wedge R_\zeta \wedge R_{\text{adiabatic}} \wedge R_{\text{isocurvature}} \wedge R_{\text{growing}} \wedge R_{\text{constraint}} \end{aligned}$$

*The aggregate bridge receipt is their conjunction with **SCALE_BRIDGE_REFINEMENT_RECEIPT** and **NO_POSTHOC_CALIBRATION_RECEIPT**. A unit string, cycle dictionary, cap angle, or common contract boolean cannot instantiate any of these receipts.*

Target 4.15 (native geometry extraction). *The remaining OPH-native theorem is the construction of*

$$\text{CosmoGeomRead}_r : \text{nf}(Q_r) \rightarrow (\mathfrak{G}_r, \mathfrak{E}_r)$$

*as a quotient-natural, positive, nondegenerate, causal-order-compatible, observer-clock-compatible, executor-independent, refinement-convergent, physically scaled source embedding. Without that theorem, the physical route is **CONDITIONAL_PHYSICAL** with an imported frozen FLRW geometry packet.*

5 Compatibility With the Core Stack

The compatibility table fixes how the inflation-free draft is read against the core papers.

Imported draft idea	Core-compatible reading
Inflation-free replacement	Phase III continuation only. The compact paper’s recovered core does not include an inflation replacement, physical CMB likelihood, H_0/S_8 branch, or dark-sector cosmology claim.
Finite normal form	Use quotient-level fixed-cutoff normal forms on the declared branch, unique only modulo boundary redundancy, implementation hiding, inert ancillary stabilization, and any required same-boundary unique-extension condition. Do not read the old draft as selecting a unique microscopic representative.
Screen spectrum and $n_s = 1 - P/48$	Screen-level theorem only after geometric q_r , normalized K_r , source release energy, refinement tilt, and angular-spectrum receipts pass. The special $P/48$ tilt is theorem-grade only after the edge-center reserve, half-collar projection, and reserve-to-RG receipts pass. Physical A_s , n_s , running, isocurvature, phase coherence, and TT/TE/EE spectra require the separate source-stress, single-clock, radial-prior, null-space, forward-residual, transfer, and likelihood receipts.
Dark/anomaly slot	Imported from <code>cosmology/oph_dark_matter_paper.tex</code> . For a source-only primordial certification run, the dark continuation is OFF unless a supplied abundance is explicitly typed <code>CONDITIONAL_SOURCE_STATE</code> ; transported Q_A is not by itself an OPH derivation of the homogeneous anomaly abundance.
Scalar anomaly rows	Rows for $\bar{\rho}_A$, $\bar{\rho}_{A,\text{eq}}$, and B_A are diagnostics unless the finite covariant parent also emits w_A , $c_{s,A}^2$, σ_A , Q_A^μ , exchange-current closure, recipient stress for nonzero exchange, gauge-independent variables, CDM-limit recovery, and refinement convergence. A transition-matrix number is $\gamma_{\text{repair step}}$ until promoted to Γ_{rec} by the physical clock, active-fiber, and common-parent response receipts.
Boltzmann and likelihood comparison	Physical promotion requires immutable source, solver, and likelihood hashes, official likelihood execution, and global CDM-limit reductions. Shard-local <code>any()</code> rollups or nonlinear averages before global pooling do not satisfy the contract.
Vacuum language	Finite reference states and free-field/lattice baselines remain reference ensembles. OPH-native vacuum promotion requires the quotient-ensemble selector plus source Euclidean slab data and a transfer/reflection-positive reconstruction gate from the compact/screen-microphysics theorem surface.

Imported draft idea	Core-compatible reading
Simulator receipts	Any physical-promotion receipt cited here must carry exact source/config/solver/likelihood hashes. The 256k run used for diagnostics records <code>git_commit=unknown</code> , so it is not used as exact replay provenance.

6 Source Corpus

This paper consolidates these active surfaces.

Surface	Imported role
<code>cosmology/oph_inflation_without_inflaton_observer_screen_synchronization.tex</code>	Main inflation-free branch: finite screen federation, flatness, horizon coherence, geometric screen spectrum, screen amplitude, radial lift, MaxEnt release, anomaly slot, transfer, and likelihood gates.
<code>cosmology/physical_cmb_theorem_program.md</code>	Theorem-contract file identifying the physical CMB source, scale, stress, kernel, transfer, and likelihood contracts.
<code>paper/recovering_relativity_and_standard_model_structure_from_observer_overlap_consistency_compact.tex</code>	Compact recovered-core claim boundary and the first surface for later propagation.
<code>paper/tex_fragments/PAPER.tex</code>	Main technical paper surface for global OPH theorem language.
<code>paper/tex_fragments/PRIMORDIAL_BRIDGE_THEOREMS.tex</code>	Primordial bridge theorem boundary and screen-to-primordial finite-source conditions.
<code>paper/screen_microphysics_and_observer_synchronization.tex</code>	Finite observer-screen carriers, records, patch ports, evidence bundles, and implementation invariance.
<code>cosmology/oph_dark_matter_paper.tex</code>	Collar-remainder dark/anomaly stress, galaxy equilibrium, dynamic stress, and cosmological kernel contracts. This paper imports those contracts by reference; it does not duplicate the dark-sector derivations.
<code>oph-physics-sim/runs/gcp_large_vis_256k_model_kernel_20260623</code>	256k-patch simulator evidence: strong CMB diagnostics, explicit false physical-prediction receipt, and full visualizer export data. The run manifest records <code>git_commit=unknown</code> ; the canonical simulator repository is https://github.com/muellerberndt/oph-physics-sim , and the public source reference available for this consolidation is <code>aeaa9e502491a277b5e21a80bb89df6d089a074d</code> .

Remark 6.1 (Simulator receipt provenance). Whenever this paper mentions simulator receipts, the receipt artifacts should be traced to the responsible commit in the canonical simulator repository.

tory, <https://github.com/muellerberndt/oph-physics-sim>. For the 256k run used here, the local artifact `runs/gcp_large_vis_256k_modkernel_20260623/manifest.json` reports `git_commit=unknown`. Therefore the numerical run is cited as diagnostic evidence with incomplete commit provenance. The available public source context for this consolidation is <https://github.com/muellerberndt/oph-physics-sim/commit/aeea9e502491a277b5e21a80bb89df6d089a074d>, but that link is not asserted to be an exact run-replay hash for the 256k artifacts. Physical-promotion receipts require exact source, config, solver, and likelihood hashes.

Remark 6.2 (Ownership boundary for the dark-matter paper). The released dark-matter paper `cosmology/oph_dark_matter_paper.tex` keeps ownership of the dark-sector theory: collar-remainder stress, galaxy equilibrium, dynamic repair-stress transport, cluster behavior, and the detailed perturbation-kernel derivations. This cosmology program paper only records the interface that paper must expose to the physical CMB/large-scale-structure pipeline. When it names $\bar{\rho}_A(a)$, $B_A(k, a)$, stress closure, repair-stress transport, or a certified $\Gamma_{\text{rec}}(k, a)$, it is naming imported source functions and receipts, not rederiving the dark sector here.

7 Imported Theorem Map

The inflation-free draft contains many theorem statements. The table gives them a public-core landing surface and keeps their dependency roles visible.

Imported theorem family	Role in this paper
Finite synchronization normal form	Defines the early observer-facing quotient state whose repair normal form replaces an assumed smooth primordial substrate, with uniqueness read modulo the compact-paper quotient equivalences.
Clocked FLRW curvature holonomy, flat-sector selection, and curvature bounds	Owns the flatness branch: zero clock-slice spatial Levi-Civita holonomy identifies $\kappa = 0$; exact selection requires a direct theorem, conditional CMH selector, or explicit flat-branch assumption. Curvature damping supplies only $ K $ or $ \Omega_K $ bounds.
Diffusive horizon no-go, same-boundary coherence, and low- k repair gap	Separates a failed purely local diffusion story from the allowed same-boundary or uniform low- k synchronization mechanisms.
Geometric screen scalar, precision operator, MaxEnt screen covariance, Mellin lift, edge-center scalar opportunities, $\mathbb{Z}_6$ reserve, and half-collar sampling	Owns the screen-level near-scale-invariant covariance. The red tilt $P/48$ is theorem-grade only after the reserve-to-RG receipts pass; physical primordial promotion waits for the source-only radial bridge receipts.
Finite release collar state, scalar release code, and scalar amplitude theorem	Owns A_ζ as a finite source artifact rather than a fitted amplitude.
MaxEnt release, entropy transport, BBN-safe release, recombination inheritance, and adiabaticity	Connects the synchronized finite screen/collar state to a hot radiation branch with ordinary low-energy thermal history.

Imported theorem family	Role in this paper
Freezeout, scalar quotient record, gauge-invariant ζ , isocurvature decay, and superhorizon conservation	Imported only through the stricter primordial bridge: to- tal stress closure, single clock, repair gap, freezeout, grow- ing mode, isocurvature, phase coherence, radial prior, null- space, and forward residual.
Finite homogeneous anomaly charge, no double count- ing, packet moments, cold anomaly, Bianchi exchange, no-slip lensing, and linear anomaly transfer	Imported from <code>cosmology/oph_dark_matter_paper.tex</code> ; this paper only records the CMB/LSS handoff interface and receipt gates.
Universe Simulation certifi- cate bundle, no-data-use theorem, finite-certificate branch, Boltzmann stability, likelihood promotion, and CMB admissibility	Owens the promotion rule: source-only finite artifacts first, frozen transfer and likelihood second, physical CMB claim last.
Tensor repair gap and finite- collar non-Gaussianity	Records secondary predictions to be tested only after the primary scalar/source/stress gates close.

8 Extended Cosmology Agenda

The local cosmology notes identify several OPH-native prediction surfaces beyond the immediate CMB gate. They are program tracks, not independent claims.

1. **CMB as observer-consensus fossil.** The highest-value target is the CMB: largest observ-
able scales should carry relics of finite observer synchronization rather than a fitted inflationary
initial spectrum.
2. **Λ from observer capacity.** The screen-capacity branch relates the de Sitter scale to finite
horizon capacity. A full cosmology paper should derive the equation of state, possible time
evolution, and any deviation from $w = -1$.
3. **Full cosmological dark sector.** The dark paper’s galaxy layer is not enough. CMB, BAO,
weak lensing, growth, and S_8 need the transported repair-stress perturbation branch.
4. **H_0 and S_8 predictions.** With the scale bridge and anomaly kernels fixed, the branch should
emit blind values or intervals for H_0 and S_8 , without tuning toward observed tensions.
5. **Large-scale topology.** Finite observer-accessible screens make global topology and matching-
circle style signatures natural tests.
6. **Consensus hierarchy of structure.** Galaxy, cluster, and supercluster formation should be
investigated as a hierarchy of stable consensus attractors, with density perturbation growth in
a fixed background as one layer of the story.
7. **Cosmological arrow of time.** OPH has native records, checkpoints, and continuation laws;
these should be used to formulate the low-entropy past condition as a finite-record boundary
condition.

8. **Cosmological neutrino background.** The particle/mass branch should eventually feed neutrino background and free-streaming predictions into the same frozen likelihood contract.

9 Finite Cosmological State

Definition 9.1 (Early OPH screen federation). *An early OPH cosmological state at regulator r is a finite patch federation*

$$\mathcal{F}_r = (V_r, E_r, \{\mathcal{A}_i, \rho_i, \mathcal{R}_i\}_{i \in V_r}, \{\mathcal{I}_e, \pi_{i,e}\}_{e \in E_r}, \mathcal{U}_r, \text{Chk}_r),$$

with finite accessible algebras, record algebras, visible overlap interfaces, repair instruments, and checkpoint data. Its physical content is the quotient under hidden carrier coordinates that do not affect visible interface records, repair maps, or checkpoint continuation.

Definition 9.2 (Source-only finite artifact). *A finite artifact X is source-only for cosmology if it is a deterministic functional of declared OPH source data, release-branch constants, and standard non-CMB physical constants, and if its dependency manifest excludes CMB, BAO, supernova, weak-lensing, RSD, SPARC, cluster, or other observational likelihood values used to evaluate the prediction.*

Definition 9.3 (Physical CMB promotion). *An OPH CMB run is physically promoted only when the primordial source, scale bridge, dark/anomaly stress kernels, Boltzmann solver input files, solver version/tolerances, and likelihood code/data hashes are all frozen before likelihood evaluation, and all source functions are source-only finite artifacts.*

Definition 9.4 (Cosmology artifact promotion ladder). *Cosmology artifacts first carry the three claim tiers of Definition 4.1. Within a tier, source artifacts also carry an explicit semantic type*

DIAGNOSTICPROXY $\prec$ FINITESCREENSOURCE $\prec$ FINITEPRIMORDIALSOURCE $\prec$ PHYSICALBOLTZMANNSOURCE

*No receipt may skip a boundary in this order. A close TT curve, a scalar CMI row, a cap-angle mode number, a repair-clock eigenvalue, or a visual freezeout surface is therefore **DIAGNOSTIC_PROXY** until the required source, physical-scale, stress, transfer, and likelihood receipts promote it step by step. Imported FLRW geometry can promote downstream labels only to **CONDITIONAL_PHYSICAL**; **OPH_NATIVE_PHYSICAL** additionally requires Target 4.15.*

Definition 9.5 (Proof-carrying source artifact). *A promotable artifact is a record*

$$\mathfrak{A}_v = (id_v, \tau_v, x_v, \mathcal{X}_v, \mathcal{U}_v, \text{Par}(v), \mathcal{D}_v, \mathcal{E}_v, h_v^{\text{theorem}}, h_v^{\text{witness}}, h_v^{\text{build}}, h_v^{\text{artifact}}),$$

where τ_v is the semantic type, x_v is the value or tensor/function, $\mathcal{X}_v$ records axes, domain, grid, and support, $\mathcal{U}_v$ records units, gauge, frame, pivot, and conventions, $\text{Par}(v)$ is the complete parent set, $\mathcal{D}_v$ is the data-class ledger, $\mathcal{E}_v$ is the numerical and theorem error envelope, and the hashes commit to theorem version, witness, build environment, and artifact payload.

Theorem 9.6 (Sound promotion by source DAG induction). *Let $G = (V, E)$ be a finite acyclic artifact graph. For each node v , let $P_v(\mathfrak{A}_v)$ be the mathematical predicate claimed by the artifact and let $V_v(\mathfrak{A}_v, w_v)$ be a sound verifier, so*

$$V_v(\mathfrak{A}_v, w_v) = 1 \implies P_v(\mathfrak{A}_v).$$

Define

$$\text{Pass}(v) = V_v(\mathfrak{A}_v, w_v) \wedge \bigwedge_{p \in \text{Par}(v)} \text{Pass}(p) \wedge \text{SourceOnly}(v).$$

Then $\text{Pass}(v) = 1$ implies $P_v(\mathfrak{A}_v)$, and every required parent predicate holds.

Proof. Order the finite DAG topologically. Root nodes satisfy their predicates by local verifier soundness. If every parent of v has passed, then the local verifier proves P_v and the parent hypotheses are available. The claim follows by induction. $\square$

Target 9.7 (Source ancestry and freeze receipts). *Physical promotion requires*

TRANSITIVE_SOURCE_ANCESTRY_RECEIPT, HERMETIC_READ_SET_RECEIPT, SOURCE_MODEL_FREEZE_RECEIPT.

The first receipt walks every ancestor of every promoted quantity and rejects any CMB, BAO, supernova, weak-lensing, growth, cluster, or similar evaluation-data dependency. The second requires an intercepted, content-addressed read set for files, network, environment, command-line configuration, caches, randomness, code, dependencies, and human-selected branch identifiers. The third freezes branch, model family, basis, pivot, cutoff, packetization, smoothing, thresholds, freezeout tolerances, and hyperparameter ranges before comparison.

10 Inflation-Free Branch

The inflation-free branch replaces inflation’s jobs rather than reproducing an inflaton. Its load-bearing theorem stack is:

1. quotient-level observer-facing normal form for the early screen/collar federation, with the compact-paper quotient equivalences;
2. zero clock-slice spatial Levi–Civita holonomy to identify the flat branch, plus a direct flatness theorem, conditional CMH selector, or explicit flat-branch assumption to select it;
3. same-boundary scalar normal form or a low- k repair gap for horizon coherence;
4. geometric screen spectrum for near scale invariance;
5. protected $\mathbb{Z}_6$ half-collar reserve for the red spectral exponent;
6. finite scalar release code for A_ζ ;
7. MaxEnt release and common release clock for a hot adiabatic start;
8. imported dark/anomaly stress variables for the pre-recombination CDM-like slot, when the `cosmology/oph_dark_matter_paper.tex` source contract permits them;
9. Boltzmann transfer and frozen likelihood for observational comparison.

Theorem 10.1 (Conditional OPH replacement of inflation). *Assume a cofinal family of finite early OPH screen federations has a quotient-level observer-facing normal-form projection on a declared branch, with uniqueness understood modulo boundary redundancy, implementation hiding, inert ancillary stabilization, and any same-boundary unique-extension condition required by the branch. Assume the clocked FLRW continuation boundary $B_{\text{cos},r}$ supplies $u_a = -N\nabla_a\chi$, $h_{ab} = g_{ab} + u_a u_b$, $K(\tau) = \kappa/a^2$, and topology policy, and that one of the exact flat-sector gates is present: DIRECT-THEOREM, CONDITIONALCMH, or EXPLICITASSUMPTION. Assume every CMB-scale scalar mode shares the same boundary normal form or satisfies a uniform low- k repair gap, the finite screen-spectrum receipts derive a geometric q_r , normalized K_r , source release energy A_q , and refinement tilt, the source-only radial bridge receipts pass, and the imported dark/anomaly sector supplies a permitted CDM-like source state through recombination. Then the branch supplies flatness, horizon coherence, near scale invariance, hot adiabatic initial data, and acoustic-transfer input without an inflaton degree of freedom.*

Proof. The proof is a dependency composition. The quotient normal-form projection gives a finite observer-facing early state on the declared branch, not a unique hidden representative. The spatial Levi-Civita holonomy theorem identifies the $\kappa = 0$ branch; the direct, CMH, or explicit-assumption gate selects it. Curvature damping is not used as a selector. Same-boundary repair or a low- k gap gives coherence on the observed CMB band. The screen-spectrum theorem package first gives

$$C_\ell^q = A_q \frac{\Gamma(\ell - \theta/2)}{\Gamma(\ell + 2 + \theta/2)}.$$

The value $\theta = P_*/48$ follows only if the edge-center reserve, half-collar scalar projection, and reserve-to-RG receipts pass; otherwise θ is measured from the actual scalar refinement operator or kept as a diagnostic hypothesis. The source-only bridge receipts then type that screen scalar as a primordial curvature source and convert A_q to A_ζ only after the exact thin-shell or finite-window radial lift, null-space, and forward-residual receipts pass. Without them it remains a screen theorem. MaxEnt release gives a hot radiation state and the common clock gives adiabatic leading perturbations. If the imported anomaly stress branch is an admissible source state through recombination, the acoustic peaks are computed by ordinary Einstein-Boltzmann transfer from those finite inputs. $\square$

Proposition 10.2 (Curvature suppression is not sector selection). *On a fixed clocked FLRW sector,*

$$K(\tau) = \kappa/a(\tau)^2, \quad \dot{K} = -2HK.$$

Any repair or expansion estimate that proves a finite bound on $|K|$ or $|\Omega_K|$ is an APPROXIMATE-BOUND. It does not change κ , does not distinguish globally inequivalent flat topologies, and may not be reported as DIRECTTHEOREM or CONDITIONALCMH.

Remark 10.3 (Theorem boundary). The theorem is conditional because several premises are open theorem issues. The branch is sharp enough for diagnostics, but not for promoting CMB agreement to a physical prediction.

11 Physical CMB Finite-Source Contract

The simulator's false physical CMB receipt identified a precise theorem contract. The following objects must be theorem-grade finite artifacts:

$$A_\zeta, \quad q_{\text{IR}}, \quad \ell_{\text{IR}}, \quad \bar{\rho}_A(a), \quad \bar{\rho}_{A,\text{eq}}(a), \quad B_A(k, a), \quad \gamma_{\text{repair step}}(k, a) \text{ or certified } \Gamma_{\text{rec}}(k, a),$$

plus full stress variables, freezeout, physical mode calibration, and N_{CRC} provenance.

Definition 11.1 (Finite primordial source package). *At regulator r , a physical primordial-source claim is carried by*

$$\mathfrak{S}_r^{\text{prim}} = (\mathcal{Q}_r, n_r, \mu_r, q_r, J_r, \Pi_r, K_r, M_r, \Xi_r, \Sigma_r, E_r^{\text{rel}}, A_{q,r}, \theta_r, q_{\text{IR},r}, t_{\text{IR},r}, \Sigma_{\star,r}, W_r, \Delta_{\zeta,r}^2, \mathcal{P}_{IJ,r}, \mathcal{E}_r).$$

Here $\mathcal{Q}_r$ is the physical quotient state space, n_r the normal-form map, μ_r the source probability law, q_r the quotient-visible scalar source, J_r the finite carrier-to-harmonic readout, Π_r the gauge/monopole/dipole projector, K_r the positive scalar precision operator, M_r the mobility or repair-response operator, Ξ_r the stochastic forcing covariance, $\Sigma_r = \text{Cov}_{\mu_r}(q_r)$, E_r^{rel} the source release-energy artifact, $A_{q,r}$ the finite screen amplitude, θ_r the spatial/RG spectral exponent, $q_{\text{IR},r}$ the IR suppression depth, $t_{\text{IR},r}$ the IR semigroup time, $\Sigma_{\star,r}$ the freezeout cut, W_r the radial window, $\Delta_{\zeta,r}^2$ the lifted primordial spectrum, $\mathcal{P}_{IJ,r}$ the species-level initial covariance, and $\mathcal{E}_r$ the complete error and refinement envelope.

Target 11.2 (Finite-source input contract). *The paper stack must define pass/fail receipts for:*

1. a source-provenance dependency DAG for all CMB source inputs;
2. globally pooled sufficient-statistic reducers before nonlinear source estimates;
3. the role of N_{CRC} as a consensus invariant, additive capacity, or theorem-side constant;
4. finite derivations of A_ζ , q_{IR} , and ℓ_{IR} ;
5. total stress closure, single-clock normal form, entropy repair gap, curvature evolution, adiabatic growing mode, isocurvature bound, phase coherence, screen-to-radial lift, radial null-space, forward residual, and finite freezeout receipts;
6. physical k , ℓ , scale-factor, and redshift calibration;
7. source-only imported dark-sector functions $\bar{\rho}_A(a)$, $\bar{\rho}_{A,\text{eq}}(a)$, $w_A(a)$, $c_{s,A}^2(k,a)$, $\sigma_A(k,a)$, Q_A^μ , $B_A(k,a)$, and the γ_{repair} step diagnostic or certified $\Gamma_{\text{rec}}(k,a)$, where the chosen dark-continuation mode permits them.

Target 11.3 (Scalar, harmonic, and IR source receipts). *The scalar-source branch must supply*

`SCALAR_SOURCE_MAP_RECEIPT`, `A5_TO_S03_INTERTWINER_RECEIPT`, `SPHERICAL_QUADRATURE_RECEIPT`, `LOW`

The map $\text{ScalarSourceMap}_r : \text{nf}(\mathcal{Q}_r) \rightarrow V_r^{\text{scalar}}$ must be quotient-invariant, refinement-natural, and normalized without per-run standardization. Harmonic reconstruction uses a weighted Gram matrix $G_L = Y_L^\dagger W Y_L$ and the low-mode projector

$$\Pi_{\geq 2} = I - Y_{01}(Y_{01}^\dagger W Y_{01})^{-1} Y_{01}^\dagger W.$$

The IR kernel is parameterized by the semigroup time

$$S_{\text{IR}} = I - q_{\text{IR}} e^{-t_{\text{IR}} L}, \quad F_{\text{IR}}(\ell) = 1 - q_{\text{IR}} e^{-t_{\text{IR}} \ell(\ell+1)}.$$

The validator must enforce $0 \leq q_{\text{IR}} \leq 1$, $t_{\text{IR}} \geq 0$, and derive the display scale

$$\ell_{\text{IR}} = \frac{-1 + \sqrt{1 + 4/t_{\text{IR}}}}{2}$$

when $t_{\text{IR}} > 0$. The old dimension count $12 + 20 + 1 = 33 \Rightarrow \ell_{\text{IR}} = 32$ is not a source theorem.

Target 11.4 (Fluctuation covariance and tilt receipts). *Deterministic settling does not choose a fluctuation ensemble. The finite quotient ensemble must supply $\mu_r(q) = m_r(q) e^{-S_r(q)} / Z_r$ with projective compatibility, sampler correctness, and implementation invariance. Repair drift does not determine covariance: $D\Sigma + \Sigma D^\top = \Xi$ requires the noise covariance Ξ . A fluctuation-dissipation branch may use $D = M_r K_r$, $\Xi = 2A_r M_r$, and $\Sigma_r = A_r K_r^{-1}$, but only with*

`QUOTIENT_ENSEMBLE_RECEIPT`, `LYAPUNOV_COVARIANCE_RECEIPT`, `FLUCTUATION DISSIPATION_RECEIPT`, `MOD`

The temporal repair gap, spatial RG exponent, primordial tilt, and physical exchange-rate kernel are distinct types:

`TEMPORAL_REPAIR_GAP`, `SPATIAL_RG_EIGENEXPONENT`, `PRIMORDIAL_SPECTRAL_TILT`, `PHYSICAL_EXCHANGE_R`

Only after the radial theorem identifies $\Delta_\zeta^2(k) \propto k^{-\theta}$ may the branch write $n_s = 1 - \theta$.

Proposition 11.5 (Why the 256k CMB curve is diagnostic). *The 256k OPH-FPE run may be used as a measurement-facing diagnostic, but not as a physical CMB prediction, because its promotion report marks the finite-source input contract false and lists missing source provenance, missing pooled reducers, missing N_{CRC} invariant role, non-finite A_ζ , missing screen-to-primordial lift, non-finite q_{IR} and ℓ_{IR} , non-finite $\bar{\rho}_A$ and B_A , missing finite covariant parent receipt, missing stress closure, missing full fluid/exchange variables, missing gauge/causal/refinement certificates, missing freezeout, missing official likelihood and CDM-limit reductions, and missing frozen solver and likelihood hashes.*

12 Physical Scale Bridge

Section 4 owns the bridge definitions, no-go propositions, and conditional physical-scale theorem. The practical rule is:

`inverse_cap_opening_angle_proxy, repair_cycle, graph_mode_index`

are nonpromotable diagnostic labels unless the structured receipt conjunctions in Target 4.14 pass. A frozen imported FLRW geometry packet can exercise the complete spectral/clock/freezeout/transfer pipeline at tier `CONDITIONAL_PHYSICAL`. Replacing that packet by a quotient-derived `CosmoGeomReadr` without changing downstream interfaces is the open `OPH_NATIVE_PHYSICAL` task.

13 Dark/Anomaly Interface Imported from the Dark-Matter Paper

The dark/anomaly slot is the largest cosmology-facing bridge between the OPH dark-matter paper and the CMB program. The released dark-matter paper `cosmology/oph_dark_matter_paper.tex` owns the derivation: collar remainder stress, settled galaxy response, dynamic transport, cluster behavior, and perturbation kernels. This paper only states the interface the CMB program requires from that paper. Cosmology needs the imported dark sector to expose a transported stress component with background density, perturbations, exchange terms, gauge-invariant variables, and the CDM-limit switch-off check.

Definition 13.1 (Imported finite covariant collar-packet parent interface). *For the purpose of this cosmology program, the finite covariant collar-packet parent is an imported `cosmology/oph_dark_matter_paper.tex`-owned finite artifact interface*

$$\mathcal{P}_A = (\mathcal{C}_r, Z_r, A_r, R_r, G_r, \pi_r, L_r, Q_r, D_r)$$

where Z_r splits anomaly and recipient packet states, π_r is the finite equilibrium packet functional, L_r is the repair generator, Q_r records energy-momentum reaction channels, and D_r records any causal auxiliary response. The interface emits packet stress moments

$$\rho_A, \quad P_A, \quad q_A, \quad \pi_A,$$

and source variables

$$\bar{\rho}_A(a), \quad \bar{\rho}_{A,\text{eq}}(a), \quad w_A(a), \quad c_{s,A}^2(k,a), \quad \sigma_A(k,a), \quad Q_A^\mu, \quad B_A(k,a), \quad \gamma_{\text{repair step}}(k,a) \text{ or certified } \Gamma_{\text{rec}}(k,a),$$

with explicit conservation or exchange equations, recipient stress for nonzero exchange, gauge projection, causal response, regulator-refinement convergence, and B_A built from the anomaly-frame baryon density $n_b^{(A)} = -u_{A\mu} J_b^\mu$.

Definition 13.2 (Issue #319 conditional source receipt). *The imported dark-sector parent first has to pass*

ISSUE_319_CONDITIONAL_SOURCE_RECEIPT.

This receipt is a conjunction of source-side receipts, not a scalar CMI row. It requires exactly one declared route,

SOURCE_ROUTE_RECEIPT,

unit metadata

ENTROPY_UNIT_RECEIPT,

the exact carried modular identity

$$\langle K_{AB} + K_{BD} - K_B - K_{ABD} \rangle = I(A : D|B),$$

*and the first-variation classification that prevents raw CMI from being silently identified with a linear fixed-reference modular-energy variation. The source route is either *FIXED_REFERENCE_MODULAR_ENERGY* or *NONLINEAR_CMI_STRESS*. The second route also requires*

CMI_TO_MODULAR_SOURCE_MATCHING_RECEIPT.

Both routes require collar-localization, BW normalization, physical diamond scale, ℓ^4 scaling, cover-independence, and stress-tomography receipts before the proper-radius normalization

$$T_A^{ab} u_a u_b = \frac{15\hbar c}{8\pi^2 \ell^4} R_{r,\ell}^{\text{src}}$$

may be imported. Classical diagonal CMI rows, cap-angle proxies, duplicated collar counts, or absolute-value rectified density rows do not satisfy this receipt by themselves.

Target 13.3 (Imported stress-parent receipt). *The simulator receipt*

FINITE_COVARIANT_PARENT_RECEIPT

must fail unless the owning dark-sector paper and its evidence bundle certify all of the following:

1. *ISSUE_319_CONDITIONAL_SOURCE_RECEIPT;*
2. *CMI_TO_MODULAR_SOURCE_MATCHING_RECEIPT* on the nonlinear-CMI route, or *FIXED_REFERENCE_MODULAR_EN* on the fixed-reference route;
3. *STRESS_TOMOGRAPHY_RECEIPT* and *SM_CURRENT_NULL_RECEIPT;*
4. *exact finite stress-energy closure for anomaly plus recipient stresses;*
5. *explicit recipient stress and equal-and-opposite exchange current whenever the exchange branch is nonzero;*
6. *detailed balance or a declared nonequilibrium exchange law;*
7. *gauge independence of the emitted perturbation variables, including the anomaly-frame baryon-density definition of B_A ;*
8. *causal response with subluminal finite characteristics;*
9. *convergence under regulator refinement;*

10. recovery of the CDM branch when exchange, pressure, sound speed, and anisotropic stress are switched off.

The finite parent receipt is source-side. It does not include official CMB likelihood execution. The later `PHYSICAL_CMB_PROMOTION_RECEIPT` additionally requires the physical scale bridge, physical ρ_A , B_A , and Γ_{rec} kernels, Boltzmann transfer, frozen source/solver/likelihood hashes, and the official likelihood run.

Remark 13.4 (Dark continuation modes). For a source-only primordial certification run, the dark-sector paper recommends `dark_continuation = OFF`. A supplied dark abundance may enter only as `dark_continuation = CONDITIONAL_SOURCE_STATE` until the dark-matter theorem surface derives the homogeneous anomaly abundance and finite kinetic stress construction. The physical cosmology branch may set `dark_continuation = PHYSICAL_PARENT` only after the finite covariant parent receipt, physical scale bridge, response/kernel receipts, and CDM-limit recovery all pass for the same frozen generation.

14 Physical Kernel Interface

The physical kernels are not scalar rows in a table. They are imported finite functionals with units, gauge-invariant definitions, paired controls, and refinement margins. Their derivation stays in `cosmology/oph_dark_matter_paper.tex`; this paper only states what the CMB/LSS pipeline must receive.

Target 14.1 (Physical anomaly kernel interface). *The released dark-matter paper must define deterministic receipts*

`RHO_A_SOURCE_RECEIPT`, `RHO_A_EQ_SOURCE_RECEIPT`, `B_A_KERNEL_RECEIPT`, `GAMMA_REC_SOURCE_RECEIPT`

These receipts pass only when $\bar{\rho}_A(a)$, $B_A(k, a)$, and $\Gamma_{\text{rec}}(k, a)$ are emitted as certified physical functions with $\bar{\rho}_{A,\text{eq}}(a)$, $w_A(a)$, $c_{s,A}^2(k, a)$, $\sigma_A(k, a)$, and Q_A^μ from the finite parent, physical scale calibration, gauge consistency, paired perturbation controls, active-fiber/physical-clock response receipts, CDM-limit recovery, and refinement convergence. The cosmology program consumes the emitted functions but does not redefine the dark-sector source law.

Diagnostic rows remain useful for debugging. They should be plotted, stress-tested, and compared to controls. They should not be accepted by the physical CMB contract unless they are promoted by the kernel receipts above. Until the physical clock, active fiber, conserved-sector decomposition, and common-parent response pole are certified, the simulator should emit the transition number as $\gamma_{\text{repair step}}$, not as Γ_{rec} . Likewise finite rows for ρ_A , $\rho_{A,\text{eq}}$, and B_A have label `DIAGNOSTIC_SCALAR_RESPONSE` until promoted by the parent and kernel receipts.

Target 14.2 (Finite covariant parent import). *The imported dark-sector parent is not a readiness boolean. The receipt consumed by this paper is the conjunction of primitive parent receipts owned*

by cosmology/oph_dark_matter_paper.tex:

*SOURCE_ROUTE_RECEIPT, ENTROPY_UNIT_RECEIPT, MODULAR_NONADDITIVITY_IDENTITY_RECEIPT,
CMI_FIRST_VARIATION_CLASSIFICATION_RECEIPT, SOURCE_LOCALIZATION_SATURATION_RECEIPT,
BW_BALL_NORMALIZATION_RECEIPT,
PHYSICAL_DIAMOND_SCALE_RECEIPT,
ELL4_SCALING_PLATEAU_RECEIPT,
COVER_INDEPENDENCE_RECEIPT, STRESS_TOMOGRAPHY_RECEIPT,
FINITE_PACKET_KINEMATICS_RECEIPT,
PACKET_MASS_SHELL_RECEIPT,
TRANSPORT_COVARIANCE_RECEIPT,
CHANNEL_FOUR_MOMENTUM_RECEIPT,
FINITE_PACKET_STRESS_READOUT_RECEIPT,
VARIATIONAL_MOMENT_STRESS_AGREEMENT_RECEIPT,
LOCAL_FRAME_COVARIANCE_RECEIPT,
CARRIER_QUOTIENT_INVARIANCE_RECEIPT,
TOTAL_STRESS_CLOSURE_RECEIPT,
EXPLICIT_RECIPIENT_STRESS_RECEIPT,
EXCHANGE_CURRENT_CLOSURE_RECEIPT,
SM_CURRENT_NULL_RECEIPT,
COSMOLOGICAL_GAUGE_INVARIANCE_RECEIPT,
FINITE_DOMAIN_OF_DEPENDENCE_RECEIPT,
SUBLUMINAL_CHARACTERISTICS_RECEIPT,
RETARDED_RESPONSE_RECEIPT, RESPONSE_STABILITY_RECEIPT,
REFINEMENT_CONVERGENCE_RECEIPT, CDM_LIMIT_RECOVERY_RECEIPT.*

The source parent does not contain solver or likelihood hashes. Those belong to the frozen generation and likelihood layer below.

15 Boltzmann Transfer and Frozen Likelihood

For finite physical source functions, the transfer problem is ordinary cosmology. OPH does not need a special CMB plotting rule. It needs a frozen handoff to CAMB, CLASS, or a declared independent Einstein-Boltzmann solver.

Target 15.1 (Frozen transfer and likelihood protocol). *A physical OPH CMB prediction requires:*

- 1. immutable hashes for source artifacts;*
- 2. immutable hashes for solver source, version, tolerances, and input files;*
- 3. immutable hashes for likelihood code, datasets, covariances, masks, and nuisance priors;*
- 4. a no-data-use manifest showing that observational likelihood values did not enter source artifact generation;*

5. global pooled reducers for nonlinear source estimates and global CDM-limit checks;
6. official likelihood execution readiness beyond a diagnostic binned-table comparison;
7. a falsification rule fixed before the likelihood run.

Definition 15.2 (Model generation). *A cosmology prediction generation is*

$$G_n = (H_{\text{source}}, H_{\text{scale}}, H_{\text{parent}}, H_{\text{kernel}}, H_{\text{init}}, H_{\text{solver}}, H_{\text{build}}, H_{\text{config}}, H_{\text{likelihood}}, H_{\text{data}}, H_{\text{priors}}, \mathcal{A}_n),$$

where $\mathcal{A}_n$ is the registry of observational datasets, residuals, posterior summaries, and diagnostic plots exposed during source development. Hashing the final artifact is not enough: if a Planck residual influenced source or model selection, the generation is calibrated on Planck even when the final bytes are frozen.

Theorem 15.3 (Physical OPH cosmology promotion). *Let R_{src} be source-only provenance and pooled-reducer receipt, R_{scale} the physical mode/clock/lift receipt, R_{parent} the finite covariant parent receipt, R_{kernel} the physical response/kernel receipt, R_{init} the regular initial-mode and Einstein-constraint receipt, R_{transfer} the Boltzmann well-posedness and numerical-convergence receipt, R_{freeze} the model-generation freeze receipt, and R_{like} the official likelihood execution receipt. Define*

$$R_{\text{physical}} = R_{\text{src}} \wedge R_{\text{scale}} \wedge R_{\text{parent}} \wedge R_{\text{kernel}} \wedge R_{\text{init}} \wedge R_{\text{transfer}} \wedge R_{\text{freeze}} \wedge R_{\text{like}}.$$

If every parent receipt is labelled PASS, then the resulting TT/TE/EE, lensing, matter power, growth, BAO, or RSD spectra are physical predictions of the frozen generation G_n . If any receipt is FAIL, OPEN_GATE, INCOMPLETE, or INVALIDATED, the spectra remain diagnostic or conditional according to the parameter ledger.

Proof. Each receipt states a typed premise needed by the next map: source fields, physical coordinates, parent stress, response kernels, initial state, transfer evolution, freeze integrity, and likelihood execution. Composition of certified maps gives a physical prediction only when every premise is available for the same frozen generation. A failed or open receipt removes at least one premise, so downstream software cannot promote the output. $\square$

Target 15.4 (Outcome and prediction-class ledger). *Likelihood outcomes are*

*INVALIDATED, NUMERICALLY_INCONCLUSIVE, FALSIFIED_ON_DECLARED_TEST,
PASSED_DECLARED_TEST.*

INVALIDATED is used when provenance or freeze integrity fails; it is not a physical falsification. Inputs are typed as

*OPH_DERIVED, EXTERNAL_FIXED, CONDITIONAL_COSMOLOGY, NUISANCE,
MEASUREMENT_CALIBRATED,*

and outputs as

*SOURCE_ONLY_OPH_PREDICTION, CONDITIONAL_OPH_PREDICTION,
FITTED_OPH_MODEL, DIAGNOSTIC_BASELINE.*

The 256k curve has label CMB0_LCDM_DIAGNOSTIC_BASELINE.

Remark 15.5 (Global reducers only). Shard-local nonlinear averages and shard-local `any()` rollups are not promotion rules. The source artifact must pool additive sufficient statistics globally, with validated units, coordinate grids, coverage, duplicate policy, interpolation policy, and covariance, before any nonlinear quantity such as amplitude, rank, condition number, isocurvature leakage, B_A , $\bar{\rho}_A$, or Γ_{rec} is evaluated; a transition diagnostic is recorded separately as $\gamma_{\text{repair step}}$.

Remark 15.6 (Comparison versus prediction). The best OPH diagnostic model in the 256k run is scientifically useful because it is close: shape correlation 0.9951542364, normalized RMSE 0.0984455548, amplitude-fit $\chi^2/\text{bin} = 41.6684770597$, zero mean peak-location error, and mean peak-height fractional error about 0.0704905. Those numbers justify fixing the theorem gates. They do not replace the gates.

16 Vacuum and Quantum-Fluctuation Boundary

The same finite-source discipline applies to vacuum claims. Seed noise, repair jitter, and reference free-field baselines are not an OPH-native quantum vacuum.

Target 16.1 (OPH-native vacuum gate). *An OPH-native vacuum simulation needs a quotient ensemble measure or density operator on the observable quotient algebra, refinement compatibility, implementation invariance, a sampler correctness proof, and source Euclidean slab data $(Q_r, m_r^0, J_r, V_r, a_{t,r})$ whose transfer operator is derived independently of the target law and sampler. Without this gate, vacuum-like fields in the simulator are reference baselines or diagnostics.*

17 Claim-Gate Summary

The paper-side gates are quotient ensemble measure, OPH-native vacuum promotion, Boltzmann transfer, frozen likelihood closure, finite-source CMB contract hardening, physical scale bridge, finite covariant collar-packet stress parent, physical anomaly kernels, flat-sector selection, screen-spectrum derivation, conditional collar source routing, repair-stress transport, and linear repair-stress perturbations. GitHub issues carry the mutable ownership, dependency, and completion state for those gates.

18 Physical Promotion Rule

Physical promotion requires exact receipt semantics for the objects the simulator is being asked to produce:

1. implement the finite-source dependency DAG and no-data-use firewall;
2. implement global pooled reducers for all nonlinear source estimates;
3. implement the `CONDITIONAL_PHYSICAL` imported-FLRW reference backend for the spectral, clock, freezeout, and transfer pipeline;
4. implement the structured physical scale bridge and freezeout certificates from Target 4.14;
5. implement the imported finite covariant parent and stress-closure receipts from the `cosmology/oph_dark_matter_paper.tex` dark-sector theory;
6. implement physical $\bar{\rho}_A$, B_A , and Γ_{rec} kernel receipts as imported dark-sector source functions;

7. freeze source, solver, and likelihood hashes;
8. run official likelihood comparisons and global CDM-limit reductions;
9. only then label the result `CONDITIONAL_PHYSICAL` or, after Target 4.15 passes, `OPH_NATIVE_PHYSICAL`.

The simulator may run diagnostics before physical promotion: visualization payloads, CMB overlays, vacuum baselines, defect worldlines, observer cameras, and H3 readouts. Those outputs carry diagnostic labels until the relevant receipt passes.

19 Conclusion

The 256k simulator almost-fits are not random noise. They expose stable structure close enough to justify the theorem gates, and they show why those gates are necessary. A physical cosmology prediction needs finite source-only objects, physical scale calibration, covariant stress closure, physical anomaly kernels, and frozen likelihoods. This paper is the staging surface for that work and the simulator contract for physical promotion.

References

- [1] B. Müller, A. Osika, K. Xue, and P. Nguyen, *Recovering Relativity and the Standard Model from Observer Overlap Consistency*, OPH compact paper, release r1502, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/recovering_relativity_and_standard_model_structure_from_observer_overlap_consistency_compact.pdf
- [2] B. Müller, A. Osika, K. Xue, B. Cassie, P. Nguyen, M. Ponder, and K. A. Anirudha, *Observers Are All You Need*, OPH synthesis paper, release r1502, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/observers_are_all_you_need.pdf
- [3] B. Müller, A. Osika, K. Xue, and P. Nguyen, *Reality as Consensus Protocol*, OPH consensus paper, release r1502, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/reality_as_consensus_protocol.pdf
- [4] B. Müller and K. Xue, *Federated Echoshedral Screen Microphysics: Patch Hardware, Records, and Observer Synchronization in OPH*, OPH microphysics paper, release r1502, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/screen_microphysics_and_observer_synchronization.pdf
- [5] B. Müller, *Observer-Patch Holography and the Dark Matter Phenomenon*, OPH cosmology paper, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/cosmology/oph_dark_matter_paper.pdf
- [6] B. Müller, *Inflation Without an Inflaton: Observer-Screen Synchronization as an OPH Cosmology Branch*, workspace cosmology draft, 2026.
- [7] B. Müller, *OPH-FPE finite screen-consensus and cosmology diagnostics*. <https://github.com/muellerberndt/oph-physics-sim>

- [8] Planck Collaboration, *Planck 2018 results. VI. Cosmological parameters*, Astron. Astrophys. 641, A6, 2020, arXiv:1807.06209. <https://arxiv.org/abs/1807.06209>
- [9] A. Lewis, A. Challinor, and A. Lasenby, *Efficient computation of CMB anisotropies in closed FRW models*, Astrophys. J. 538, 473–476, 2000, arXiv:astro-ph/9911177. <https://arxiv.org/abs/astro-ph/9911177>
- [10] D. Blas, J. Lesgourgues, and T. Tram, *The Cosmic Linear Anisotropy Solving System (CLASS). Part II: Approximation schemes*, JCAP 07, 034, 2011, arXiv:1104.2933. <https://arxiv.org/abs/1104.2933>