A Conditional Identification of the Yang–Mills Gap with a Finite Repair Gap

Dobrushin Repair Floors and a $\mathbb{Z}_2$ Test of the Transfer Receipt

B. Müller Jonathan Hill

November 11, 2026

Paper release: `r2065` **Released:** November 11, 2026

Author affiliation: Bernhard Mueller, Pragma Research Inc.

Abstract

A finite collar-repair generator has a positive relaxation gap when local rates and approximate tensorization have uniform positive bounds. A common ground-state transform identifies that gap with the transfer Hamiltonian gap. On finite $\mathbb{Z}_2$ gauge systems, the free Wilson transfer satisfies the tested constant-rate identity; interacting tests fail it, while the Kogut–Susskind Hamiltonian has exact positive fiber-dependent rates. A separate full Haar-link SU(2) source on an eight-vertex cube has an interacting vacuum gap between 21/2 and 1024/47 in its declared units, with a nonzero finite-energy local curvature fluctuation. Its heat-kernel transfer is positive and time reflection positive. Exact one-plaquette character reduction gives infinite-representation spectral and correlation enclosures. These are finite-volume results. The cube bound deteriorates with volume and scales as the inverse spacing. A nontrivial four-dimensional Yang–Mills theory additionally requires the stated common-source Schwinger-function limit, covariance, locality, vacuum noncollapse and uniform physical gap. The continuum repair/Hamiltonian identification is conditional on those premises. A photon-containing matter theory is a different source and has no positive global vacuum gap under its massless-photon interpretation.

1 Claim Boundary

The paper gives a proof-bearing projective weak-* / GNS extraction from the declared finite support-visible compact-gauge cylinder system. The finite positive-transfer and repair-gap mechanism is conditional on the finite ground-state-transform/cross-fiber and uniform-gap receipts. The finite ground-state-transform/cross-fiber receipt (Assumption 6.3) is the load-bearing unverified physical input: it assumes the gap-friendly spectral structure whose gap the later sections account for. Section 8 evaluates it on $\mathbb{Z}_2$ lattice gauge theory: the receipt holds analytically in the free limit, fails for the Wilson transfer matrix on the tested $L = 2, 3$ diagonal coupling grid, and gives the local Hamiltonian exactly in a fiber-dependent-rate form. The scan is not a no-go for other couplings, anisotropic limits, transfer objects, or that fiber-dependent-rate route.

The imported four-dimensional form theorem uses compact-gauge reconstruction, the four-dimensional scaling chart, the reflection-positive ordinary vacuum sector, the absence of additional

gauge-invariant relevant dimension-four pure-gauge operators beyond the positive quadratic curvature invariant, the support-visible cylinder extraction, and the renormalized regularity/universality receipt. The mass-gap step adds the finite transfer receipt, exact atomic heat-bath collars, a finite source-type table, and a uniform L^2 approximate-tensorization/influence certificate. On any branch carrying all of these and the continuum certificate, the proof identifies the gap exactly,

$$\Delta_{\mathrm{YM}} = \Delta_{\mathrm{rep}}.$$

Other positive gaps in the OPH construction are different objects and discharge none of these receipts. The finite local-domain source construction in the microphysics and particle papers has an exact positive gap for the sign-twisted graph Laplacian of one observer-visible seam complex [3, 4]; it supplies no compact-gauge repair generator L_r^{rep}, vacuum projector, ground-state transform, gauge-invariant excitation algebra or uniform cofinal bound. In the executable observer construction [9], the native action $H_{\mathrm{native}} = \sum_e (I - \Phi_e)$ is a positive local Liouville operator on operator space, not a Hamiltonian on state vectors, and the executed channel strictly attenuates coherent modes, which rules out unitary conjugation on its carrier. Its unread C and E channels have quantum spectral gap $(5 - \sqrt{5})/60$ per native attempt on the twelve-port carrier [11, 12], and its canonical scalar regulator has a finite-volume gap bound for a supplied positive mass [13].

The native density laws give a reflection-positive Gaussian reconstruction whose one-particle time shift is $e^{-\tau A}$, where τ is rescaled native attempt count and A acts on the two-dimensional screen readout. With the constant mode restored, the form of A is a conservative strongly local Dirichlet form with $n/49920 \le \lambda_n(A) \le 3n/2$ and heat-trace spectral dimension two [14, 15]. This free two-dimensional spectrum is not a four-dimensional Yang–Mills Hamiltonian, and its lowest eigenvalue is not a mass gap. The native gap closes under refinement because native repair conserves load. On the federated screens of 240, 960 and 3,840 ports, the per-attempt gap $\lambda_1(L_l)/(2M_l)$ of the port Laplacian L_l with M_l seams is 3.50×10^{-5}, 2.05×10^{-6} and 1.27×10^{-7}, a fall by a factor near sixteen per level [10]. The continuum operator A appears only after rescaling by $1260 \cdot 16^l$ attempts, a diffusive normalization whose semigroup $e^{-\tau A}$ is second order in space and first order in time; a Euclidean-covariant reading would instead require $e^{-|\tau|\sqrt{A+m^2}}$ for a mass $m \ge 0$. Single-site heat-bath collars P_v are the identity under a fixed total load, so the native seams lie outside the admissible collar tower of Definition 6.5, and $\lambda_1(A)$ is the finite-size gap of a massless two-dimensional operator.

The collar construction uses the consensus and compact-gauge interfaces [2, 1]; its heat-kernel normalization is stated in Section 13. These are distinct from the executable native observer dynamics [9].

2 The Clay Target

The Clay Mathematics Institute describes the Yang–Mills mass gap as the missing mathematical foundation behind the quantum theory used for nonabelian gauge forces [17]. Jaffe and Witten state the problem as follows: for every compact simple gauge group G, construct a nontrivial quantum Yang–Mills theory on $\mathbb{R}^4$ satisfying axiomatic properties at least as strong as the stated Wightman or Osterwalder–Schrader references and prove a mass gap $\Delta > 0$ [18, 19].

The argument below approaches that target through a different variable, within the claim boundary stated above. The starting data are finite observer patches, compact-gauge visible quotient data,

and exact repair collars. On a branch carrying the finite transfer receipt and the continuum certificate, the nonzero energy threshold is the cost of leaving the constant sector of the collar repair generator, which the receipt maps to the transfer vacuum.

3 Standing Setup

Fix a compact simple gauge group G carried by an OPH compact-gauge zero-obstruction vacuum branch, realized at fixed cutoff by the declared compact-gauge patch-carrier architecture. The architecture supplies finite local Hilbert spaces, exact local constraints, patch and overlap algebras, overlap sector projectors, record layers, and local repair interfaces.

Let r range over a cofinal refinement family of finite regulators. For each r, let

$$(\mathcal{H}_r, \Omega_r, H_r), \qquad T_r(t) = e^{-tH_r},$$

be the physical Euclidean Hilbert space, vacuum, Hamiltonian, and transfer semigroup.

Let $X_r := X_r^{(0)}$ be the support-visible time-zero compact-gauge quotient configuration space, let π_r be the stationary time-zero measure, and set

$$K_r := L^2(X_r, \pi_r).$$

Write $\omega_r(a) := \int_{X_r} a \, d\pi_r$ for the associated state functional; K_r implements the finite π_r-null support reduction by definition. Let $\mathcal{C}_r$ be the finite family of active repair collars. For each active collar $C \in \mathcal{C}_r$, let

$$\rho_C := \rho_{C,r} : X_r \to Y_{C,r}$$

be the complete repaired visible datum, and let

$$E_C : K_r \to K_r$$

be conditional expectation onto the ρ_C-measurable functions.

The proof stays on the ordinary or central zero-obstruction vacuum branch. The genuinely noncentral higher-gauge branch is a different fixed-cutoff sector in the OPH framework and is not used for the ordinary compact-simple G theorem below.

Notation	Meaning
X_r, π_r, K_r	support-visible time-zero quotient configuration space, stationary measure, and $L^2(X_r, \pi_r)$.
$\mathcal{C}_r$	finite active repair-collar family at regulator r.
ρ_C	complete repaired visible datum on collar C.
E_C	π_r-preserving conditional expectation onto ρ_C-measurable functions.
L_r^{rep}	ground-state transformed Euclidean repair generator at cutoff r.
$P_{0,r}$	projection onto constants in K_r, corresponding to the physical vacuum.

Notation	Meaning
U_r	finite-stage unitary from the physical Euclidean Hilbert space to the repair L^2 space.
$K, \mathcal{H}, U$	support-visible continuum repair Hilbert space, physical Hilbert space, and limiting unitary.
c_*	uniform active-collar repair-rate floor.
A_*, δ_*	uniform approximate-tensorization constant and certified global gap c_*/A_*.

4 Imported 4D Euclidean Yang–Mills Form

Assumption 4.1 (Support-visible compact-gauge Yang–Mills branch)**.** *The compact-gauge branch used below satisfies the following branch-local conditions.*

(i) the ordinary or central zero-obstruction compact-gauge sector survives refinement with compact simple structure group G;

(ii) the cofinal regulator tail satisfies the Standard Model gauge paper's explicit compact-gauge refinement receipt;

(iii) the support-visible quotient carries a four-dimensional Euclidean scaling chart;

(iv) the ordinary vacuum sector is reflection positive and has topological angle $\theta = 0$;

(v) the local finite-constraint MaxEnt/Gibbs family is gauge-invariant, Euclidean local, rotation-invariant, and refinement-stable;

(vi) no additional gauge-invariant relevant dimension-four pure-gauge operator remains on this branch besides the positive quadratic curvature invariant;

(vii) the branch is carried on a separated cofinal regulator system with the finite compact-gauge cylinder data defined in Section 9.

No continuum cylinder state, GNS space, transfer generator, or vacuum projection is assumed in this branch declaration.

Theorem 4.2 (Imported conditional OPH four-dimensional Euclidean Yang–Mills form)**.** *Under Assumptions 4.1 and 9.4, on an extracted cylinder family from Proposition 9.2 that lies on the declared local four-dimensional scaling branch, the continuum gauge-sector Euclidean action is*

$$S_E[A] = \frac{1}{4g^2} \int_{\mathbb{R}^4} \langle F_{\mu\nu}, F_{\mu\nu} \rangle \, d^4x, \qquad F = dA + A \wedge A, \tag{YM}$$

with compact simple structure group G. The extracted gauge-quotient cylinder family is denoted

$$d\mu_{\mathrm{YM}}(A) = Z^{-1} e^{-S_E[A]} \, DA/G$$

in the OPH support-visible GNS representation. This notation abbreviates the certified cylinder family; it is not a literal infinite-dimensional Lebesgue measure. Its support-visible continuum transfer semigroup is the corresponding Euclidean Yang–Mills semigroup.

Proof. This is the support-visible compact-gauge Yang–Mills form theorem of the Standard Model gauge paper, where it appears as the four-dimensional Euclidean Yang–Mills form theorem [1]. The proof spine is recalled here because it fixes the target Hamiltonian for the mass-gap step.

The present Yang–Mills branch takes compact simple G as branch data. When G is imported from OPH reconstruction, that import is conditional on the compact-gauge refinement receipt; on that tail the zero-obstruction transportable bosonic sector category and its faithful forgetful fiber functor reconstruct G [1]. At fixed cutoff, the declared compact-gauge patch-carrier presentation gives support-visible link holonomies and plaquette holonomies. In the refinement limit, the zero-obstruction gluing law makes infinitesimal rectangle holonomies multiplicative and path-local. The Cauchy/regularity certificate in items (Y1)–(Y3) upgrades the generalized-holonomy cluster state to a local connection A on the four-dimensional scaling chart, and the infinitesimal plaquette defect is

$$U_{\mu\nu}(\varepsilon, x) = \mathbf{1} + \varepsilon^2 F_{\mu\nu}(x) + O(\varepsilon^3), \qquad F = dA + A \wedge A.$$

The Euclideanized MaxEnt/local-Gibbs branch gives a local finite-range action density built from support-visible gauge-invariant collar data. Gauge quotienting permits only class functions of the curvature and its covariant derivatives. The four-dimensional scaling chart, Euclidean rotation invariance, locality, and reflection positivity leave one relevant dimension-four positive quadratic invariant in the pure gauge sector:

$$\langle F_{\mu\nu}, F_{\mu\nu} \rangle.$$

The possible topological density $\langle F \wedge F \rangle$ is reflection odd and belongs to a separate topological-angle sector; it is absent on the ordinary reflection-positive zero-obstruction vacuum branch used here. Higher curvature powers and covariant-derivative terms are irrelevant operators under the declared continuum scaling; their disappearance in the extracted state is part of the uniform renormalized remainder control in (Y1)–(Y3). Normalizing the unique positive quadratic invariant defines the coupling g and gives (YM).

The finite-stage cylinder measures are the gauge-register / quantum-link Gibbs measures pushed to the support-visible quotient. Proposition 9.2 proves the compatible support-visible weak-$*$ / GNS cylinder extraction. The generator and transfer-semigroup identification use the separate dynamic items (Y4)–(Y5) of Assumption 9.4; weak-* compactness alone does not provide them. □

Corollary 4.3 (Yang–Mills equations on the compact-gauge branch)**.** *With an external conserved gauge current J, the OPH curvature-square action on the same support-visible compact-gauge branch gives*

$$DF = 0, \qquad D{*}F = g^2 {*} J,$$

equivalently

$$D_\mu F^{\mu\nu} = g^2 J^\nu$$

in local coordinates after the usual continuation to the Lorentzian field-equation convention.

Proof. The curvature $F = dA + A \wedge A$ obeys the covariant Bianchi identity $DF = 0$. Varying the quadratic curvature action

$$S[A, J] = -\frac{1}{2g^2} \int \langle F \wedge {*}F \rangle + \int \langle A \wedge {*}J \rangle$$

with respect to compactly supported variations of A gives

$$\delta S = \int \langle \delta A \wedge (g^{-2} D{*}F - {*}J) \rangle$$

up to a boundary term. Stationarity gives $D{*}F = g^2{*}J$. The source-free equation is the case $J = 0$. □

Remark 4.4 (Two separate claims). The proof has two separate claims. Theorem 4.2 identifies the support-visible continuum gauge sector with four-dimensional Euclidean Yang–Mills under its branch hypotheses. The spectral argument applies to that Hamiltonian and proves a positive gap only when Assumption 9.4 supplies the OS/transfer/nontriviality bridge.

Remark 4.5 (Abelian boundary and Maxwell branch). The ordinary electromagnetic $\mathrm{U}(1)_Q$ branch is the abelian boundary case of the same compact-gauge curvature package. Restricting $F = dA + A \wedge A$ to $\mathfrak{u}(1)_Q$ gives

$$F_Q = dA_Q, \qquad dF_Q = 0, \qquad d{*}F_Q = g_Q^2{*}J_Q$$

after varying the quadratic electromagnetic action with current J_Q. This recovers Maxwell's equations after canonical electromagnetic normalization. The Clay-facing mass-gap claim concerns compact simple nonabelian G on the support-visible branch above. On the separate abelian branch, the displayed Maxwell kinetic action, an ordinary Lorentz vacuum, and the absence of a Higgs, Stückelberg, or medium mass give two transverse classical modes with quadratic pole $k^2 = 0$. The abstract group $\mathrm{U}(1)_Q$ alone does not imply that action, phase, or pole. A photon particle additionally requires a positive-energy physical quantization and a positive-residue massless spectral pole; none of those quantum receipts is part of the Clay-facing Yang–Mills gap statement.

5 Fixed-Cutoff Repair Equals Projection

Proposition 5.1 (Local exact repair equals conditional expectation)**.** *For each active collar C, the exact-Markov repair map on the support-visible quotient is the π_r-preserving conditional expectation E_C.*

Proof. On the exact-Markov branch, repair preserves exactly the repaired visible datum ρ_C, changes only complementary invisible fiber data, and acts on the quotient-first physical algebra rather than on microscopic representatives. Let Φ_C be the Heisenberg repair map and let $\mathcal{N}_C$ be the repaired local fixed algebra. Exact repair semantics require $\mathcal{N}_C$ to be exactly the ρ_C-measurable subalgebra. They give

$$\Phi_C(a) = a \quad (a \in \mathcal{N}_C),$$
$$\Phi_C(\mathcal{A}_r^{G,\mathrm{sv}}) \subseteq \mathcal{N}_C, \qquad \omega_r \circ \Phi_C = \omega_r,$$

and Φ_C is $\mathcal{N}_C$-bimodular. Therefore, for $a \in \mathcal{N}_C$ and $x \in \mathcal{A}_r^{G,\mathrm{sv}}$,

$$\omega_r(a^*\Phi_C(x)) = \omega_r(a^*x).$$

Since $\Phi_C(x) \in \mathcal{N}_C$, this equation characterizes its class in K_r as the orthogonal projection of x onto the ρ_C-measurable subspace, uniquely modulo π_r-null functions. That orthogonal projection is the π_r-preserving conditional expectation E_C. □

6 Exact Euclidean-Consensus Law

Lemma 6.1 (Fiber-homogeneous orbit condition)**.** *Fix an active collar C and a repaired value $y \in Y_{C,r}$. On the support-visible quotient, let the complete conditional fiber be*

$$F_C(y) = \rho_{C,r}^{-1}(y)$$

with conditional law $\nu_{C,y}$. If this is either finite uniform or standard atomless, has no remaining observable labels, and the primitive relaxation is invariant under every $\nu_{C,y}$-preserving automorphism, then the full fiber, including every holonomy coordinate actually resampled by repair, is homogeneous for the local repair receipt.

Proof. This is the complete-fiber homogeneity receipt. Quotienting removes declared implementation labels, but the proof must also check that no remaining source constraint or relaxation rate distinguishes points in the fiber. Under that check, the conditioned state and primitive relaxation carry the stated full measure-preserving symmetry. □

Lemma 6.2 (Scalar relaxation on a homogeneous conditional fiber)**.** *Let (F, ν) be either a finite uniform probability space or a standard atomless probability space. Let E_F be expectation onto constants, and let D_F be a bounded positive self-adjoint Markov relaxation generator such that*

$$\ker D_F = \mathrm{Ran}(E_F)$$

and D_F commutes with every measure-preserving automorphism of (F, ν). Then there is a scalar $c_F > 0$ such that

$$D_F = c_F(I - E_F).$$

Proof. For a finite uniform fiber, the symmetric-group representation on the mean-zero subspace is irreducible. For a standard atomless fiber, use equal-measure finite partitions. Automorphisms within pieces and permutations of pieces force every bounded commutant operator to be scalar on the mean-zero step functions; compatibility under refinements gives one scalar, and these step functions are dense in $L_0^2(F, \nu)$. Positivity and the kernel condition make the scalar strictly positive. The formula follows. □

Assumption 6.3 (Finite ground-state-transform and cross-fiber receipt)**.** *For every regulator, the reflection-positive finite transfer matrix supplies a unitary $U_r : \mathcal{H}_r \to K_r$, with $U_r\Omega_r = \mathbf{1}_r$, such that*

$$U_r H_r U_r^{-1} = \sum_{C \in \mathcal{C}_r} D_C.$$

Each D_C is a bounded positive self-adjoint detailed-balance Markov generator acting on the complete complementary conditional fiber, including every record or holonomy coordinate changed by repair; its fixed algebra is exactly the $\rho_{C,r}$-measurable algebra, and its fiberwise scalar $c_C(y)$ is independent of the repaired value y on the support of π_r. A Gibbs state by itself does not imply this finite transfer-matrix/decomposition receipt.

Theorem 6.4 (Exact Euclidean repair law under homogeneous fibers)**.**

Under the fiber-homogeneous orbit condition for every active collar and Assumption 6.3, there are positive constants $c_C > 0$ such that the ground-state transformed physical Euclidean generator is exactly

$$L_r^{\mathrm{rep}} = \sum_{C \in \mathcal{C}_r} c_C(I - E_C), \tag{1}$$

and therefore

$$U_r e^{-tH_r} U_r^{-1} = e^{-tL_r^{\mathrm{rep}}} \qquad (t \geq 0), \tag{2}$$

for a unitary $U_r : \mathcal{H}_r \to K_r$ *with* $U_r \Omega_r = \mathbf{1}_r$.

Proof. The finite receipt supplies the ground-state transform and its primitive local pieces. Each D_C is supported on one collar C, preserves exactly the repaired visible datum $\rho_{C,r}$, and relaxes the complete complementary conditional fiber. Hence

$$\ker D_C = \mathrm{Ran}(E_C).$$

Lemma 6.1 gives the full hidden-fiber permutation symmetry. Applying Lemma 6.2 fiberwise gives $D_{C,y} = c_C(y)(I - E_{C,y})$. The cross-fiber receipt makes $c_C(y) = c_C$, so

$$D_C = c_C(I - E_C).$$

The scalar is positive because C is active. Branch homogeneity makes it a collar-type scalar, so summing over the active collars gives (1), and exponentiation of the positive self-adjoint generator gives (2). □

Definition 6.5 (Source-defined admissible atomic collar tower)**.** *For every regulator and allowed boundary condition* b*, let the source give an atomic register set* V_r *with finite local alphabets, a finite-range Gibbs law* $\pi_{r,b}$*, and one rooted active collar* $C(v)$ *per* $v \in V_r$*, with*

$$P_{v,r,b} f := \mathbb{E}_{\pi_{r,b}}[f \mid x_{V_r \setminus \{v\}}].$$

Its source type is the isomorphism class of the rooted interaction-radius neighbourhood, boundary/sector flags, local alphabets, nonzero potential templates, complete repaired readback, conditional kernel, and normalized rate. The tower is admissible when:

(G1) *all active types over every location, allowed boundary, system size, and cofinal refinement stage belong to one printed finite set* $\mathfrak{T}_{\mathrm{act}}$*;*

(G2) $\mathcal{C}_r = \{C(v) : v \in V_r\}$*,* $E_{C(v)} = P_{v,r,b}$*, and type-equivalent collars have the same rate* $c_{v,r,b} = c_{\tau(v)} \geq c_* > 0$*;*

(G3) *if* $a_{vu}^{r,b}$ *is the supremum total-variation change of the root-*v *conditional kernel when exterior configurations differ only at* u*, outward-rounded rational bounds from the finite type table prove*

$$\sup_{r,b,v} \sum_{u \neq v} a_{vu}^{r,b} \leq \eta_* < 1. \tag{3}$$

The Dobrushin Poincaré comparison [24] then derives the uniform approximate-tensorization bound

$$\|f - \pi_{r,b}(f)\|_2^2 \leq A_* \sum_{v \in V_r} \|(I - P_{v,r,b}) f\|_2^2, \qquad A_* = (1 - \eta_*)^{-1}; \tag{AT}$$

(G4) *coarse shadow preserves the listed type transitions and conditional kernels and creates no unlisted type.*

Qualitative finite-range mixing and collar-CMI decay are not item (G3).

Proposition 6.6 (Finite collar classification and local-rate floor)**.** *The admissible collar classes are exactly the nonempty fibers of $\tau : \bigsqcup_r V_r \to \mathfrak{T}_{\mathrm{act}}$. Hence there are at most $|\mathfrak{T}_{\mathrm{act}}|$ classes and*

$$c_{v,r,b} \geq c_* := \min_{t \in \mathfrak{T}_{\mathrm{act}}} c_t > 0 \tag{4}$$

uniformly across location, boundary condition, system size, and cofinal refinement.

Proof. The type tuple is a complete source signature. Items (G1)–(G2) therefore identify precisely a finite image with a positive finite minimum, while item (G4) prevents refinement from producing a new or rate-degenerate type. □

7 Finite-Stage Gap

Theorem 7.1 (Uniform collar-projection and transfer gap)**.** *For an admissible collar tower satisfying the finite ground-state-transform receipt, set*

$$\delta_* := c_*(1 - \eta_*) = \frac{c_*}{A_*} > 0 \tag{5}$$

in the influence-certified case. Uniformly across the complete family,

$$L^{\mathrm{rep}}_{r,b} \geq \delta_*(I - P_{0,r,b}), \qquad \|e^{-tL^{\mathrm{rep}}_{r,b}} - P_{0,r,b}\|_{2\to 2} \leq e^{-t\delta_*}. \tag{6}$$

Proof. Every $P_{v,r,b}$ is an orthogonal conditional expectation. The Dobrushin comparison applied to item (G3) gives (AT), so for $f \perp \mathbf{1}$,

$$\langle f, L^{\mathrm{rep}}_{r,b} f\rangle = \sum_v c_{v,r,b}\|(I - P_{v,r,b})f\|_2^2 \geq \frac{c_*}{A_*}\|f\|_2^2.$$

This proves the operator bound, and the spectral theorem gives the transfer bound. The four uniformities follow because items (G1)–(G4) use one type table and one approximate-tensorization modulus on the entire cofinal tower. □

Proposition 7.2 (Finite countermodels for the two hypotheses)**.** *If uniform mixing is removed, the faithful two-site family, with $0 < \varepsilon \leq 1/2$,*

$$\pi_\varepsilon(00) = \pi_\varepsilon(11) = \frac{1-\varepsilon}{2}, \qquad \pi_\varepsilon(01) = \pi_\varepsilon(10) = \frac{\varepsilon}{2}$$

has exact heat-bath spectrum $\{0, 2\varepsilon, 2(1-\varepsilon), 2\}$, hence gap $2\varepsilon \to 0$. If locality is removed, put the product fair-bit law on $\{0,1\}^m$, list its 2^m states in cyclic Gray-code order, and average over the two alternating perfect matchings, with projections E_0, E_1. Product mixing is exact and separated CMI is zero, but deciding which bit to change reads the whole configuration. Exact cycle Fourier modes give

$$\mathrm{gap}\big((I - E_0) + (I - E_1)\big) = 1 - \cos(2\pi/2^m) \longrightarrow 0. \tag{7}$$

Thus both countermodels are finite, and neither hypothesis can be deleted.

Remark 7.3 (What the certificate must contain)*.* Finite-range Gibbs form and collar-CMI decay do not imply (3) and do not control projection angles. A numerical receipt must certify outward-rounded bounds $c_*^{\mathrm{lo}} > 0$ and $\eta_*^{\mathrm{hi}} < 1$, yielding the explicit certified floor $\delta_*^{\mathrm{lo}} = c_*^{\mathrm{lo}}(1 - \eta_*^{\mathrm{hi}})$. Point estimates do not qualify.

Remark 7.4 (Finite calibration boundary). The exact-rational 244-type four-dimensional Ising calibration has $c_* = 1$, $\eta_* \le 1/2$, and hence $\delta_* \ge 1/2$ for its declared finite family. The separate theorem-contract witness retains the floor 3/8. Both interfaces use one exact row validator that retains every influence and its integer multiplicity. These are checks of the supplied finite arithmetic; source-kernel completeness and rate bounds are declared premises. Both interfaces reject physical-source scope, so descriptive metadata cannot promote a calibration to a physical compact-simple-gauge receipt. A physical instance requires source collar kernels, refinement closure, gauge and zero-mode data, and independent continuum and transfer receipts. None of these data is supplied by the calibration.

8 Machine-Checked Boundaries and a Finite Gauge Diagnostic

Three finite results constrain how the receipts of Assumption 6.3 and Definition 6.5 can be discharged. Two are machine-checked theorems in the OPH Lean formalization [5]; the third is a finite numerical computation on two small interacting gauge systems, with an analytic free-limit control [6].

Noncommuting collars. The formalization contains the commuting-projection special case of Theorem 7.1, in which pairwise commuting collar projections give $L_r^{\mathrm{rep}} \ge c_*(I - P_0)$. In that Lean theorem, `thm_7_3_finite_gap`, the strictly positive collar rates are direct hypotheses. The separately machine-checked uniform-fiber lemma proves a positive scalar for its own uniform-fiber matrix datum, but no Lean theorem bridges that matrix and scalar to the collar projections and rates of the commuting theorem. The module's minimal zero-projection witness has joint fixed space $\{0\}$; on its one-point coordinate type the constant functions are the whole space. It therefore witnesses only the stripped finite operator premises and supplies neither a compact-gauge realization nor a continuum or physical mass gap.

The same formalization also contains a three-state rational countermodel: two proper symmetric idempotent Markov projections that fail to commute, whose unit-rate generator has eigenvalues 1/2 and 3/2 in place of the commuting subset sums 0, 1, and 2. Spectral accounting by subset sums does not transfer to the noncommuting collars of Definition 6.5. The Dobrushin route in Theorem 7.1 is the route that applies, and commutation is used nowhere in it. This provenance boundary does not rule out that route, the fiber-dependent-rate route below, or another receipt-complete tower.

Uniform gap and carrier growth. A second machine-checked theorem states that a refinement interface carrying a uniform bound on carrier cardinality cannot also carry a cofinal family with strict carrier growth. An admissible collar tower has strictly growing carriers along refinement, so the uniform floor (4) must be stated through the finite type table, as in Proposition 6.6, and never through a bound on carrier size.

The finite transfer receipt on $\mathbb{Z}_2$ gauge theory. Assumption 6.3 can be evaluated by finite-matrix methods on a specified finite gauge system. The accompanying float64 eigendecomposition and matrix-logarithm computation evaluates it for $\mathbb{Z}_2$ lattice gauge theory on an $L \times L$ periodic spatial torus, $L \in \{2, 3\}$, in the Gauss-law sector, with one collar per spatial link and the π_r-preserving heat bath on the fiber $\{o, X_l o\}$, where X_l flips link l and o ranges over gauge orbits.

The ground-state transform is the Doob transform by the Perron vector Ω, with $\pi_r = \Omega^2$. Two transfer objects are tested: the reflection-positive Wilson transfer matrix $T = e^{\beta_s P/2} K e^{\beta_s P/2}$, with $K(s, s') = \prod_l e^{\beta_t s_l s'_l}$, P the spatial plaquette sum, and $H = -\log(T/\lambda_{\max})$; and the Kogut–Susskind Hamiltonian $H = -\lambda \sum_l X_l - \sum_p U_p$.

For the Wilson transfer matrix the receipt holds exactly at $\beta_s = 0$, where $K = \prod_l (e^{\beta_t} I + e^{-\beta_t} X_l)$ is a constant multiple of $\exp(\frac{1}{2} \log\coth \beta_t \sum_l X_l)$ and every rate equals the dual coupling $\log\coth \beta_t$; the relative residual is below 10^{-12} for both tori. At every tested interacting point $\beta_s = \beta_t \in \{0.1, 0.3, 0.5, 0.7, 1\}$ it fails. In these finite examples $\log T$ couples configurations at multiple Hamming distances, and the best constant-rate approximant $\sum_l c_l (I - E_l)$ leaves a relative Frobenius residual of 0.038 at $\beta_s = \beta_t = 0.1$, 0.084 at 0.5, and 0.24 at 1.0 on the $L = 3$ torus.

For the Kogut–Susskind Hamiltonian the single-flip structure survives the transform exactly,

$$L^{\mathrm{rep}} = \sum_l c_l(o)\,(I - E_l), \qquad c_l(o) = \lambda(r_l(o) + r_l(o)^{-1}), \qquad r_l(o) = \frac{\Omega(o)}{\Omega(X_l o)},$$

so the scalar cross-fiber condition of Assumption 6.3 holds only where $|\log \Omega(o) - \log \Omega(X_l o)|$ has one fixed value under each flip of link l. The measured spread $\max_o c_l(o) / \min_o c_l(o)$ on the $L = 3$ torus is 1.036 at $\lambda = 2$, 1.17 at $\lambda = 1$, and 2.05 at $\lambda = 1/2$. The Dobrushin sum η_* of the single-link conditional kernels is below one, among the interacting test points, at $\beta_s = \beta_t = 0.1$, where $\eta_* = 0.59$, and at $\lambda = 2$, where $\eta_* = 0.82$; at the other tested interacting couplings item (G3) fails on this system. The free control is assessed separately.

The diagnostic fixes three facts on the two tested finite systems and coupling grid. It is exact in the free limit. The tested Wilson transfer matrices fail through multi-flip terms, and the tested local Hamiltonians fail the scalar cross-fiber clause through fiber-dependent rates. This finite grid does not exclude anisotropic couplings, alternate transfer objects, fiber-dependent positive rates, or another compact-gauge tower. A receipt on a physical compact-gauge tower may therefore admit fiber-dependent rates, replacing each scalar c_C by a positive fiber-constant function bounded below by c_*, or name a transfer object other than the Wilson matrix whose ground-state transform is a constant-rate heat-bath sum. The floor in Theorem 7.1 survives the first option without change, since its proof uses only $c_{v,r,b} \geq c_*$ and each rate function is constant on its own collar fiber; multiplication then commutes with the corresponding conditional expectation. In the tested Kogut–Susskind transform, $c_l = \lambda(r_l + r_l^{-1}) \geq 2\lambda$ by AM–GM. The variable-rate comparison is machine-checked at its finite level in the Lean module `KogutSusskindFiberRateComparison` [5]: the single-fiber rate identity $c_l = \lambda(r + r^{-1})$ with the sharp bound $c_l \geq 2\lambda$ and equality exactly at $r = 1$, and, for a finite independent product of two-point fibers with fiber-dependent rates, a product-generator spectral gap equal to the minimum fiber rate with explicit eigenstructure, hence bounded below by 2λ uniformly in every fiber ratio. Concrete two- and three-fiber instances with distinct ratios carry computed exact gaps in a released receipt tied to the finite-transfer conventions of this section. The comparison is proved on the independent fiber product, and the finite descent-and-restriction part of the gauge-orbit quotient step is machine-checked in the Lean module `GaugeOrbitQuotientGap` [5]: the $\mathbb{Z}_2$ XOR-mask action, which agrees with the gauge action of the committed non-self-loop $L = 2, 3$ transfer receipts, is formalized as a permutation subgroup whose every element is a mask translation; with fiber weights symmetric on every moved link, a condition proved equivalent to ratio one there under the $\pi = \Omega^2$ reading and exactly the equality case of the 2λ floor, the product generator commutes with every gauge translation and restricts to the gauge-invariant subspace, and every product-space Dirichlet constant survives that restriction, so the minimum fiber rate and the 2λ floor bound the quotient generator on mean-zero invariant

observables, uniformly in the unmoved-link ratios. A worked abstract two-bit global-complement instance has exact invariant-subspace constant $c_0 + c_1 = 4$ at $\lambda = 1$, strictly above the floor 2, so the bound is one-sided. This instance is not a physical one-site periodic lattice: repeated self-loop incidences cancel over $\mathbb{Z}_2$, and the finite diagnostic is defined only for $L \geq 2$. The quotient is formalized as the invariant subspace; the orbit-space carrier formulation, the committed receipt tori with several star generators, the non-product orbit law with orbit-dependent rates, and the approximate-tensorization comparison are the named residual steps, and testing the anisotropic Wilson-to-Hamiltonian limit is the other constructive route. The scalar form of Theorem 6.4 is what cross-fiber equality buys; the floor itself does not require scalar rates.

9 Continuum Extraction

Definition 9.1 (Finite compact-gauge state spaces and local cylinders)**.** *At regulator r, make the following construction both for the finite four-dimensional Euclidean slab $\Lambda_r^{(4)} = (V_r^{(4)}, E_r^{(4)})$ and for its time-zero spatial slice $\Lambda_r^{(0)} = (V_r^{(0)}, E_r^{(0)})$. In the formulas below $\Lambda_r = (V_r, E_r)$ denotes either choice until the superscript is restored. Let D_r be the finite repaired-record and zero-obstruction sector register. In the gauge-register presentation, let*

$$\widetilde{X}_r \subseteq G^{E_r} \times D_r$$

be the closed set obeying the finite Gauss, overlap, and repaired-record constraints. With $\mathcal{G}_r = G^{V_r}$ acting at edge endpoints and $\sim_{\mathrm{ov}}$ the closed relation of support-visible overlap/presentation indistinguishability, define

$$X_r := (\widetilde{X}_r/\mathcal{G}_r)/\!\sim_{\mathrm{ov}}$$

as the support-visible quotient. This is compact Hausdorff. The regulator has finitely many cells, but X_r need not be a finite set when G is a compact Lie group.

For each active collar C, let

$$\rho_{C,r} : X_r \to Y_{C,r}$$

be the continuous complete repaired readback. The complementary conditional fiber contains every record or holonomy coordinate actually resampled by repair and is either finite uniform or a standard atomless probability space with its conditioned MaxEnt/Haar law. The readback retains exactly the repaired fixed datum, not every pre-repair holonomy. A quantum-link carrier enters this proposition only through its commuting Euclidean cylinder/readout subalgebra; the noncommutative carrier algebra would require a separate GNS-representation-kernel support reduction.

For a bounded cell region $O \subset \Lambda_r$, let $\mathfrak{C}_r^{G,\mathrm{sv}}(O)$ be generated by the gauge-invariant Peter–Weyl matrix coefficients contracted into spin networks, Wilson-loop characters, and retained gauge-invariant boundary carriers in O, together with the overlap-sector projectors and continuous functions of the complete repaired collar readbacks in O. Define

$$\mathcal{A}_r^{G,\mathrm{sv}}(O) := \overline{\mathfrak{C}_r^{G,\mathrm{sv}}(O)}^{\|\cdot\|_\infty} \subseteq C(X_r).$$

Equivalently, start from the freely presented gauge-invariant readout-generator *-algebra, quotient by the kernel of its evaluation homomorphism on X_r, and complete. This kernel is a closed two-sided overlap-trivial ideal chosen before a state and is distinct from the limiting GNS null ideal below. The relation $\sim_{\mathrm{ov}}$ identifies exactly the configurations on which all declared spin-network,

sector, and repaired-readback generators agree. The resulting self-adjoint unital algebra separates points of X_r, so Stone–Weierstrass gives

$$\mathcal{A}_r^{G,\mathrm{sv}}(\Lambda_r) = C(X_r).$$

For any selected stage measure π_r, its GNS cylinder closure is therefore the full $L^2(X_r, \pi_r)$, after the finite π_r-null support reduction.

Write the two resulting systems as

$$(X_r^{(4)}, \mathcal{A}_r^{G,\mathrm{sv},(4)}(O)) \quad \textit{and} \quad (X_r^{(0)}, \mathcal{A}_{r,0}^{G,\mathrm{sv}}(B)).$$

Euclidean-history restriction gives a continuous time-zero map $\tau_{0,r} : X_r^{(4)} \to X_r^{(0)}$.

For $r \preceq s$, coarse shadow multiplies the ordered fine-edge holonomies above each coarse edge and restricts the repaired records and sector labels. Gauge covariance and refinement associativity give

$$p_{sr} : X_s \to X_r, \qquad p_{tr} = p_{sr} \circ p_{ts},$$

with the four-dimensional/time-zero commuting square

$$\tau_{0,r} \circ p_{sr}^{(4)} = p_{sr}^{(0)} \circ \tau_{0,s},$$

and unital local *-maps on the separated support-visible quotient,

$$\iota_{rs}^{O} : \mathcal{A}_r^{G,\mathrm{sv}}(O) \to \mathcal{A}_s^{G,\mathrm{sv}}(O_s), \qquad \iota_{rs}^{O}(a) = a \circ p_{sr}, \qquad \iota_{rt}^{O} = \iota_{st}^{O_s} \iota_{rs}^{O}.$$

These maps preserve local isotony, gauge invariance, repaired readbacks, and the zero-obstruction sector. When this cylinder system is used on the Standard Model gauge paper's refinement-limit gauge branch, its sector-projector restriction must match the separately certified block-multiplicity injection in the compact-gauge refinement receipt [1]; that injection is not a consequence of deterministic coarse shadow. We use a countable cofinal regulator sequence and countable bounded-region exhaustion. The unrestricted directed version uses a subnet and product compactness instead. If every declared coarse configuration has a fine extension, p_{sr} is surjective and the full pullback is injective. Otherwise the C^*-inductive limit automatically quotients the common refinement-null kernel; Proposition 9.2 does not require prelimit injectivity. That noninjective quotient branch is not, by itself, the gauge paper's receipt-certified sector ladder. Write $\mathcal{A}_r^{G,\mathrm{sv}} := \mathcal{A}_r^{G,\mathrm{sv}}(\Lambda_r)$.

Proposition 9.2 (Projective weak-* extraction and support-visible GNS gluing)**.** *Let $\mu_s^{(4)}$ be the declared finite quotient-Gibbs probability measure on $X_s^{(4)}$, let*

$$\pi_s := (\tau_{0,s})_{\#}\mu_s^{(4)}, \qquad \mu_s^{(0)} := \pi_s,$$

and, for $d \in \{4, 0\}$, define

$$\omega_s^{(d)}(a) := \int_{X_s^{(d)}} a \, d\mu_s^{(d)}, \qquad \omega_{s\downarrow r}^{(d)} := \omega_s^{(d)} \circ \iota_{rs}^{(d)}.$$

The following statements hold for both values of d; the superscript is suppressed in the proof.

(i) *for $q \preceq r \preceq s$,*

$$\omega_{s\downarrow r} \circ \iota_{qr} = \omega_{s\downarrow q};$$

(ii) *a cofinal subsequence, or subnet, has weak-* limits* $\omega_{s_\alpha\downarrow r} \to \overline{\omega}_r$ *on every fixed local cylinder, and*

$$\overline{\omega}_r = \overline{\omega}_t \circ \iota_{rt} \qquad (r \preceq t);$$

(iii) *the local limits glue to a state* ω_∞ *on*

$$\mathcal{A}_\infty^{G,\mathrm{cyl}}(O) := \overline{\varinjlim_r \mathcal{A}_r^{G,\mathrm{sv}}(O_r)},$$

and the union of these isotone local algebras is dense in the global cylinder algebra;

(iv) *after the separate support reduction*

$$\mathcal{N}_\omega(O) := \{a \in \mathcal{A}_\infty^{G,\mathrm{cyl}}(O) : \omega_\infty(a^*a) = 0\}, \qquad \mathcal{A}_\infty^{G,\mathrm{supp}}(O) := \mathcal{A}_\infty^{G,\mathrm{cyl}}(O)/\mathcal{N}_\omega(O),$$

the induced state and representation are faithful. The compatible local GNS spaces have a Hilbert direct limit $(K^{(d)}, \Pi^{(d)}, \mathbf{1}^{(d)})$*, and the union of their cylinder images is dense. Write* $K^E := K^{(4)}$ *for the Euclidean-history GNS space and* $K := K^{(0)}$ *for the time-zero repair GNS space. The latter vector becomes the physical vacuum only through the transfer/OS certificate.*

If, in a finite quotient presentation, the four-dimensional measures have declared weights

$$\mu_r^{(4)}(x) = Z_r^{-1} m_r(x) e^{-S_r(x)}$$

and satisfy the fiber-sum identity

$$\sum_{x' :\, p_{sr}(x') = x} m_s(x') e^{-S_s(x')} = \alpha_{sr} m_r(x) e^{-S_r(x)} \tag{FS}$$

with α_{sr} *independent of* x*, then* $(p_{sr}^{(4)})_\# \mu_s^{(4)} = \mu_r^{(4)}$*. The time-zero square gives* $(p_{sr}^{(0)})_\# \pi_s = \pi_r$*, so both original state families are projective. Without (FS), compactness produces a projective cluster family; it does not prove* $\overline{\omega}_r = \omega_r$ *or uniqueness of the continuum state. For continuous compact-holonomy stages, replace (FS) by the corresponding disintegration receipt* $(p_{sr}^{(4)})_\# \mu_s^{(4)} = \mu_r^{(4)}$*.*

Proof. For $q \preceq r \preceq s$, coarse-shadow functoriality gives

$$(\omega_s \circ \iota_{rs}) \circ \iota_{qr} = \omega_s \circ \iota_{qs},$$

proving (i).

For a fixed local cylinder algebra, its state space is weak-* compact by Banach–Alaoglu. The chosen local algebras are separable because compact metrizable G has a countable Peter–Weyl test algebra and a finite regulator has only finitely many record and sector generators. Their state spaces are therefore weak-* metrizable. Successively extract a subsequence for the first, second, and later cylinders, choosing the k-th diagonal index beyond the k-th regulator stage. The diagonal subsequence is cofinal and converges on all cylinders. For a general directed regulator set, use a convergent subnet in the compact product of the local state spaces; unresolved early coordinates may be filled arbitrarily because each fixed coordinate is genuinely resolved on a cofinal tail. Weak-* continuity of precomposition and part (i) yield

$$\overline{\omega}_r \circ \iota_{qr} = \lim_\alpha \omega_{s_\alpha\downarrow r} \circ \iota_{qr} = \lim_\alpha \omega_{s_\alpha\downarrow q} = \overline{\omega}_q,$$

which proves (ii).

For a representative $[a]_r$ in the algebraic inductive limit, define

$$\omega_\infty([a]_r) := \overline{\omega}_r(a).$$

Compatibility makes this well defined. Positivity and normalization are checked in one finite algebra containing the element. For every $t \succeq r$,

$$|\overline{\omega}_r(a)| = |\overline{\omega}_t(\iota_{rt}(a))| \le \|\iota_{rt}(a)\|.$$

Taking the infimum along the tail gives the inductive-limit norm bound even for noninjective bonding maps, so the state extends uniquely to the C^*-completion. This proves (iii).

On a commutative local cylinder algebra, the Riesz representation theorem identifies the null ideal with the continuous functions that vanish on the measure support. The support quotient is therefore $C(\operatorname{supp}\mu_O)$, where μ_O is the representing local Radon measure, and its induced state is faithful. If $O \subseteq O'$, state compatibility gives

$$a \in \mathcal{N}_\omega(O) \quad \Longleftrightarrow \quad \iota_{OO'}(a) \in \mathcal{N}_\omega(O').$$

Thus the inclusions descend injectively and the support quotients form an isotone faithful local net. For $r \preceq t$, define

$$V_{rt}[a]_r := [\iota_{rt}(a)]_t.$$

It is well defined and isometric because

$$\|V_{rt}[a]_r\|^2 = \overline{\omega}_t(\iota_{rt}(a^*a)) = \overline{\omega}_r(a^*a).$$

These maps compose, preserve $\mathbf{1}$, and intertwine the cylinder representations. Their Hilbert direct limit is the GNS completion of the cylinder union, proving (iv).

Finally, suppressing the $d = 4$ superscript, summing (FS) over x gives $Z_s = \alpha_{sr} Z_r$. Division by the partition functions then gives $(p_{sr}^{(4)})_\# \mu_s^{(4)} = \mu_r^{(4)}$, equivalently $\omega_s^{(4)} \circ \iota_{rs}^{(4)} = \omega_r^{(4)}$. The commuting time-zero square gives the corresponding $d = 0$ identity. □

Remark 9.3 (What compactness does not preserve). The overlap-trivial presentation quotient and the GNS support quotient are different. Even faithful finite states can converge to a nonfaithful state. Proposition 9.2 constructs the cylinder algebra, its state, the repair-side GNS space K, and the constant vacuum vector. It does not construct $\mathcal{H}$, a limit generator, a physical intertwiner, or vacuum-projection convergence. Those require the finite dynamic receipts below.

Assumption 9.4 (Four-dimensional Yang–Mills continuum certificate)**.** *In addition to the finite compact-gauge repair data, one cofinal regulator family supplies all of the following.*

(Y1) *For each bounded region O, renormalized gauge-invariant cylinder maps from one fixed countable test *-algebra $\mathfrak{C}_0^G(O)$ to $\mathcal{A}_r^{G,\mathrm{sv}}(O_r)$, with a uniform finite-volume/lattice-spacing Cauchy bound. The maps commute with the coarse-shadow ι_{rs} up to those vanishing Cauchy defects. This makes the extracted cluster family unique on the declared tests.*

(Y2) *Exact finite reflection positivity, or a positive self-adjoint transfer matrix, with reflection-compatible renormalization maps.*

(Y3) *Uniform restoration of Euclidean covariance, locality, regularity, clustering, and the declared vacuum-sector condition.*

(Y4) *Generalized Mosco convergence of both the physical transfer forms q_r^{phys} and repair forms q_r^{rep} to closed nonnegative continuum forms, or exact coherent direct-limit semigroup identities under state-preserving repair and physical refinement isometries, with one refinement-independent physical Euclidean-time normalization. In the exact alternative the isometries send $\mathbf{1}_r$ to $\mathbf{1}_s$ and Ω_r to Ω_s.*

(Y5) *Convergence of the finite intertwiners U_r and their inverses on dense form cores to a unitary U, together with $U_r\Omega_r = \mathbf{1}_r$ and strong convergence of $P_{0,r} = |\mathbf{1}_r\rangle\langle\mathbf{1}_r|$ to $P_0^K = |\mathbf{1}\rangle\langle\mathbf{1}|$ under the declared Hilbert identifications. The physical vacuum projection is $P_0^{\mathcal{H}} := U^{-1}P_0^K U = |\Omega\rangle\langle\Omega|$. On the dense time-zero cylinder core, the finite Markov/OS identity*

$$\omega_r^{(4)}\Big((\iota_{0,r}f)^*\tau_t^{(r)}(\iota_{0,r}g)\Big) = \langle f, e^{-tL_r^{\mathrm{rep}}}g\rangle_{K_r} = \langle U_r^{-1}f, e^{-tH_r}U_r^{-1}g\rangle_{\mathcal{H}_r}$$

holds and passes to the limit. Thus the OS-reconstructed time-zero Hilbert space and Hamiltonian are the same $(\mathcal{H}, H)$ as the transfer-form limit, not a second pair with reused notation.

(Y6) *A nontriviality certificate: renormalized local observables A_r, all images of one fixed bounded cylinder test under the maps in (Y1), whose centered vectors converge under the Hilbert identifications and satisfy*

$$\inf_r \mathrm{Var}_{\omega_r}(A_r) > 0, \qquad \sup_r q_r^{\mathrm{phys}}(A_r\Omega_r) < \infty.$$

The explicit multiresolution reference branch of the Standard Model gauge paper [1], with its fixed-cutoff regulator certificate in the screen microphysics paper [3], supplies a model for items (Y1)–(Y2) and an exact transfer tower when its shell identities are implemented. Identification of its continuum limit with the four-dimensional Yang–Mills OS theory is the additional content of (Y3)–(Y6).

Theorem 9.5 (Conditional continuum transfer identification)**.** *Under Assumption 9.4, the generalized Mosco limit, or the exact coherent direct limit in item (Y4), produces nonnegative self-adjoint generators H and L^{rep} and a unitary U satisfying*

$$Ue^{-tH}U^{-1} = e^{-tL^{\mathrm{rep}}} \qquad (t \geq 0), \tag{8}$$

and hence

$$UHU^{-1} = L^{\mathrm{rep}}. \tag{9}$$

If the finite repair generators satisfy

$$L_r^{\mathrm{rep}} \geq \delta_*(I - P_{0,r})$$

with one $\delta_* > 0$ and the vacuum projections converge strongly, then

$$H \geq \delta_*(I - P_0^{\mathcal{H}}).$$

Proof. Generalized Mosco convergence is equivalent to generalized strong-resolvent and transfer-semigroup convergence on the declared Hilbert identifications. In the exact alternative, the coherent finite contraction semigroups define strongly continuous direct-limit semigroups on the dense cylinder images, and uniform contractivity extends them to the completed Hilbert spaces. Their

bounded operators are symmetric and positive on the dense finite-stage union, hence self-adjoint and positive after extension. Convergence of U_r and U_r^{-1} on dense form cores passes the finite-stage intertwining relation to the limit, so uniqueness of self-adjoint semigroup generators gives $UHU^{-1} = L^{\mathrm{rep}}$. For a repair-form Mosco recovery sequence $\psi_r \to \psi$,

$$q_r^{\mathrm{rep}}(\psi_r) \geq \delta_*(\|\psi_r\|^2 - \|P_{0,r}\psi_r\|^2).$$

Taking the limit and using strong convergence of the vacuum projections gives the continuum form inequality. In the exact direct-limit alternative, the finite bound is equivalently

$$\|e^{-tL_r^{\mathrm{rep}}}(I - P_{0,r})\| \leq e^{-\delta_* t};$$

coherence passes this estimate to the direct-limit semigroup, and the spectral theorem gives the same continuum form inequality. □

Theorem 9.6 (Spectral exclusion from local Euclidean correlations)**.** *Let a reconstructed local theory have a nonnegative self-adjoint Hamiltonian H, normalized vacuum Ω, and strongly continuous time translations, with $H\Omega = 0$. Suppose the centered local vectors $v_A = (A - \langle\Omega, A\Omega\rangle I)\Omega$ have dense linear span in $\Omega^\perp$. At each finite regulator use one declared source, ground state, and physical time normalization*

$$K_r = s_r(H_r^{\mathrm{source}} - E_{0,r}I),$$

where $s_r > 0$ specifies source time units per physical time unit before the comparison. Compute finite Euclidean correlations using e^{-tK_r}. For every fixed local test A, suppose its centered Euclidean correlations satisfy, for fixed $h > 0$, $\delta > 0$, and every integer $k \geq 0$,

$$C_{A,r}(kh) \longrightarrow C_A(kh) := \langle v_A, e^{-khH}v_A\rangle, \qquad 0 \leq C_{A,r}(kh) \leq B_A e^{-\delta kh} + \varepsilon_{A,r}(k),$$

where $B_A < \infty$ is independent of the regulator and $\varepsilon_{A,r}(k) \to 0$ for each fixed k. All implementation and record errors are included in $\varepsilon_{A,r}(k)$. Then

$$H \geq \delta(I - |\Omega\rangle\langle\Omega|).$$

The vacuum is unique in this cyclic representation. No equality of finite and continuum spectra is assumed.

Proof. First take the regulator limit at fixed k, giving $C_A(kh) \leq B_A e^{-\delta kh}$. If μ_A is the positive spectral measure of H in v_A, then for $0 \leq E < \delta$,

$$\mu_A([0, E]) \leq e^{khE}C_A(kh) \leq B_A e^{-kh(\delta - E)} \longrightarrow 0.$$

Thus $\mathbf{1}_{[0,E]}(H)$ annihilates every centered local vector. Boundedness of this projection extends the conclusion to $\Omega^\perp$; it acts as the identity on Ω. Increasing E to δ proves the form inequality. The regulator limit precedes the large-time limit, so no uniform error bound at arbitrarily large time is inferred. Exact finite transfer controls and the full criterion are given in [7]. □

Corollary 9.7 (A finite threshold from correlation noncollapse)**.** *Under Theorem 9.6, suppose one fixed local test also has correlation convergence at times $t_j \downarrow 0$, a finite-stage variance floor $\|v_{A,r}\|^2 \geq v_0 > 0$, and a uniform physical energy bound $\langle v_{A,r}, K_r v_{A,r}\rangle \leq M < \infty$, in that same clock. Then*

$$0 < \delta \leq \inf(\mathrm{Spec}(H) \setminus \{0\}) \leq M/v_0 < \infty.$$

Proof. Time-zero convergence preserves the variance floor. The elementary bound $1 - e^{-tE} \le tE$ gives

$$0 \le \frac{C_{A,r}(0) - C_{A,r}(t_j)}{t_j} \le M.$$

Pass to the regulator limit at fixed j. The spectral functions $(1 - e^{-t_j E})/t_j$ increase to E, so monotone convergence gives $\|H^{1/2} v_A\|^2 \le M$. Its nonzero spectral measure has mass at least v_0, hence mean energy at most M/v_0. The nonzero spectral threshold cannot exceed that mean. □

These results are an alternative route to a gap of the reconstructed physical Hamiltonian. Their correlation-identification premise requires the same local theory and time translations as the reconstruction. They establish neither the existence of that theory nor the equality $\Delta_{\mathrm{YM}} = \Delta_{\mathrm{rep}}$, which retains the unitary-intertwining hypotheses of Theorem 9.5. A uniform finite gap is sufficient for the decay bound, but is not necessary: low-energy prelimit modes whose original weights vanish in every limiting local test can disappear from the cyclic continuum representation. Conversely, decay for a single fast probe does not exclude slower spectrum outside its generated subspace.

10 Axiomatic Reconstruction and Nontriviality

Theorem 10.1 (Conditional Osterwalder–Schrader reconstruction on the support-visible compact–gauge branch)**.** *Under Assumptions 4.1 and 9.4, the continuum support-visible compact-gauge cylinder family is Euclidean invariant, reflection positive, regular on gauge-invariant local cylinder observables, nontrivial, and cyclic for the vacuum sector. Hence Osterwalder–Schrader reconstruction gives a four-dimensional quantum Yang–Mills theory*

$$(\mathcal{H}, \Omega, H, \mathcal{A}^G_{\mathrm{loc}})$$

on the support-visible gauge-invariant local algebra, with $H \ge 0$ and e^{-tH} equal to the certified continuum Euclidean transfer semigroup.

Proof. Euclidean covariance, locality, reflection positivity, regularity, clustering, transfer convergence, and nontriviality are precisely the certificate items in Assumption 9.4. Proposition 9.2 supplies the four-dimensional Euclidean cylinder state K^E and its time-zero GNS restriction K; the certificate supplies the missing regularity and transfer controls. The finite/limiting time-zero identity in (Y5) identifies the OS time-zero inner product and translation semigroup with the transfer-form limit $(\mathcal{H}, H)$. The standard Osterwalder–Schrader reconstruction theorem therefore produces the Hilbert space, vacuum, local algebra, and positive Hamiltonian, and identifies the physical time-translation semigroup with the certified Euclidean transfer semigroup [20, 21]. □

Proposition 10.2 (Nontriviality of the support-visible compact-gauge theory)**.** *Under Assumption 9.4, the support-visible compact-gauge local algebra on the zero-obstruction vacuum branch strictly contains the vacuum scalars and admits a non-vacuum finite-energy local excitation.*

Proof. The compact-gauge witness theorem of the Standard Model gauge paper supplies finite support-visible gauge-invariant local observables on the receipt-conditional compact-gauge branch [1]. The executable observer's finite gauge example is a U(1) code, and the separate sixty-state native link design has exact local non-Abelian Gauss constraints with raising-operator links; neither supplies these Wilson cylinder observables or non-Abelian Yang–Mills dynamics [16, 9]. At

finite cutoff the witness gives a support-visible gauge-invariant local observable, such as a nonconstant Wilson/plaquette cylinder observable, whose vacuum variance is positive. Its GNS vector is orthogonal to the vacuum after subtracting its expectation value. The finite-range local-Gibbs generator assigns finite energy to finite-cylinder excitations. The continuum nontriviality step is not a consequence of weak-$*$ state convergence alone; it is the variance-floor and finite-energy item (Y6) in Assumption 9.4. With that certificate, the local cylinder vectors survive in the support-visible continuum, so the continuum local algebra is strictly larger than $\mathbb{C}I$ and contains non-vacuum finite-energy local excitations. □

11 Main Theorem

Theorem 11.1 (Conditional positive four-dimensional compact-gauge Yang–Mills mass gap)**.** *Let G be a compact simple gauge group carried by a support-visible compact-gauge OPH vacuum branch satisfying the standing setup above and the continuum certificate of Assumption 9.4, the finite transfer receipt of Assumption 6.3, and all hypotheses of Theorem 6.4, Proposition 6.6, and Proposition 7.1. The theory reconstructed in Theorem 10.1 is nontrivial, and its continuum support-visible Hamiltonian H satisfies*

$$H \geq \delta_*(I - P_0^{\mathcal{H}}), \tag{10}$$

where $P_0^{\mathcal{H}}$ projects onto the physical vacuum. Therefore

$$\mathrm{Spec}(H) \cap (0, \delta_*) = \varnothing, \qquad \Delta_{\mathrm{YM}} \geq \delta_* > 0. \tag{11}$$

On that certified continuum branch the repair gap and Yang–Mills gap are exactly equal:

$$\Delta_{\mathrm{YM}} = \Delta_{\mathrm{rep}}. \tag{12}$$

Proof. At every finite stage, Proposition 7.1 gives

$$L_r^{\mathrm{rep}} \geq \delta_*(I - P_{0,r}).$$

By Theorem 9.5, on the extracted cylinder/GNS space, this lower bound passes to the support-visible continuum:

$$L^{\mathrm{rep}} \geq \delta_*(I - P_0^K).$$

Using (9), $UHU^{-1} = L^{\mathrm{rep}}$. Conjugating the lower bound by U^{-1} gives (10), and the spectral statement (11) follows. Since unitary equivalence preserves the nonzero spectrum and identifies the vacuum with the constant sector, the infimum of the nonzero spectrum is the same on both sides, giving (12). □

Remark 11.2 (Group-uniform form). The proof is group-uniform. Once a compact simple G is carried by a compact-gauge zero-obstruction OPH branch, no step uses special properties of the realized Standard Model quotient. The inputs are compact-gauge support-visible quotient locality, exact local repair on collars, the finite ground-state-transform/cross-fiber receipt, the source-type and uniform approximate-tensorization receipts, refinement coherence, projective cylinder extraction, the transfer/vacuum certificate, and the OS regularity/noncollapse certificate.

12 Exact Gap Accounting

On a branch satisfying Assumption 9.4, the proof gives more than a positive lower bound. It identifies the Hamiltonian whose gap is being measured:

$$H = U^{-1} L^{\mathrm{rep}} U.$$

Therefore

$$\mathrm{Spec}(H) \setminus \{0\} = \mathrm{Spec}(L^{\mathrm{rep}}) \setminus \{0\}.$$

The Yang–Mills gap is exactly the infimum of the nonzero repair spectrum:

$$\Delta_{\mathrm{YM}} := \inf\big(\mathrm{Spec}(H) \setminus \{0\}\big) = \inf\big(\mathrm{Spec}(L^{\mathrm{rep}}) \setminus \{0\}\big) =: \Delta_{\mathrm{rep}}.$$

The finite-stage approximate-tensorization argument proves $\Delta_{\mathrm{rep}} \geq \delta_* > 0$. The exact accounting statement is the conditional equality $\Delta_{\mathrm{YM}} = \Delta_{\mathrm{rep}}$; the inequality is the positivity proof for that same quantity. Without the continuum certificate, the same equations are finite-regulator repair accounting instead of a Clay-admissible Yang–Mills theorem.

13 Relation to the 2D Yang–Mills Bridge

The companion OPH results also contain an exact edge-sector identity:

$$Z_{\mathrm{edge}}(t) = \sum_R d_R^2 e^{-tC_2(R)} = K_t(1),$$

which is the compact-group heat kernel at the identity and matches the standard two-dimensional Yang–Mills heat-kernel partition form. Peter–Weyl supplies the heat-kernel identity, and Gross–Taylor gives the standard large-N worldsheet dictionary when a separate large-N_{edge} branch with remainder control is declared [22, 23].

That 2D result is a normalization and worldsheet-effective bridge. The spectral theorem above concerns the support-visible compact-gauge Hamiltonian and obtains its lower bound by identifying Euclidean transfer with the repair generator.

14 A full Haar-link finite Yang–Mills source

On the full gauge-invariant Haar link space of an open three-dimensional SU(2) cube, the positive Hamiltonian $H = \alpha \sum_l C_l + \beta \sum_p (1 - \mathrm{Re}\,\mathrm{Tr}\, U_p/2)$ has a unique vacuum. The complete electric physical gap is 3α. At $g^2 = 8$, spacing one and eight vertices, min–max and the constant-state residual give an interacting vacuum gap between $21/2$ and $1024/47$ in source units. A local plaquette has vacuum variance at least $47/256$ and unnormalized centered-vector energy form at most 4. Every link representation is retained. The lower bound deteriorates with volume and both energy bounds scale as the inverse spacing; no four-dimensional continuum mass gap follows. The source, domains and quantitative certificates are given in `model/openwave_pure_su2_source`.

The same pure-gauge source admits $T_h = e^{-hV/2} e^{-hE} e^{-hV/2}$ on the full compact-group link space. Every electric character coefficient is positive, and reflected cylinder histories are Gram forms of half-history vectors. Gauge projection commutes with the transfer. The strong product limit gives

$(T_{t/n})^n \to e^{-tH}$ in the declared Hamiltonian clock. Complete character-space controls verify this construction at three time steps. This establishes finite-regulator reflection positivity; it supplies no joint spatial continuum limit or restoration of Euclidean covariance. The source, domains and quantitative certificates are given in `model/openwave_su2_transfer`.

For a pure SU(2) plaquette, exact Gauss reduction gives the full character basis χ_n, $n \geq 0$, and Jacobi entries $H_{nn} = \alpha n(n+2) + \beta$, $H_{n,n+1} = -\beta/2$. Rational Schur bounds on the infinite tail and Sturm counts enclose four complete-source energy levels to width below 2^{-40}, at four declared couplings. Local curvature has a certified positive variance and finite energy, with controlled Euclidean correlation records. The representation cutoff is removed in these bounds; spatial volume and lattice-spacing limits are separate requirements. The source, domains and quantitative certificates are given in `model/openwave_su2_spectral`.

This construction instantiates the finite physical-vacuum and local-noncollapse interfaces without identifying its finite graph with the continuum branch. Uniform volume control, renormalized four-dimensional Schwinger functions, covariance, locality and the continuum transfer remain hypotheses of that branch.

15 Physical scope

A Clay-level conclusion requires the finite transfer receipt and a uniform positive rate and approximate-tensorization floor, followed by the Schwinger-function, reflection-positivity, covariance, locality, nontriviality and transfer/intertwiner convergence requirements in Assumption 9.4. The projective weak-* / GNS extraction alone does not supply them. Under all those premises the same Hamiltonian satisfies

$$\Delta_{\mathrm{YM}} = \Delta_{\mathrm{rep}} \geq c_*/A_* > 0.$$

This is a conditional statement for pure Yang–Mills with compact simple G. The full theory with an unbroken electromagnetic sector and a massless photon has no positive global vacuum gap. Its sector gaps must specify the state, sector and limiting procedure.

The coordinate-field construction used in the finite observer controls has sixty fermionic modes per cell and a fifteen-dimensional product-group representation [8]. Choosing a compact simple factor of its Lie algebra does not exhibit a reducing Hilbert sector of its coupled Wilson action. Fixing the other quantum link variables to the identity is not normalizable in Haar L^2, and projecting them to constant words need not preserve plaquette multiplication. A pure Yang–Mills branch therefore requires a separately declared source or a proved decoupling map, with its own vacuum and clock.

That finite source has spacing $a_n = 2^{-n}$, side $L_n = 2^{2n+1}$, per-link word degree $\ell_n = n + 1$, and fixed electric and magnetic coefficients 105 and 1/7. Every Hamiltonian term scales as a_n^{-1}, so

$$H_n = a_n^{-1} \widehat{H}(L_n, \ell_n; 105, 1/7).$$

Its physical-sector gap in the declared clock is $\widehat{\Delta}_n/a_n$. If the uniform finite-gap route and a noncollapsing finite-energy local witness both apply, they require

$$a_n \delta \leq \widehat{\Delta}_n \leq a_n M/v_0.$$

A constant positive dimensionless gap would instead send the physical threshold to infinity and contradict that witness. Neither a positive finite gap nor a fitted change of clock proves a finite

continuum mass. The fixed coefficients supply no running-coupling or asymptotically free short-distance theorem [7].

Its stationary polynomial preparation is also distinct from the exact vacua in Theorem 9.6. For example, $H = \mathrm{diag}(0, 1)$, $\rho = \mathrm{diag}(1 - \epsilon, \epsilon)$, and $A = |0\rangle\langle 1|$ give

$$\mathrm{Tr}\Big(\rho A^* e^{-tH} A e^{tH}\Big) = \epsilon e^t.$$

The ground-energy excess is ϵ, yet this ordered correlation contains a negative energy transfer. Ground-energy excess alone therefore supplies neither a vacuum correlation nor control of exponential moments as spectral widths grow. Prepared approximants require their own errors vanishing at each fixed Euclidean time; their source and retained reference cannot be replaced by a different vacuum during the limit.

References

[1] B. Müller, M. Poneder, A. Osika, K. Xue, P. Nguyen, D. Matscheko, and D. A. Friedman, *Standard Model Gauge Structure under Complete-Response and Observer-Transport Premises*, 2026.
Source: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/paper/deriving_standard_model_gauge_structure_from_observer_overlap_consistency.tex.
PDF: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/paper/deriving_standard_model_gauge_structure_from_observer_overlap_consistency.pdf.

[2] B. Müller, K. Xue, J. Kim, K. A. Anirudha, D. Matscheko, and J. Hill, *Reality as a Consensus Protocol: The Fixed-Point Computation That Implements Physics*, 2026.
Source: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/paper/reality_as_consensus_protocol.tex.
PDF: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/paper/reality_as_consensus_protocol.pdf.

[3] B. Müller, A. Osika, K. Xue, B. Cassie, B. de La Fournière, and D. A. Friedman, *Twelve-Port Icosahedral Carriers and Native Seam Repair: The Carrier Layer of Observer Patch Holography*, 2026.
Source: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/paper/screen_microphysics_and_observer_synchronization.tex.
PDF: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/paper/screen_microphysics_and_observer_synchronization.pdf.

[4] B. Müller, *OPH-FPE: finite simulator and receipt engine for Observer-Patch Holography physics experiments*, 2026.
Source: https://github.com/muellerberndt/oph-physics-sim.
Finite local-domain revision and data: https://github.com/muellerberndt/oph-physics-sim/tree/550fe77dc67dc8bef2dd8927bcf72ec98cfa3506, https://github.com/muellerberndt/oph-physics-sim/tree/550fe77dc67dc8bef2dd8927bcf72ec98cfa3506/data/local_domain.
Canonical evidence archives: https://github.com/FloatingPragma/observer-patch-h

ography/tree/r2065/evidence, including the exact local-domain package and checker at https://github.com/FloatingPragma/observer-patch-holography/tree/r2065/evidence/local_domain.

[5] B. Müller et al., *Observer Patch Holography: machine-checked Lean formalization*, separate modules for the uniform-fiber lemma, commuting-projection inequality, finite-gap assembly and stripped witness, conditional continuum chain, and cofinal spectral-tail boundary, 2026. https://github.com/FloatingPragma/observer-patch-holography/tree/r2065/Lean. Modules `KogutSusskindFiberRateComparison` and `GaugeOrbitQuotientGap`: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/Lean/Screen/KogutSusskindFiberRateComparison.lean, https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/Lean/Screen/GaugeOrbitQuotientGap.lean.

[6] B. Müller, *Finite $\mathbb{Z}_2$ lattice-gauge diagnostic for the ground-state-transform receipt*, 2026. https://github.com/FloatingPragma/observer-patch-holography/tree/r2065/code/yang_mills.

[7] FloatingPragma, *Euclidean decay, the physical clock and a continuum spectral gap.* Proof: `model/particle_yang_mills_gap_contract/PROOF.md`. Exact finite transfer controls: `model/particle_yang_mills_gap_contract/transfer.py`.

[8] FloatingPragma, *One declared source, state, reference and clock.* `model/particle_common_contract/PROOF.md`.

[9] FloatingPragma, *The runnable OPH model*, 2026. https://github.com/FloatingPragma/observer-patch-holography/tree/r2065/model.

[10] FloatingPragma, *What propagates on a federation of identical elementary carriers*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/federated_observer/SCALING.md; receipt: https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/federated_observer/results.json

[11] FloatingPragma, *A local action for native quantum observations*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/native_action/README.md.

[12] FloatingPragma, *Quantum mixing of the native C/E repairs*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/observer/QUANTUM_MIXING.md.

[13] FloatingPragma, *A source-executed canonical scalar field regulator*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/relativistic_scalar/README.md.

[14] FloatingPragma, *A spatial fluctuation field from one native history*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/physical_space_time/OCCUPATION_FIELD.md.

[15] FloatingPragma, *Joint time and field laws on matched source screens*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/observer_time_fields/REFINEMENT.md.

[16] FloatingPragma, *OPH reverse-engineering table: model and theorem premises*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2065/model/PREMISES.md.

[17] Clay Mathematics Institute, "Yang–Mills & the Mass Gap." https://www.claymath.org/millennium/yang-mills-the-maths-gap/.

[18] A. Jaffe and E. Witten, "Quantum Yang–Mills Theory," official Clay Mathematics Institute problem description. https://www.claymath.org/wp-content/uploads/2022/06/yangmills.pdf.

[19] J. Carlson, A. Jaffe, and A. Wiles (eds.), *The Millennium Prize Problems*, Clay Mathematics Institute and American Mathematical Society. https://www.claymath.org/library/monographs/MPPc.pdf.

[20] K. Osterwalder and R. Schrader, "Axioms for Euclidean Green's functions," *Communications in Mathematical Physics* **31** (1973), 83–112.

[21] K. Osterwalder and R. Schrader, "Axioms for Euclidean Green's functions II," *Communications in Mathematical Physics* **42** (1975), 281–305.

[22] F. Peter and H. Weyl, "Die Vollständigkeit der primitiven Darstellungen einer geschlossenen kontinuierlichen Gruppe," *Mathematische Annalen* **97** (1927), 737–755.

[23] D. J. Gross and W. Taylor, "Two-dimensional QCD is a string theory," *Nucl. Phys. B* **400** (1993), 181–208, arXiv:hep-th/9301068.

[24] L. Wu, "Poincaré and transportation inequalities for Gibbs measures under the Dobrushin uniqueness condition," *Annals of Probability* **34** (2006), 1960–1989, arXiv:math/0611635.

AI Assistance Disclosure

This research project used research-grade commercial models, including Anthropic's Fable and OpenAI's GPT-6 Astra, for research support, software development, editing, and synthesis. The authors are responsible for the paper's claims, methods, and final text.