The de Sitter Time-Advance Sign from a Finite Screen with Fixed Capacity

Bernhard Mueller*

November 2, 2026

Paper release: r2060 **Released:** November 2, 2026

Author affiliation: Bernhard Mueller, Pragma Research Inc.

Abstract

On a finite screen with fixed capacity, maximizing generalized entropy over sector probabilities gives $S_{\rm gen}^{\max} = \log M$, and uniform transfer of a fraction f of the horizon capacity to an observer changes both this maximum and the logarithmic sector observable $\mathcal{A}$ by $\log(1-f) < 0$. The unperturbed horizon is therefore a one-sided boundary maximum along the transfer direction. The exact gradient and Hessian of $\mathcal{A}$ show that this statement cannot be replaced by an interior-maximum claim for that observable: at fixed horizon capacity its tangent Hessian is positive, while its homogeneous curvature is negative, and the joint counting entropy is a nonstrict maximum. Under a horizon-capacity dictionary this fixed-budget law supplies a conditional sign mechanism for the time advance and increasing regularized out-of-time-ordered correlator of negative de Sitter shocks, a sign that conflicts with the positivity and cyclicity required by a proposed static-patch trace. For an icosahedral screen, a normalized finite shock spectrum follows conditionally from exact gauge transport of the rotation triplet and a nearest-neighbour Laplacian identification, with the pure-de-Sitter normalization and the line-graph spectrum identity exact. The calculation does not derive the coefficient or gravitational attachment and supplies no static-patch trace.

Keywords: de Sitter horizon, shock wave, static patch, finite screen, generalized entropy, icosahedral graph

One theory and its executable hypothesis. OPH proposes closure: one complete history is both a timeless self-referential structure and its own internal computation, and observers inside it recover and realize its specification. The hypothesis that makes this proposal concrete and testable is executable.

> *A simple, fully specified local repair and memory law on the proposed carrier architecture admits continuing observers whose compatible records and interventions have the observed* $3+1$*-dimensional quantum-field description and a common classical-gravity*

*Corresponding author: bernhard@floatingpragma.ai

limit. The preferred matter mechanism is localized topological defects of native overlap transport, with rest energy derived from their excess maintenance dynamics in the same physical clock and energy dictionary.

One source process must meet every part of this statement: records, clocks, matter, energy and responsive geometry come from the same law, state and refinement. Coherence, protected codes, radial directions, fields, clocks and couplings may be specified as part of its design; their implementation and observable consequences must be established on that process. A failed specified law rejects that law and its tested interpretation; it rejects the wider proposal only when the rejected choice is proved necessary.

The tower. The papers form one tower over the executable model. Each layer states a hypothesis and proves finite theorems under displayed premises. A conclusion passes to the next layer only after that premise is proved on the same model process; until then it is a conditional implication.

0. **Closure and agreement.** One history reads itself into existence, and agreement on shared records defines the public world. This is the interpretation and acceptance criterion; no theorem selects a unique universe.
1. **Repair mathematics.** Local agreement repair reaches global consistency, and nonzero cycle holonomy is the residue that local repair cannot remove. Fixed-point, normal-form, frustration and finite event-algebra theorems hold on their stated domains.
2. **Carrier and repair law.** Federated twelve-port icosahedral carriers with the native C/E seam repair are the substrate. The seam rule follows from four stated requirements, the quantum record laws are classified, and the uniform native repair selects a rank-three slow-response band.
3. **Executable observer.** Observers are continuing native processes that prepare, read, retain and act on records. Finite complete-output executions, overlapping-observer correspondence, event clocks and checkpoint continuation are verified; one continuing observer with every service and resource accounted is the open join.
4. **Clocks and fields.** Modular flow of observer algebras supplies candidate local clocks, and native repair supplies a positive continuum generator of spectral dimension two on the screen. A declared radial extension gives a conditional smooth $1+3$ scalar description. Complete finite source factors join spatial CAR dynamics, particle creation, repair and a common calibrated clock. Physical validation of that clock and energy normalization, and an interacting chiral $3+1$ limit, remain open.
5. **Gauge structure and matter.** Gauge structure is the transport of overlap data, and stable massive matter is a localized defect whose boundary transport carries a holonomy that admitted repairs cannot erase. The compact Lie type $\mathfrak{u}(1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(3)$ follows under complete-response premises, and fixed nonzero holonomy forces positive residual mismatch. A specified coherent ordering memory protects native exchange-history sectors, and a colored extension implements their fermionic exchange. A localized physical defect, its maintenance-to-mass law and a predicted measured spectrum remain unestablished; the earlier double-load candidate is transient.

6. **Gravity.** Einstein dynamics is the coarse response of the same process. The null-balance and tensor-completion implication is proved under its premises; realizing those premises on one refinement tower is open, and the static radial product fails null balance.

7. **Cosmology and tests.** The same process must fix the vacuum, dark sector, primordial response and black-hole records, and pass independent prospective tests. The cosmology papers and the fine-structure, Koide, de Sitter and Yang–Mills supplements carry conditional branches, diagnostics and retrospective comparisons at this layer.

Completion and stopping. The aim is a consistent and complete theory: every layer realized on one process, or a proved obstruction that removes OPH or a stated branch of it. Results that no join needs leave the active chain with their provenance. A universal obstruction over a frozen admissible class, an unavoidable excluded prediction, or a contradiction among indispensable premises stops that class. A failed candidate, an unconstructed map or a resource stop has a narrower meaning. The ordered joins, retained foundations and stopping rules are recorded with the executable model in the public repository.[1]

Fixed physical boundaries. Repair counts, host runtime and entropy production are distinct from energy; the maintenance-to-energy map must be derived with one common signal speed and no per-species conversion. The modular parameter is dimensionless until a specified and implemented clock calibration relates it to duration. Stationary repair or record entropy is distinct from the fixed-volume generalized-entropy variation that the Einstein construction consumes. A massless photon makes the infinite-volume theory gapless, so sector gaps and thresholds name their state and limit; confinement precludes free massless gluons as observable particles; a quantum graviton lies outside the classical Einstein theorem.

This paper's hypothesis and place in the tower

This paper is a supplement at layers six and seven. Its hypothesis is that the screen of a de Sitter observer is a finite screen of fixed total capacity, shared between the horizon and the observer, with the horizon area read as the logarithmic sector observable. Under that hypothesis the sign of the negative de Sitter shock, and the dark-energy equation of state on the fixed branch, follow from finite capacity transfer.

The finite results hold unconditionally on their stated domains: the pure de Sitter normalization $\mu^2 = \lambda_{\ell=1}$, the entropy maximum $\log M$, the uniform-transfer law $\log(1-f)$, the analytic gradient and Hessian of the sector observable, and the regular line-graph spectrum identity. The time-advance sign, the icosahedral shock spectrum and the equation-of-state statements are conditional implications of the physical dictionary and of the two discrete-shock assumptions stated where they are used.

For the conclusion to apply, the executable model must realize from one native process a closed screen-capacity ledger shared by horizon and observer, an area map that turns the whole-screen count into $A_{\mathrm{horizon}}/4G$ with a derived coefficient, a map from transferred capacity to the observer mass of the gravitational shock calculation, a dynamical generator that changes sector dimensions,

[1] https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/claims/completion_program.json

and the gauge transport and kinetic operator of the discrete shock. These are open joins. An established exclusion of $(w_0, w_a) = (-1, 0)$ rejects the fixed-capacity branch, and an established phantom equation of state $w < -1$ rejects the monotone-capacity branch. A proof that the native whole-screen count admits no area map with a nonzero universal coefficient would remove the gravitational reading while leaving the finite identities intact.

1 The sign problem

Chen, Stanford, Tang, and Yang study a static-patch proposal in which a Euclidean gravitational path integral with an observer worldline computes a Hilbert-space trace [1]. Their shock-wave calculation includes observer recoil and gravitational backreaction. The resulting out-of-time-ordered correlator conflicts with cyclicity and positivity: its leading regularized contribution increases. Geometrically, the negative shock advances the signal in the horizon null coordinate.

The finite-screen calculation below isolates a capacity mechanism for this sign. Three levels are kept separate throughout:

1. exact entropy and graph identities for a finite sector model;
2. an analytic relaxation of integer sector dimensions for stating gradients and curvatures;
3. the physical dictionary identifying screen capacity transfer with a gravitational horizon shock.

The first two levels are mathematical statements. The third is a conditional attachment. In particular, a finite capacity premise does not evade the trace obstruction in Ref. [1]. Finite-dimensionality is part of the premise tested there.

2 The smooth de Sitter normalization

Write the horizon shock equation in spacetime dimension d as

$$\left(-\nabla^2_{S^{d-2}} - \mu^2\right) X^+ = 8\pi G\, T^-, \qquad \mu^2 = \frac{d-2}{2}|f'(r_c)|r_c = (d-2)\kappa r_c. \tag{1}$$

For pure de Sitter space of radius L,

$$f(r) = 1 - \frac{r^2}{L^2}, \qquad r_c = L, \qquad \kappa = \frac{1}{L}.$$

Lemma 2.1 (Radius-independent shock mass)**.** *For pure de Sitter space,*

$$\mu^2 = d - 2 = \lambda_{\ell=1}\left(S^{d-2}\right).$$

Consequently, μ^2 is independent of L and of the cosmological constant.

Proof. The product κr_c equals one. Equation (1) therefore gives $\mu^2 = d - 2$. Scalar spherical harmonics on S^{d-2} have

$$\lambda_\ell = \ell(\ell + d - 3),$$

so $\lambda_1 = d - 2$. □

The smooth $\ell = 1$ shock is generated by a de Sitter isometry and is pure gauge, with the corresponding source component removed by the gravitational constraint [1]. This explains the zero of the operator in Eq. (1). Transferring that normalization to a discrete screen requires a separate premise, stated in Section 4.

3 Finite entropy and the capacity-transfer direction

3.1 The exact entropy maximum

Let a finite screen have positive integer sector dimensions $d_1, \dots, d_n$, and let p_i be a probability distribution over the sectors. Set

$$M = \sum_{i=1}^{n} d_i, \qquad S_{\mathrm{gen}}(p, d) = -\sum_{i=1}^{n} p_i \log p_i + \sum_{i=1}^{n} p_i \log d_i.$$

This is the counting entropy of sectors with one-dimensional bulk blocks, or with d_i counting the whole sector. For blocks $\rho = \bigoplus_i p_i(\rho_i \otimes I_{d_i}/d_i)$ with bulk dimensions b_i, the full entropy is $\log M - D_{\mathrm{KL}}(p\|w/M) - \sum_i p_i D_{\mathrm{KL}}(\rho_i\|I_{b_i}/b_i)$ with $w_i = b_i d_i$ and $M = \sum_i w_i$, so saturation also requires maximally mixed bulk blocks [4].

Proposition 3.1 (Finite sector entropy maximum)**.** *For fixed d_i,*

$$p_i^\star = \frac{d_i}{M}, \qquad \max_p S_{\mathrm{gen}}(p, d) = \log M.$$

Proof. With $q_i = d_i/M$,

$$S_{\mathrm{gen}}(p, d) = \log M - \sum_i p_i \log \frac{p_i}{q_i} = \log M - D_{\mathrm{KL}}(p\|q).$$

Nonnegativity of relative entropy proves the result. □

Thus a logarithmic capacity $N = \log M_0$ agrees exactly with the extremal generalized entropy when the closed screen carries total dimension M_0. The same identity compares every admissible integer partition with total count M: $S_{\mathrm{gen}}(p, d) \le \log M$, with equality exactly at $p = d/M$. In the positive-real relaxation, every twice differentiable joint path at fixed M through $p = d/M$, with $u = p'(0)$ and $v = d'(0)$, has $S'_{\mathrm{gen}}(0) = 0$ and $S''_{\mathrm{gen}}(0) = -\sum_i (u_i - v_i/M)^2/p_i \le 0$, so saturation is a nonstrict joint maximum at fixed M [4]. Neither statement constructs an operation that changes integer sector dimensions.

3.2 Exact gradient and Hessian

At the maximizing distribution, the expectation of the logarithmic sector operator is

$$\mathcal{A}(d) := \sum_{i=1}^{n} \frac{d_i}{M} \log d_i. \tag{2}$$

The dimensions are integers in the finite model. Extend them to positive real variables solely for the following differential calculation.

Proposition 3.2 (Analytic curvature of the sector observable)**.** *The exact gradient of Eq.* (2) *is*

$$\frac{\partial \mathcal{A}}{\partial d_i} = \frac{\log d_i + 1 - \mathcal{A}}{M}. \tag{3}$$

Writing $g_i = \partial_i \mathcal{A}$*, its Hessian is*

$$\frac{\partial^2 \mathcal{A}}{\partial d_i \partial d_j} = \frac{\delta_{ij}}{M d_i} - \frac{g_i + g_j}{M}. \tag{4}$$

At the symmetric point $d_i = d$*,*

$$\operatorname{Hess} \mathcal{A} = \frac{1}{nd^2}\left(I - \frac{2}{n}J\right), \tag{5}$$

where $J = \mathbf{1}\mathbf{1}^T$*. Its eigenvalue on* $\mathbf{1}$ *is* $-1/(nd^2)$*, and every vector orthogonal to* $\mathbf{1}$ *has eigenvalue* $+1/(nd^2)$*.*

Proof. Differentiate $\mathcal{A} = M^{-1}\sum_i d_i \log d_i$ to obtain Eq. (3). A second derivative gives Eq. (4). At $d_i = d$, one has $M = nd$, $\mathcal{A} = \log d$, and $g_i = 1/(nd)$, which yields Eq. (5). □

Two consequences prevent a false extremum statement. First, $\nabla \mathcal{A} = \mathbf{1}/(nd)$ at the symmetric point, so the point is not stationary in the full positive orthant. Second, a fixed-horizon-capacity variation obeys $\mathbf{1}^T \delta d = 0$, and hence

$$\delta d^T (\operatorname{Hess} \mathcal{A}) \delta d = \frac{\|\delta d\|^2}{nd^2} > 0$$

for every nonzero tangent variation. The symmetric point is locally minimizing along the fixed-M redistribution directions. The negative eigenvalue belongs to the homogeneous direction, which changes M.

3.3 The one-sided budget maximum

The de Sitter sign concerns transfer between the horizon and the observer, rather than redistribution among horizon sectors. Let the closed total capacity be M_0. Assign $c_{\text{obs}} \geq 0$ units in the finite capacity ledger to the observer, leaving

$$M_h = M_0 - c_{\text{obs}}$$

on the horizon. Suppose the transfer depletes every horizon sector by the same factor $1 - f$, where $f = c_{\text{obs}}/M_0$. For integer dimensions, restrict to values for which the depleted dimensions remain integers. Then

$$S^{\max}_{\text{gen},h}(f) - S^{\max}_{\text{gen},h}(0) = \log(1 - f), \tag{6}$$

and

$$\mathcal{A}((1 - f)d) - \mathcal{A}(d) = \log(1 - f). \tag{7}$$

Both relations are exact.

Theorem 3.3 (Finite transfer law and analytic one-sided maximum)**.** *Every admissible positive integer transfer has a smaller horizon entropy maximum and uniformly depleted sector observable*

$\mathcal{A}$ than the zero-transfer allocation. On the positive-real interpolation $0 \le c_{\mathrm{obs}} < M_0$, the change is strictly decreasing and concave:

$$\frac{d}{dc_{\mathrm{obs}}} \log(M_0 - c_{\mathrm{obs}}) = -\frac{1}{M_0 - c_{\mathrm{obs}}} < 0, \qquad \frac{d^2}{dc_{\mathrm{obs}}^2} \log(M_0 - c_{\mathrm{obs}}) = -\frac{1}{(M_0 - c_{\mathrm{obs}})^2} < 0.$$

The allocation $c_{\mathrm{obs}} = 0$ is a one-sided boundary maximum on both the admissible finite transfers and the analytic transfer path.

For transfers of 2%, 5%, and 10%,

$$\log(0.98) = -0.020203\ldots, \qquad \log(0.95) = -0.051293\ldots, \qquad \log(0.90) = -0.105361\ldots.$$

The exact finite order and its analytic interpolation are the budget statements relevant to the shock sign. Proposition 3.2 supplies compatible negative homogeneous curvature, while also showing why the transfer result cannot be described as an interior maximum of $\mathcal{A}$ at fixed horizon M.

3.4 Conditional gravitational interpretation

In the horizon coordinates used in Ref. [1], the constant term of the shock operator carries the sign of $B'(0)/B(0) = 2r'(0)/r_c$. For a cosmological horizon, the transverse area decreases from the horizon toward the observer, and the shock has the time-advance sign. This radial profile belongs to the gravitational dictionary. The native screen energy excludes a boundedly invertible L^2 energy congruence with the boundary action of a smooth compact three-dimensional Dirichlet bulk, so a bulk reading of the screen needs a radial observation algebra that the native energy does not supply [5]. Under the two physical identifications

$$\text{closed screen capacity} = \text{fixed horizon–observer budget}, \qquad \mathcal{A} = \frac{A_{\mathrm{horizon}}}{4G},$$

Theorem 3.3 supplies the corresponding conditional sign mechanism. Identifying the capacity coordinate c_{obs} with the observer mass used in the gravitational shock calculation is an additional physical dictionary. The finite entropy theorem and transfer law establish none of these identifications. On the executable source, fixed total count and $\log d$ have no established physical volume or area map. Its full return reference has entropy deficit at most $2025/(64V)$ for V ports, and on a refining base-face patch its mutual information per crossing seam tends to zero, so no nonzero universal coefficient multiplies boundary-seam counts of regions on the two-dimensional source [4]. The identification above reads the whole screen count, which those controls leave open.

4 Conditional icosahedral spectrum

Consider the graph Laplacian of the twelve vertices of the icosahedron. Its eigenspaces decompose under the alternating group A_5 on five letters as

$$\mathbb{R}^{12} \simeq \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{5} \oplus \mathbf{3}',$$

with raw Laplacian eigenvalues

$$0, \qquad 5 - \sqrt{5}, \qquad 6, \qquad 5 + \sqrt{5},$$

respectively. This graph is the seam graph of the twelve-port carrier of the executable observer. Under its uniform independent seam schedule both native repairs have quantum spectral gap $(5-\sqrt{5})/60$ per attempt; that gap describes mixing on a fixed carrier, not a shock spectrum [7].

Assumption 4.1 (Discrete gauge transport DS-GAUGE)**.** *The* **3** *generated by the screen isometries consists of exact gauge zero modes of the discrete shock operator, in direct analogy with the smooth* $\ell = 1$ *isometry modes.*

Assumption 4.2 (Nearest-neighbour kinetic identification DS-LAPLACIAN)**.** *The discrete shock kinetic operator is the nearest-neighbour combinatorial Laplacian on the port graph, or its edge-sector line graph, with one positive overall scale.*

These two assumptions are independent. Assumption 4.1 fixes the scale by placing the **3** at the smooth value 2. Assumption 4.2 selects the finite operator whose remaining eigenvalues are being compared.

Proposition 4.3 (Conditional normalized shock spectrum)**.** *Under Assumptions 4.1 and 4.2, the normalized port Laplacian and shock spectra are*

A_5 *block*	*multiplicity*	λ_{norm}	λ_{shock}
1	1	0	-2
3	3	2	0
5	5	$3+\dfrac{3}{\sqrt{5}}$	$1+\dfrac{3}{\sqrt{5}}$
3′	3	$3+\sqrt{5}$	$1+\sqrt{5}$

The invariant mode agrees exactly with the smooth value -2*. The* **5** *shock eigenvalue is* $1+3/\sqrt{5} = 2.341640\ldots$*, and the additional* **3′** *triplet lies at* $1+\sqrt{5} = 2\varphi = 3.236067\ldots$*.*

Proof. Multiply the graph Laplacian by $2/(5-\sqrt{5})$, then subtract $\mu^2 = 2$. Direct simplification gives the displayed values. □

The value -2 carries no enhancement relative to the continuum invariant mode. Any different comparison obtained by leaving the graph scale unfixed is a normalization artifact.

4.1 Ports and edge sectors

The logarithmic sector operator naturally suggests an edge carrier. The low-lying spectrum is unchanged by this choice.

Lemma 4.4 (Regular line-graph spectrum)**.** *Let* G *be a* k*-regular graph with* $k \geq 2$*,* n *vertices, and* m *edges. The line graph* $L(G)$ *has the Laplacian spectrum of* G*, together with the eigenvalue* $2k$ *repeated* $m-n$ *times.*

Proof. Let B be the unoriented vertex-edge incidence matrix. Then

$$BB^T = A_G + kI, \qquad B^TB = A_{L(G)} + 2I.$$

The nonzero spectra of BB^T and B^TB agree, and the spectrum of the $m \times m$ matrix B^TB contains $m-n$ additional zeros relative to the spectrum of the $n \times n$ matrix BB^T, counted with multiplicity.

Since $L(G)$ is $(2k-2)$-regular, converting adjacency eigenvalues to Laplacian eigenvalues gives the claim. □

For the icosahedron, $k=5$, $n=12$, and $m=30$. The edge-sector spectrum is therefore

$$\{0^1, (5-\sqrt{5})^3, 6^5, (5+\sqrt{5})^3, 10^{18}\}.$$

It contains the full port spectrum plus a high block at 10. In particular, the normalization-free ratio between the **5** and rotation-triplet Laplacian eigenvalues is

$$\frac{\lambda_{\mathbf{5}}}{\lambda_{\mathbf{3}}} = \frac{6}{5-\sqrt{5}} = \frac{3(5+\sqrt{5})}{10} = 2.170820\ldots$$

on both carriers. The dodecahedral face graph gives $\varphi^2 = 2.618033\ldots$, while the smooth $\lambda_{\ell=2}/\lambda_{\ell=1}$ ratio is 3.

Lemma 4.4 closes the port-versus-edge ambiguity once the nearest-neighbour Laplacian premise is accepted. It does not establish that premise.

5 The repair generator has a different domain

Let X_r be the configuration space at fixed sector structure, with measure π_r, and let P_v be the single-site heat-bath conditional expectation. The declared heat-bath generator

$$L^{\mathrm{rep}} = \sum_v (I - P_v)$$

acts on

$$K_r = L^2(X_r, \pi_r).$$

Since each P_v is an orthogonal projection,

$$\langle f, L^{\mathrm{rep}} f\rangle = \sum_v \|(I-P_v)f\|^2.$$

Consequently,

$$\ker L^{\mathrm{rep}} = \bigcap_v \mathrm{Ran}\, P_v.$$

Reducing this intersection to the constants requires a separate irreducibility or intersection proof and is unnecessary for the scope statement here.

The executable native action is the seam sum $H_{\mathrm{native}} = \sum_e (I - \Phi_e)$, a positive local Liouville operator built from the record-discarded native repair Φ_e of each seam [6]. It acts on the operator algebra of a fixed carrier and conserves total occupation, whose fixed-number sectors its connected exchanges generate. It therefore changes no sector dimension either, and the domain statement below applies to it.

A horizon shock changes the sector dimensions d_α in

$$L_C = \sum_\alpha (\log d_\alpha) P_\alpha.$$

Such a change deforms the configuration space and lies outside $L^2(X_r, \pi_r)$ at fixed sector structure. Neither generator therefore acts on the sector-dimension shock. Their spectral gaps neither exclude nor supply the time-advance mode. The capacity-transfer law in Theorem 3.3 is a response input on the enlarged sector-dimension space; no dynamical generator on that space is supplied.

A concrete stationary repair control. The balanced conservative twelve-port family illustrates the domain distinction with its actual local repair law. On a connected weighted port graph, let $z = n - (r/p)\mathbf{1}$ and $\kappa = r(p-r)/[p(p-1)]$. Fair endpoint exchange at each Poisson attempt has density drift $Gz = -Lz/2$, uniform stationary occupancy, and $\mathrm{Cov}(z(t), z(0)) = \kappa e^{-|t|L/2}\Pi$. Here t is the Poisson schedule parameter, one expected attempt per seam per unit; it is a model coordinate, not a physical, modular or observer clock time. At fixed graph, its integrated fluctuation record has covariance $4\kappa L^{+}$. This derives a graph Green covariance from local port readback and repair histories; it neither deforms sector dimensions nor identifies the record with a horizon shock.

For a stationary reversible continuous-time chain and centered current-state observable f, spectral resolution gives

$$\langle f, e^{tG} f\rangle_\pi = \sum_j a_j^2 e^{-\gamma_j t}, \qquad \gamma_j \geq 0.$$

It is nonnegative and nonincreasing. A nondecreasing global time change preserves that property. The discrete-time version requires nonnegative eigenvalues, satisfied by a lazy kernel $(I+K)/2$; the nonlazy two-state flip instead gives $(-1)^m$. These are controls for a specified classical autocorrelation, not constraints on an out-of-time- ordered quantum correlator or an enlarged sector-dimension dynamics. Exact finite witnesses and source-pinned operator controls are archived with the repository evidence [10].

The inverse covariance also retains geometric anisotropy: $[L^{-1}, U] = -L^{-1}[L, U]L^{-1}$ on the mean-zero subspace. The native level-four seven-mode band has 14.642% splitting, whereas an independently supplied round-metric control has 0.01561%. Band averages or a $3, 5, 3$ multiplicity pattern therefore do not by themselves supply the rotational covariance assumed by a round horizon attachment. The conditional shock-spectrum identification must keep its metric and representation premises.

6 The fixed-capacity equation of state

The capacity reading of the cosmological-constant term supplies a conditional branch test through the dark-energy equation of state, and its algebraic law is machine-checked. Read the dark-energy density through the capacity relation as $\rho_{\mathrm{DE}}(a) = \kappa/N(a)$, with κ the single positive constant packaging $c^4\, 3\pi/(8\pi G\, l_P^2)$, and define the effective equation of state through the FRW continuity relation, $w(a) = -1 - \frac{1}{3}\,\mathrm{d}\ln\rho_{\mathrm{DE}}/\mathrm{d}\ln a$, which is the effective-fluid quantity projected by a $w_0 w_a$CDM analysis within its chosen cosmological likelihood. Three exact conditional statements follow, each proved in `FixedCapacityWLaw` of the Lean corpus with the capacity value cancelling throughout.

First, on the fixed-capacity branch, constant N, the equation of state is $w(a) = -1$ identically, and the CPL representation is forced to $(w_0, w_a) = (-1, 0)$ exactly, with zero continuous freedom within that branch. This is the immediate consequence of the assumed $\rho_{\mathrm{DE}} = \kappa/N$ identification and the continuity definition of w; it does not derive fixed capacity, the density identification, or a closed dark sector. Second, the exact converse holds on the drift branch: $w(a) = -1 + \frac{1}{3}\,\mathrm{d}\ln N/\mathrm{d}\ln a$, so any measured departure from -1 is a capacity-drift readback only conditional on those same identifications; the power law $N \propto a^{\varepsilon}$ gives the exact line $w = -1 + \varepsilon/3$. Third, if the record capacity is monotone nondecreasing, then $w(a) \geq -1$ at every epoch: this conditional branch forbids a phantom equation of state, and an established $w(a_0) < -1$ forces capacity loss at a_0 through the strict readback.

A prospective test would require the dataset combination, redshift range, statistic, thresholds, fallbacks, and direction-neutral systematics policy to be fixed before exposure. The comparisons below are retrospective diagnostics and carry no prospective score. Exclusion of $(-1, 0)$ in any direction conflicts with the fixed branch; a posterior-mean quadrant is not a valid extra gate. Conversely, compatibility with $(-1, 0)$ does not validate fixed capacity or OPH because that point is also the standard ΛCDM null. No comparison is frozen or armed by this paper. The fixed branch is registered separately in the frozen-prediction ladder [8]. One registration (FZ-01) freezes the point $(w_0, w_a) = (-1, 0)$ for a later DESI DR3, CMB and supernova combination with a thawing-mean condition. A second (FZ-13) registers the conditional fixed-global-capacity branch with direction-neutral empirical credible ellipsoids; it scores only the fixed branch, the monotone branch cannot rescue it, and because the point coincides with ΛCDM its survival earns no distinctive OPH support.

For a seen-data diagnostic, a script checked the official SHA-256 values for four DESI DR2 chains in each of four w_0w_aCDM combinations. It then applied the exact CPL endpoint condition $w_0 \geq -1$ and $w_0 + (2/3)w_a \geq -1$ on $0 \leq z \leq 2$. The posterior-weight fractions in that conditional monotone subset are 0.00705%, 0.1125%, 0.01206%, and 0.02054% for BAO+CMB and its Pantheon+, Union3, and DESY5 combinations, respectively [2]. Two combinations have at least one chain with no raw sample in the rare subset. The respective raw tail counts are 6, 126, 15, and 11. Union3 is represented in all four chains but is sparse. These seen-data fractions depend on the collaboration's model, likelihood, and priors; they are not model evidences, frequentist exclusions, direct capacity measurements, or a prediction score. On the same official chains, a Gaussian moment summary places $(-1, 0)$ at two-sided normal equivalents 2.61, 2.60, 3.38 and 3.93 for BAO+CMB and its Pantheon+, Union3 and DESY5 combinations. These data precede both registrations, so the values are retrospective tension diagnostics for the fixed branch; they are not the empirical-quantile rule registered in FZ-13 and carry no frozen verdict [9].

Four separately authenticated official DESI DR2 BAO+CMB base-ΛCDM chains give, under a direct samplewise nonlinear transform,

$$\Lambda\ell_P^2 = (2.96770 \pm 0.03978) \times 10^{-122},$$

with weighted 2.5–97.5 percent interval $[2.88997, 3.04530] \times 10^{-122}$. This automatically retains the sampled H_0–Ω_Λ covariance. It is a target-informed retrospective ΛCDM posterior display, not an OPH likelihood or theory uncertainty [2, 3]. The conversion between N and $\Lambda\ell_P^2$ is the de Sitter horizon entropy $N = 3\pi/(\Lambda\ell_P^2)$ in nats, so it holds by definition and fixes no value of N; no independent determination of N exists, and the comparisons below test candidate formulas for N. The FZ-01 capacity band $N \in [3.30836, 3.31137] \times 10^{122}$ corresponds to $\Lambda\ell_P^2 \in [2.84618, 2.84878] \times 10^{-122}$, below the 2.5 percent point of this display, and the closure candidates $N = 3.2921 \times 10^{122}$ and 3.3001×10^{122} give values 2.64 and 2.81 posterior standard deviations below its mean. DESI DR2 precedes the FZ-01 freeze, so this is a retrospective tension under baseline ΛCDM; the FZ-01 capacity rule uses its own frozen posterior and precision condition [8, 9].

A second conditional diagnostic composes the capacity law with the local electroweak-closure tangent. It requires three extra physical premises: (B1) the closure capacity is the same physical N; (B2) the finite closure relation is an epochwise physical law; and (B3) physical evolution selects the committed root branch and co-variation convention while the local clock reads the same homogeneous cosmological α with the required time map. The mathematical derivative is interval-certified independently of those physical premises. An outward-rounded implicit-function calculation proves the selected declared root branch continuously differentiable for $|\Delta \ln \alpha| \leq 10^{-5}$, with

$$\frac{\mathrm{d}\ln N}{\mathrm{d}\ln \alpha} \in [-0.214173865, -0.206176031], \qquad \frac{\mathrm{d}\ln \alpha}{\mathrm{d}\ln N} \in [-4.850224325, -4.669103775].$$

The reciprocal is formed only after zero is excluded. Using the conservative slope bound and $H_0 = 67.4\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$, all three physical premises together let the optical-clock drift limit imply $|1 + w_0| \leq 6.94 \times 10^{-10}$ [12]. The static closure solver does not select the physical cosmic branch, and its certified local domain cannot be extrapolated over the order-one changes suggested by some CPL histories. The diagnostic restricts but does not empty the monotone branch.

7 Claim boundary

Statement	Scope
$\mu^2 = d - 2 = \lambda_{\ell=1}$ in pure de Sitter	exact smooth-geometry identity
$S_{\mathrm{gen}}^{\max} = \log M$	exact finite entropy maximum over sector probabilities
Uniform-depletion drop $\log(1 - f)$	exact finite law for admissible integer dimensions
Gradient and Hessian of $\mathcal{A}(d)$	exact analytic-relaxation identity
One-sided horizon-budget maximum	exact order on admissible finite transfers and exact analytic interpolation; gravitational reading conditional
Port/edge low-spectrum equality	exact regular line-graph theorem
Icosahedral shock spectrum	conditional on exact discrete gauge transport and the scaled nearest-neighbour Laplacian identification
Shock coefficient and physical screen attachment	not derived
Static-patch trace representation	excluded from the claim; the cited trace obstruction stands

The coefficient multiplying the shock response requires a generalized-entropy normalization and a physical horizon-area dictionary on one common domain. The finite calculation fixes neither. It also gives no Hilbert-space trace whose correlators satisfy the conditions tested in Ref. [1].

Closed tested classes and surviving alternatives

The external trace obstruction concerns the proposal tested in Ref. [1]: a Euclidean static-patch path integral with an observer worldline interpreted as a positive cyclic Hilbert-space trace for the stated correlators. Horizon-screen descriptions without that trace interpretation are outside the tested class.

The repair-domain statement concerns the declared fixed-sector L^{rep} on $L^2(X_r, \pi_r)$. A separately constructed dimension-changing generator is not excluded. The graph-spectrum statement concerns the scaled nearest-neighbour icosahedral port Laplacian and its line graph under exact gauge transport. Weighted, longer-range, refinement-limit, and other source-derived kinetic operators are possible and require their own spectra. One refinement-limit operator is constructed: along a common subsequence of refining source screens the native cell Green operators converge in operator norm to a compact injective G, and $A = G^{-1}$ defines a regular, strongly local Dirichlet form with

$n/49920 \le \lambda_n(A) \le 3n/2$ and heat-trace spectral dimension two [5]. That limit selects no unique whole-sequence limit and no physical metric, so a shock reading of A needs its own gauge-transport and normalization premises. On the executed federated screens of 240, 960 and 3,840 ports, the numerically computed ratio of the second to the first low band of the native port Laplacian is 2.68, 2.92 and 2.97, against $6/(5-\sqrt{5}) \approx 2.17$ on one carrier and 3 on the round sphere [11]. The lowest band is a near triplet whose relative spread grows from 0.92 to 2.31 percent over these levels, so the declared cross-carrier routing carries no carrier-wide A_5 covariance, and rotational covariance of a refinement limit is a premise.

These boundaries delimit one trace interpretation, one fixed-sector generator, and one conditional finite kinetic operator. The exact finite entropy, transfer, and line-graph identities hold independently of those physical attachments.

8 Conclusion

A closed capacity budget has a distinguished transfer direction. Moving capacity from a horizon screen to its observer changes the extremal entropy and the logarithmic sector observable $\mathcal{A}$ by $\log(1-f)$. The sign is negative for every allowed positive finite transfer and throughout the positive-real interpolation, so the zero-observer allocation is a one-sided boundary maximum. The analytic Hessian of $\mathcal{A}$ has negative homogeneous curvature and positive fixed-horizon-capacity tangent curvature, while the joint counting entropy is a nonstrict maximum along fixed-capacity joint variations. The sign result therefore rests on the capacity transfer, not on an interior maximum of $\mathcal{A}$.

The pure de Sitter normalization $\mu^2 = \lambda_{\ell=1}$ is exact. An icosahedral spectrum follows only after exact discrete gauge transport and a nearest-neighbour kinetic operator are assumed. The regular line-graph identity then makes its low spectrum independent of choosing ports or edge sectors. These results supply a conditional finite-screen mechanism for the time-advance sign, with the ledger-to-mass map, coefficient, gravitational attachment, and sector-dimension dynamics as separate premises. The exact identities hold if those attachments fail; such a failure would reject the corresponding physical interpretation inside its stated class.

Credited prior work

The results of this paper build on the following established work. Each entry names an established result as it is used here and cites its sources.

- Constant-Lambda equations of state, CPL fits, optical-clock drift limits, official cosmology chains, and interval implicit-function methods [17, 18, 27, 26, 14, 28].
- Finite Shannon entropy, weighted sector entropy, and logarithmic calculus [31, 19].
- Gibbons-Hawking entropy and gravitational shock-sign analyses [23, 20, 25, 22, 13].
- Pure de Sitter horizon-shock equations and scalar spherical harmonics [25, 30, 13, 15].
- Regular graph line-graph spectra and icosahedral adjacency spectra [16, 24].
- Typed operator domains and fixed-sector Markov or repair generators [29, 21].

References

[1] Y. Chen, D. Stanford, H. Tang, and Z. Yang, *Negative shocks versus static patch holography*, arXiv:2607.14042 [hep-th], 2026. https://arxiv.org/abs/2607.14042

[2] DESI Collaboration, *DESI DR2 Results II: Measurements of Baryon Acoustic Oscillations and Cosmological Constraints*, arXiv:2503.14738, 2025. https://arxiv.org/abs/2503.14738. Official chains and checksum manifest: https://data.desi.lbl.gov/public/papers/y3/bao-cosmo-params/README.html. Manifest authenticated by the replay: https://data.desi.lbl.gov/public/papers/y3/bao-cosmo-params/dr2_vac_dr2_bao-cosmo-params_v1.0.sha256sum

[3] B. Müller et al., *Observer Patch Holography: formal proofs and executable receipts*, 2026. https://github.com/FloatingPragma/observer-patch-holography/tree/r2060/code/cosmology/fixed_capacity_wlaw

[4] FloatingPragma, *Tangent first laws and finite entropy changes*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/code/geometry/ENTROPY_FIRST_LAW_AUDIT.md

[5] FloatingPragma, *A spatial fluctuation field from one native history*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/model/physical_space_time/OCCUPATION_FIELD.md

[6] FloatingPragma, *A local action for native quantum observations*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/model/native_action/README.md

[7] FloatingPragma, *Quantum mixing of the native C/E repairs*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/model/observer/QUANTUM_MIXING.md

[8] FloatingPragma, *The frozen-prediction register*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/claims/frozen_prediction_register.json

[9] FloatingPragma, *Cosmology postdiction ledger*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/code/cosmology/postdiction_ledger/COSMOLOGY_POSTDICTION_LEDGER.md

[10] FloatingPragma, *Observer-dynamics finite witnesses and operator controls*, 2026. https://github.com/FloatingPragma/observer-patch-holography/tree/r2060/evidence/observer_dynamics_20260925/sim-analysis

[11] FloatingPragma, *What propagates on a federation of identical elementary carriers*, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/model/federated_observer/SCALING.md; receipt: https://github.com/FloatingPragma/observer-patch-holography/blob/r2060/model/federated_observer/results.json

[12] M. Filzinger et al., *Improved Limits on the Coupling of Ultralight Bosonic Dark Matter to Photons from Optical Atomic Clock Comparisons*, Phys. Rev. Lett. **130**, 253001 (2023). https://doi.org/10.1103/PhysRevLett.130.253001

[13] Aalsma, L., and Shiu, G. (2020). "Chaos and complementarity in de Sitter space." *Journal of High Energy Physics* **2020**, 152. https://doi.org/10.1007/JHEP05(2020)152

[14] Adame, A. G., and others (2025). "DESI 2024 VI: cosmological constraints from the measurements of baryon acoustic oscillations." *Journal of Cosmology and Astroparticle Physics* **2025**, 021. https://doi.org/10.1088/1475-7516/2025/02/021

[15] Atkinson, K., and Han, W. (2012). *Spherical Harmonics and Approximations on the Unit Sphere: An Introduction*. Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-25983-8

[16] Brouwer, A. E., and Haemers, W. H. (2012). *Spectra of Graphs*. Springer, New York (Universitext). https://doi.org/10.1007/978-1-4614-1939-6

[17] Carroll, S. M. (2001). "The cosmological constant." *Living Reviews in Relativity* **4**, 1. https://doi.org/10.12942/lrr-2001-1

[18] Chevallier, M., and Polarski, D. (2001). "Accelerating universes with scaling dark matter." *International Journal of Modern Physics D* **10**, 213–223. https://doi.org/10.1142/S0218271801000822

[19] Cover, T. M., and Thomas, J. A. (2006). *Elements of Information Theory*. Wiley. https://doi.org/10.1002/047174882X

[20] Dray, T., and 't Hooft, G. (1985). "The gravitational shock wave of a massless particle." *Nuclear Physics B* **253**, 173–188. https://doi.org/10.1016/0550-3213(85)90525-5

[21] Ethier, S. N., and Kurtz, T. G. (1986). *Markov Processes: Characterization and Convergence*. Wiley. https://doi.org/10.1002/9780470316658

[22] Gao, S., and Wald, R. M. (2000). "Theorems on gravitational time delay and related issues." *Classical and Quantum Gravity* **17**, 4999–5008. https://doi.org/10.1088/0264-9381/17/24/305

[23] Gibbons, G. W., and Hawking, S. W. (1977). "Cosmological event horizons, thermodynamics, and particle creation." *Physical Review D* **15**, 2738–2751. https://doi.org/10.1103/PhysRevD.15.2738

[24] Godsil, C., and Royle, G. (2001). *Algebraic Graph Theory*. Springer. https://doi.org/10.1007/978-1-4613-0163-9

[25] Hotta, M., and Tanaka, M. (1993). "Shock-wave geometry with nonvanishing cosmological constant." *Classical and Quantum Gravity* **10**, 307–314. https://doi.org/10.1088/0264-9381/10/2/012

[26] Lange, R., Huntemann, N., Rahm, J. M., Sanner, C., Shao, H., Lipphardt, B., et al. (2021). "Improved limits for violations of local position invariance from atomic clock comparisons." *Physical Review Letters* **126**, 011102. https://doi.org/10.1103/PhysRevLett.126.011102

[27] Linder, E. V. (2003). "Exploring the expansion history of the universe." *Physical Review Letters* **90**, 091301. https://doi.org/10.1103/PhysRevLett.90.091301

[28] Neumaier, A. (1990). *Interval Methods for Systems of Equations*. Cambridge University Press. https://doi.org/10.1017/CBO9780511526473

[29] Reed, M., and Simon, B. (1980). *Methods of Modern Mathematical Physics I: Functional Analysis*. Academic Press. ISBN 0-12-585050-6.

[30] Sfetsos, K. (1995). "On gravitational shock waves in curved spacetimes." *Nuclear Physics B* **436**, 721–745. https://doi.org/10.1016/0550-3213(94)00573-w

[31] Shannon, C. E. (1948). "A Mathematical Theory of Communication." *Bell System Technical Journal* **27**, 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x

AI Assistance Disclosure

This research project used research-grade commercial models, including Anthropic's Fable and OpenAI's GPT-6 Astra, for research support, software development, editing, and synthesis. The authors are responsible for the paper's claims, methods, and final text.