

Observer-Patch Holography and the Dark Matter Phenomenon

Bernhard Mueller

May 3, 2026

Paper release: r1496 **Released:** June 20, 2026

Abstract

In OPH, gravity is the stress bookkeeping of stable agreement between finite observer patches. Imperfect agreement leaves a record-repair remainder on overlap collars. That remainder is electromagnetically dark and gravitates as an information-defect stress correction.

The continuation gives a galaxy-limit equation, cluster and cosmology contracts, and a coefficient target. In old, settled, low-acceleration galaxies the correction can approximate the MOND acceleration law. In clusters and cosmology it must be treated as transported stress with its own equations. A scalar source table is not enough for CMB or growth predictions; the full likelihood evaluation requires a finite covariant packet parent, source provenance, global-pooling checks, and frozen source/solver/likelihood hashes.

What This Paper Contributes

The dark matter evidence is not one observation. Galaxy rotation curves, the baryonic Tully-Fisher relation, the radial acceleration relation, cluster lensing, CMB peaks, BAO, growth, and S_8 all ask for a gravitating sector that ordinary luminous matter does not supply. This paper treats the MOND-like galaxy law as a clue about equilibrium response and the Λ CDM evidence as a clue about transported stress.

The OPH move is to give that stress an observer-patch origin. Imperfect record repair leaves a collar remainder. It has no luminous charge, so electromagnetic probes read no matter signal. It gravitates because OPH puts stable collar remainders into the effective stress bookkeeping. In settled low-acceleration galaxies the same mechanism yields the RAR-shaped response. In clusters and cosmology it becomes a transported component with its own relaxation and likelihood contracts. The paper is a continuation surface, so the joint fit remains the decisive gate.

Contents

1	Plain Picture	3
2	Three Claim Layers	3
3	The Dark Matter Challenge	3
4	Standard Routes	4
4.1	Cold Dark Matter	4
4.2	MOND	4

5	The OPH Route	5
6	Static Galaxy Derivation	6
6.1	Source From The Carried Collar Remainder	6
6.2	The Acceleration Variable	7
6.3	Poisson Activation	7
6.4	Conservative Potential Equation	8
6.5	Newtonian And Deep-IR Limits	8
6.6	Effective Dark Density	9
7	The Finite-Collar Coefficient	9
7.1	Unit Branch	9
7.2	$\mathbb{Z}_6$ Reserve Branch	9
7.3	Finite Thickness	10
8	Lensing And Relativistic Stress	11
9	Clusters And Transport	12
10	CMB, BAO, Growth, And S_8	13
11	Abundance And Kernel Targets	15
11.1	Static FLRW Boundary	15
11.2	Flat Capacity-Saturated State Selection	16
11.3	Environmental Kernel	17
11.4	Compressed CMB, BAO, Growth, And S_8 Diagnostic	18
12	Likelihood Contracts	18
12.1	SPARC Disk-Potential Contract	18
12.2	Cluster Map Contract	18
12.3	Boltzmann Contract	18
13	Numerical Galaxy Comparison	19
13.1	SPARC RAR	19
13.2	Rotation Diagnostics	19
13.3	Systematic Likelihood Scaffold	20
13.4	Empirical Implementation Scorecard	20
13.5	Implementation Work Packages	21
13.6	Correction Audit	21
14	Claim Boundaries	22
15	Proof Ledger	23
15.1	Positive Galaxy Source	23
15.2	Quotient-Edge Scalar Opportunity	24
15.3	Uniform Finite-Thickness Reserve	24
15.4	No-Slip Settled Stress	25
15.5	Repair Relaxation Scale	26
15.6	Environmental Kernel Around A Nonzero Field	26

15.7	Background Abundance Green Function	27
15.8	Linear Relaxation Transfer	27
15.9	Static-To-Linear Separation	28
15.10	Parent Collar Functional	28
15.11	Finite Repair Matrix	29
15.12	Finite Repair-Graph Gap	30
15.13	Jacobi Repair Clock	30
15.14	Homogeneous Anomaly State Selection	31
16	Falsifiers	32
17	Reproducibility	32

1 Plain Picture

A galaxy does two jobs in this model. Its baryons source the ordinary weak field, and its finite observer-overlap collars carry a record-repair load. The load is dark to electromagnetic probes because ordinary luminous charges are absent. It gravitates because OPH puts carried collar remainders into the effective stress bookkeeping.

The RAR shape has a simple origin. The relevant repair opportunities live on cuts, so their broad count scales like $r_M/r = \sqrt{g_b/a_{0,\text{OPH}}}$. Independent local opportunity counts give a Poisson zero-count law. The active fraction is $1 - \exp[-\lambda_{\text{collar}} \sqrt{g_b/a_{0,\text{OPH}}}]$. The conservative field equation promotes that scalar response into a potential equation. The simple algebraic acceleration law is exact only when symmetry removes the curl term.

2 Three Claim Layers

The first layer is the static galaxy law. It contains the RAR shape, the BTFR limit, the effective dark density, the no-slip lensing rule, and the $\mathbb{Z}_6$ /Poisson coefficient.

The second layer is the dynamic dark component. It treats the anomaly as a transported stress that relaxes toward the static equilibrium over τ_{rec} . Cluster offsets use this layer.

The third layer is cosmological dark matter. It uses the same stress in linear perturbation theory, with a background abundance, a relaxation rate, and an equilibrium kernel. CMB, BAO, lensing, growth, and S_8 test this layer.

3 The Dark Matter Challenge

In Newtonian gravity and in the weak-field limit of general relativity, a test particle in a circular orbit satisfies

$$\frac{v^2(r)}{r} = g(r) = \frac{GM(< r)}{r^2}. \quad (1)$$

If almost all gravitating matter is visible baryonic matter, then outside the bright disk the enclosed mass should stop growing quickly and the rotation speed should fall as $v(r) \propto r^{-1/2}$. Disk galaxies do not behave that way. Their outer rotation curves often remain flat, so the inferred dynamical mass grows approximately linearly with radius:

$$M_{\text{dyn}}(< r) = \frac{v^2 r}{G}. \quad (2)$$

The galaxy problem is only one part of the evidence. Clusters need more gravity than the observed gas and galaxies provide. Weak lensing maps in merging systems such as the Bullet Cluster show gravitational potential peaks separated from the X-ray gas [12]. The CMB and large-scale structure prefer a gravitating component that is mostly non-baryonic, close to pressureless, and present before galaxies settle [5]. Planck’s base Λ CDM parameter set gives $\Omega_c h^2 = 0.120 \pm 0.001$ and $\Omega_b h^2 = 0.0224 \pm 0.0001$, so most matter is not baryonic in that model [5].

The challenge includes more than “flat rotation curves”. A good theory has to explain:

- galaxy rotation curves and the baryonic Tully-Fisher relation;
- the tight radial acceleration relation, abbreviated RAR;
- galaxy and cluster lensing;
- cluster offsets in mergers;
- CMB acoustic peaks, BAO, matter power, weak lensing, and S_8 .

4 Standard Routes

4.1 Cold Dark Matter

The standard route adds a new non-baryonic matter component. In its simplest form it is cold, effectively collisionless, and pressureless on cosmological scales. Λ CDM uses the same component for CMB peaks, large-scale structure, clusters, lensing, and galaxy halos.

The particle identity is absent. Direct detection has produced no confirmed dark matter particle. Galaxy data show a tight connection between baryonic acceleration and observed acceleration, much tighter than a naive halo-plus-disk picture suggests. Halo fitting also introduces galaxy-level nuisance freedom, while the RAR looks like a law.

4.2 MOND

MOND should be treated as a clue, not as a failure. Its basic phenomenology starts from a critical acceleration a_0 , and in the deep low-acceleration regime it gives

$$g \simeq \sqrt{a_0 g_b}. \quad (3)$$

For a point baryonic mass this implies

$$v^4 = GM_b a_0, \quad (4)$$

which is exactly the baryonic Tully-Fisher scaling [10]. This is MOND’s great success: it goes straight at the galaxy regularity.

The pressure points start when the same static acceleration rule is asked to do more jobs than a galaxy equilibrium law can naturally do:

- *Relativistic completion and lensing.* A nonrelativistic force law does not by itself say which metric photons follow, or how the modified field contributes to the stress side of Einstein’s equations. Relativistic completions such as TeVeS exist [11], but the lensing sector then depends on extra fields and closure assumptions.

- *Clusters.* MOND reduces the cluster mass discrepancy but does not remove the need for additional gravitating stress in clusters. Merging systems are especially sharp: if the anomaly is an instantaneous function of the local baryonic acceleration, the lensing peak wants to remain too tightly tied to the visible baryonic source.
- *The early universe.* The CMB acoustic peaks, BAO, matter growth, CMB lensing, and weak-lensing amplitudes need a homogeneous and perturbative gravitating component before settled galaxies exist. A static RAR law does not by itself define that background abundance or its linear response.
- *Non-spherical and time-dependent systems.* The algebraic relation $g \simeq \sqrt{a_0 g_b}$ is clean in high-symmetry systems. In real three-dimensional fields, curl terms, environmental dependence, transport, and relaxation can matter.

OPH does not treat the static RAR formula as the master law. It approximates MOND in the regime where the OPH stress sector has relaxed onto the old, settled, low-acceleration galaxy branch. It need not approximate MOND in mergers, clusters, or the early universe, because those systems are not static one-dimensional galaxy equilibria. The underlying object is an effective stress sector carried by finite overlap repair. In old relaxed disks it projects to a baryon-linked acceleration law. In lensing it enters the metric-potential bookkeeping. In cluster mergers it is transported and relaxes over a finite repair time. In FLRW and perturbation theory it has to be specified as a homogeneous charge, load current, and response kernel.

This is the sense in which the MOND pressure points dissolve in OPH. OPH treats those failures as wrong-regime extrapolations, without making the same algebraic law more complicated. The static galaxy formula is allowed to be a very good equilibrium approximation, while lensing, clusters, and cosmology are handled by the corresponding OPH stress, transport, and perturbation equations. Those equations must pass their own falsifiers. The claim is structural separation, with no immunity from data.

5 The OPH Route

Observer-Patch Holography starts from finite observer patches and the demand that neighboring patches agree on overlap data. The relevant OPH inputs for the dark branch are:

- finite overlap collars and fixed-cutoff collar algebras;
- edge and cut registers on those collars;
- read, compare, repair, commit, and refresh operations;
- a finite carried modular remainder on the collar;
- the recovered Einstein branch with a screen-capacity de Sitter scale.

The dark-sector proposal is:

The dark source is a transported modular/collar information-defect remainder. It gravitates because it contributes to the effective stress bookkeeping. It is electromagnetically dark because it is a record-repair remainder, not an ordinary luminous matter species.

No new particle species is postulated. The effective dark stress is the gravitating part of finite-observer information repair.

$$G_{ab} + \Lambda g_{ab} = 8\pi G \left(T_{ab}^{\text{SM}} + T_{ab}^{\mathcal{I}} + T_{ab}^{\text{repair}} \right). \quad (5)$$

The symbol A in formulas below denotes this anomaly or information-defect component. It is an effective stress sector, with no literal particle species claim.

This differs from particle dark matter and MOND in a useful way. It is stress-first like dark matter, so lensing and cosmology have a natural slot. It is baryon-linked like MOND only on the settled galaxy branch, because that equilibrium response is activated by collar/cut repair opportunities sourced by baryonic acceleration. The important contrast with MOND is not the square-root galaxy law; OPH intentionally reproduces that approximation where the equilibrium assumptions apply. The contrast is where the law lives. In MOND, the galaxy formula is often treated as the modification itself. In OPH it is the equilibrium projection of a transported stress component. That is why lensing, clusters, and cosmology can have their own equations without giving up the galaxy regularity.

6 Static Galaxy Derivation

6.1 Source From The Carried Collar Remainder

On the support-visible BW branch the primary theorem is the cap automorphism identity. In the special type-I or effective local representation where the geometric generator may be written inside the cap algebra, the OPH modular decomposition has the local form

$$K_C = 2\pi B_C + K_C^{(\text{anom})} + \text{const.} \quad (6)$$

At finite regulator stage, the nonadditive modular part is carried on the overlap collar. The dark branch packages the galaxy-scale positive remainder as

$$R_C := I(A : D|B) \geq 0, \quad (7)$$

where $I(A : D|B)$ is conditional mutual information. This is a phenomenological carried-defect variable on the continuation branch: small CMI supplies a recovered comparison state and a fixed-collar replacement modulus, not a one-shot exact Markov normal form or an exact finite-cutoff Einstein source. The corresponding effective rest-energy density is written

$$\rho_A c^2 = \frac{15}{8\pi^2 \ell^4} R_C. \quad (8)$$

The weak-field Poisson equation becomes

$$\nabla^2 \Phi = 4\pi G(\rho_b + \rho_A). \quad (9)$$

Conditional theorem 1, attractive sign. If the carried galaxy remainder is $R_C = I(A : D|B)$ and the coarse-grained domain measure is nonnegative, then the anomaly is attractive in the Newtonian weak-field equation.

Proof. Conditional mutual information is nonnegative, so $R_C \geq 0$. Equation (8) gives $\rho_A \geq 0$. Equation (9) then shows that the anomaly enters with the same attractive sign as baryonic mass. $\square$

The source identification is a theorem target.

6.2 The Acceleration Variable

Define the baryonic acceleration ratio

$$x := \frac{g_b}{a_{0,\text{OPH}}}. \quad (10)$$

For a point baryonic source,

$$g_b(r) = \frac{GM_b}{r^2}, \quad r_M := \sqrt{\frac{GM_b}{a_{0,\text{OPH}}}}, \quad (11)$$

so

$$x = \left(\frac{r_M}{r}\right)^2, \quad \sqrt{x} = \frac{r_M}{r}. \quad (12)$$

The OPH support assumption is that settled-galaxy repair opportunities live on codimension-one collar/cut support. A screen-area count would scale as x . A cut-count scales as $\sqrt{x}$, as in (12).

6.3 Poisson Activation

Let the mean number of independent local repair opportunities be

$$\mu(x) = \lambda_{\text{collar}}\sqrt{x}. \quad (13)$$

The Poisson law does not follow from fixing this mean alone. Fixed mean alone would select a geometric distribution on nonnegative integers. The OPH claim needs the stronger collar-count premise: independent increments on disjoint collar intervals, rare local repair events, and refinement stability. Partition an interval I with mean $\mu(I)$ into m equal subintervals. If each subinterval has one activation opportunity with probability $\mu(I)/m + O(m^{-2})$, and the probability of two or more events in one subinterval is $O(m^{-2})$, then for every fixed n ,

$$\Pr[N(I) = n] = \lim_{m \rightarrow \infty} \binom{m}{n} \left(\frac{\mu(I)}{m}\right)^n \left(1 - \frac{\mu(I)}{m}\right)^{m-n} = e^{-\mu(I)} \frac{\mu(I)^n}{n!}. \quad (14)$$

In particular, the zero-count probability is

$$\Pr[N(I) = 0] = \lim_{m \rightarrow \infty} \left(1 - \frac{\mu(I)}{m}\right)^m = e^{-\mu(I)}. \quad (15)$$

The probability that at least one repair opportunity is active is

$$p(x) = 1 - \Pr[N = 0] = 1 - \exp[-\lambda_{\text{collar}}\sqrt{x}]. \quad (16)$$

The activation count is only the counting part of the static law. The source closure says that active scalar repair opportunities recover baryonic flux in the settled one-dimensional branch:

$$g_b = p(x)g_{\text{obs}}. \quad (17)$$

Thus

$$g_{\text{obs}} = \nu_{\text{OPH}}(x)g_b, \quad \nu_{\text{OPH}}(x) = \frac{1}{1 - \exp[-\lambda_{\text{collar}}\sqrt{x}]}. \quad (18)$$

Conditional theorem 2, activation law. Assume codimension-one collar/cut support, independent increments on disjoint collar intervals, rare local repair events, refinement stability, and the settled scalar flux-recovery closure (17). Then the settled one-dimensional galaxy response is (18).

Proof. Codimension-one support gives a mean count proportional to $\sqrt{x}$. The scalar coefficient is λ_{collar} . Independent increments and rare-event refinement give the Poisson law (14). The active probability is the complement of the zero-count event, which is (16). Solving $g_b = p(x)g_{\text{obs}}$ in one-dimensional symmetry gives (18). $\square$

6.4 Conservative Potential Equation

For general disks and three-dimensional baryonic profiles, the algebraic vector rule

$$\mathbf{g}_{\text{alg}} = \nu_{\text{OPH}}(|\mathbf{g}_b|/a_{0,\text{OPH}})\mathbf{g}_b \quad (19)$$

need not be curl-free because

$$\nabla \times \mathbf{g}_{\text{alg}} = \nabla \nu_{\text{OPH}} \times \mathbf{g}_b. \quad (20)$$

A gravitational acceleration must be derived from a potential. The primary static OPH equation is therefore

$$\nabla^2 \Phi = \nabla \cdot \left[\nu_{\text{OPH}} \left(\frac{|\nabla \Phi_b|}{a_{0,\text{OPH}}} \right) \nabla \Phi_b \right], \quad \mathbf{g}_{\text{obs}} = -\nabla \Phi, \quad (21)$$

with

$$\nabla^2 \Phi_b = 4\pi G \rho_b. \quad (22)$$

In one-dimensional symmetry, (21) integrates to

$$g_{\text{obs}} = \nu_{\text{OPH}}(g_b/a_{0,\text{OPH}})g_b + C. \quad (23)$$

The boundary condition $g_{\text{obs}} \rightarrow 0$ when the enclosed baryonic flux vanishes sets $C = 0$, so (18) is exact for spherical, planar, or one-dimensional symmetry. The SPARC tables in this note use the algebraic diagnostic; a full disk-potential solver is required for publication-grade rotation and lensing claims.

6.5 Newtonian And Deep-IR Limits

For $x \gg 1$, $p(x) \rightarrow 1$, hence

$$g_{\text{obs}} \rightarrow g_b. \quad (24)$$

For $x \ll 1$,

$$p(x) = \lambda_{\text{collar}}\sqrt{x} + O(x), \quad (25)$$

so

$$g_{\text{obs}} \simeq \frac{g_b}{\lambda_{\text{collar}}\sqrt{g_b/a_{0,\text{OPH}}}} = \sqrt{\frac{a_{0,\text{OPH}}}{\lambda_{\text{collar}}^2}}g_b = \sqrt{a_{0,\text{eff}}g_b}, \quad (26)$$

where

$$a_{0,\text{eff}} = \frac{a_{0,\text{OPH}}}{\lambda_{\text{collar}}^2}. \quad (27)$$

For a point source,

$$v^4 = GM_b a_{0,\text{eff}}. \quad (28)$$

Theorem 3, BTFR. The OPH activation law implies the baryonic Tully-Fisher scaling in the deep-IR limit.

Proof. For a circular orbit, $v^2/r = g_{\text{obs}}$. In the deep-IR limit, $g_{\text{obs}} = \sqrt{a_{0,\text{eff}} GM_b/r^2}$. Multiplying by r gives $v^2 = \sqrt{GM_b a_{0,\text{eff}}}$, hence (28). $\square$

6.6 Effective Dark Density

The anomaly can be represented as an equilibrium effective density:

$$\mathbf{g}_A = (\nu_{\text{OPH}} - 1)\mathbf{g}_b, \quad \rho_{A,\text{eq}} = -\frac{1}{4\pi G} \nabla \cdot [(\nu_{\text{OPH}} - 1)\mathbf{g}_b]. \quad (29)$$

For a spherical system,

$$M_{\text{dyn}}(r) = \nu_{\text{OPH}}(x) M_b(r), \quad M_A(r) = [\nu_{\text{OPH}}(x) - 1] M_b(r). \quad (30)$$

In the deep-IR point-source limit, $M_{\text{dyn}}(r) \propto r$, which is the effective mass profile behind flat rotation curves.

For a point baryonic source, the exact equilibrium expression is

$$M_A(r) = \frac{M_b}{\exp(\lambda_{\text{collar}} r_M/r) - 1}, \quad (31)$$

and

$$\rho_A(r) = \frac{M_b \lambda_{\text{collar}} r_M \exp(\lambda_{\text{collar}} r_M/r)}{4\pi r^4 [\exp(\lambda_{\text{collar}} r_M/r) - 1]^2}. \quad (32)$$

For an extended spherical baryonic profile, positivity of the settled effective density is the monotonicity condition

$$\frac{d}{dr} ([\nu_{\text{OPH}}(x(r)) - 1] M_b(r)) \geq 0. \quad (33)$$

Profiles that violate this condition need finite pressure, shear, or transport terms instead of the settled dust equilibrium alone.

7 The Finite-Collar Coefficient

7.1 Unit Branch

The default branch is

$$\lambda_{\text{collar}} = 1, \quad a_{0,\text{eff}} = a_{0,\text{OPH}}. \quad (34)$$

This branch sets the finite-collar opportunity density to one.

7.2 $\mathbb{Z}_6$ Reserve Branch

The OPH gauge structure contains the realized quotient

$$G_{\text{phys}} = \frac{SU(3) \times SU(2) \times U(1)}{\mathbb{Z}_6}. \quad (35)$$

The protected shared-edge reserve has mean

$$\epsilon_{\mathbb{Z}_6} = \frac{\bar{\ell}_{\text{shared}}}{|\mathbb{Z}_6|} = \frac{P/4}{6} = \frac{P}{24}. \quad (36)$$

Using

$$P = 1.630968209403959, \quad (37)$$

gives

$$\epsilon_{\mathbb{Z}_6} = 0.06795700872516496. \quad (38)$$

Conditional bridge 4, quotient-edge co-registration. Galaxy scalar repair opportunities are quotient-edge cut-register events on the same edge-center resolution that carries the protected $\mathbb{Z}_6$ reserve. The $\mathbb{Z}_6$ reserve therefore co-registers with the scalar activation count.

Conditional argument. The scalar source is a quotient-local recoverability defect, so pure $\mathbb{Z}_6$ center motion changes no realized matter or Higgs state and is inactive for scalar sourcing. If the scalar count and the protected reserve are forced onto the same center-projector resolution, the reserve removes active scalar opportunities from that same layer. $\square$

Proof obligation. The fixed-cutoff screen-net construction must prove that galaxy scalar activation cannot live in a separate bulk, screen-area, or noncentral collar channel.

Conditional bridge 5, local Poisson reserve survival. If the reserve occupancy in a co-registered local scalar slot at transverse collar coordinate y is Poisson with mean $\epsilon_{\mathbb{Z}_6}(y)$, then the local scalar survival factor is

$$\lambda_{\text{slot}}(y) = \Pr[N_{\mathbb{Z}_6}(y) = 0] = \exp[-\epsilon_{\mathbb{Z}_6}(y)]. \quad (39)$$

Proof. A Poisson variable with mean $\epsilon_{\mathbb{Z}_6}$ has zero-count probability $\exp(-\epsilon_{\mathbb{Z}_6})$. A scalar opportunity survives only when its co-registered reserve count is zero. This proves the local survival factor. $\square$

7.3 Finite Thickness

Let y be the transverse collar coordinate and $w(y)$ the normalized activation weight:

$$\int dy w(y) = 1. \quad (40)$$

The finite-thickness branch gives

$$\lambda_{\text{collar}} = \int dy w(y) \exp[-\epsilon_{\mathbb{Z}_6}(y)]. \quad (41)$$

If

$$\int dy w(y) \epsilon_{\mathbb{Z}_6}(y) = P/24, \quad (42)$$

then Jensen's inequality gives

$$\lambda_{\text{collar}} \geq \exp(-P/24). \quad (43)$$

Equality holds when $\epsilon_{\mathbb{Z}_6}(y)$ is constant on the weighted collar.

Conditional bridge 6, exact finite-thickness coefficient. On the product transverse regulator with one quotient trace, the exact finite-thickness coefficient is $\lambda_{\text{collar}} = \exp(-P/24)$.

Conditional argument. If the finite-thickness reserve density is uniform on the scalar-weighted collar layer, then $\epsilon_{\mathbb{Z}_6}(y) = \epsilon_{\mathbb{Z}_6}$. Equation (41) becomes

$$\lambda_{\text{collar}} = \exp(-\epsilon_{\mathbb{Z}_6}) \int dy w(y) = \exp(-\epsilon_{\mathbb{Z}_6}). \quad (44)$$

Using $\epsilon_{\mathbb{Z}_6} = P/24$ proves the claim. $\square$

On that exact branch,

$$\lambda_{\text{collar}} = \exp(-P/24) = 0.934300639489386 \dots \quad (45)$$

and by (27)

$$a_{0,\text{eff}} = 1.179018696 \times 10^{-10} \text{ m s}^{-2}. \quad (46)$$

Failure consequence. Without finite-thickness uniformity, the coefficient is the integral (41). Exact $\exp(-P/24)$ is not available as a theorem-grade number.

8 Lensing And Relativistic Stress

A density source by itself fixes the Newtonian Poisson equation. Lensing also depends on the spatial stress. The OPH no-slip claim uses the transported record-number stress and its spatial moments.

The conditional OPH stress closure is a kinetic record measure. At each coarse-grained event, the finite collar packet emits a positive measure $d\mathcal{E}_x(v)$ on the future unit timelike shell. Define

$$J_{\mathcal{I}}^a(x) = \int v^a d\mathcal{E}_x(v), \quad T_{\mathcal{I}}^{ab}(x) = \int v^a v^b d\mathcal{E}_x(v). \quad (47)$$

This stress is symmetric and future-positive. Relative to an observer u^a , it decomposes as

$$T_{\mathcal{I}}^{ab} = \rho_{\mathcal{I}} u^a u^b + P_{\mathcal{I}} h^{ab} + q_{\mathcal{I}}^{(a} u^{b)} + \pi_{\mathcal{I}}^{ab}. \quad (48)$$

The settled galaxy branch is the monokinetic rest branch $d\mathcal{E}_x(v) = \rho_{\mathcal{I}}(x) \delta(v - u_{\mathcal{I}}) d\Omega$. Then

$$T_{\mathcal{I}}^{ab} = \rho_{\mathcal{I}} u_{\mathcal{I}}^a u_{\mathcal{I}}^b, \quad P_{\mathcal{I}} = 0, \quad q_{\mathcal{I}}^a = 0, \quad \pi_{\mathcal{I}}^{ab} = 0. \quad (49)$$

In Newtonian gauge,

$$ds^2 = a^2 [-(1 + 2\Psi) d\eta^2 + (1 - 2\Phi) dx^2]. \quad (50)$$

The off-diagonal spatial Einstein equation gives

$$k^2(\Phi - \Psi) = 12\pi G a^2 \sum_i (\rho_i + P_i) \sigma_i. \quad (51)$$

The $\mathcal{I}$ -sector contribution to the right-hand side vanishes on (49). If baryonic, neutrino, radiation, and repair-sector anisotropic stresses are absent or handled separately, then

$$\Phi = \Psi. \quad (52)$$

The lensing potential is therefore the same potential that controls dynamics:

$$\Phi_{\text{lens}} = \frac{\Phi + \Psi}{2} = \Phi. \quad (53)$$

This is where OPH behaves differently from a force-only MOND law. The correction is attached to stress bookkeeping, so the same potential controls slow matter and lensing on settled no-slip galaxy states. Cluster mergers and cosmological perturbations use the full decomposition (48); a velocity-spread branch can carry pressure, heat flux, or anisotropic stress.

9 Clusters And Transport

The static galaxy law is an equilibrium law. Merging clusters need a transported information-defect state. In the cold transported branch, the kinetic stress (47) reduces to a density and velocity field:

$$\partial_t \rho_{\mathcal{I}} + \nabla \cdot (\rho_{\mathcal{I}} \mathbf{v}_{\mathcal{I}}) = -\frac{\rho_{\mathcal{I}} - \rho_{\mathcal{I},\text{eq}}[\rho_b]}{\tau_{\text{rec}}}, \quad (54)$$

$$\partial_t \mathbf{v}_{\mathcal{I}} + (\mathbf{v}_{\mathcal{I}} \cdot \nabla) \mathbf{v}_{\mathcal{I}} = -\nabla \Phi, \quad (55)$$

$$\nabla^2 \Phi = 4\pi G(\rho_b + \rho_{\mathcal{I}}). \quad (56)$$

Velocity-spread branches use the moment equations of (48) instead of (54).

The repair-commit theorem gives the form of the relaxation term. Once the finite collar packet algebra is declared, the fixed-cutoff packet state space, proposal rule, equilibrium weights, reversible transition matrix, and spectral gap are determined by packet microphysics. Let γ_{rec} be that dimensionless gap and t_{commit} the physical commit time. Then

$$\Gamma_{\text{rec}} = \frac{\gamma_{\text{rec}}}{t_{\text{commit}}}, \quad \tau_{\text{rec}} = \frac{t_{\text{commit}}}{\gamma_{\text{rec}}}. \quad (57)$$

Bianchi consistency requires a repair-exchange stress:

$$\nabla_a (T_b^{ab} + T_{\mathcal{I}}^{ab} + T_R^{ab} + \dots) = 0. \quad (58)$$

In quasi-covariant notation,

$$\nabla_a T_{\mathcal{I}}^{ab} = -\Gamma_{\text{rec}}(\rho_{\mathcal{I}} - \rho_{\mathcal{I},\text{eq}})u_{\mathcal{I}}^b, \quad (59)$$

and T_R^{ab} carries the opposite exchange flux.

An OPH-side no-fit clock candidate is the Jacobi mismatch rate. Let $E_{ij} = C_{0i0j}$ be the electric Weyl tidal tensor measured in the local free-fall frame, equivalently the trace-free Newtonian Hessian in the weak field. The homogeneous FLRW Ricci focusing term is subtracted, because it does not create a local overlap-visible cluster offset. The scalar rate

$$\Gamma_J = \left(\frac{E_{ij}E^{ij}}{6} \right)^{1/4}, \quad \tau_J = \Gamma_J^{-1} \quad (60)$$

is invariant under local rotations and has the right units. For a Newtonian point source,

$$E_{ij}E^{ij} = 6 \left(\frac{GM}{r^3} \right)^2, \quad \tau_J = \sqrt{\frac{r^3}{GM}}. \quad (61)$$

The numerical scale is:

Scale	$M/M_{\odot}$	r in kpc	τ_J in Gyr
Inner spiral disk	6.0×10^{10}	10	0.0608673
Outer spiral disk	6.0×10^{10}	30	0.316276
Dwarf scale	1.0×10^9	5	0.166692
Cluster core	1.0×10^{14}	300	0.244986
Massive cluster aperture	1.0×10^{15}	1000	0.471476

For $d_0 = 200$ kpc and $t = 0.2$ Gyr, the massive-cluster aperture row gives $d(t) = 130.859$ kpc.

The conditional physical-time conversion is

$$\Gamma_{\text{rec}}(x, t) = \Gamma_J(x, t), \quad t_{\text{commit}}(x, t) = \frac{\gamma_{\text{rec}}(x, t)}{\Gamma_J(x, t)}. \quad (62)$$

The static RAR law does not determine this clock. Any positive relaxation rate leaves the same equilibrium density. The Jacobi clock requires the premise that repair timing is set by the overlap-visible tidal mismatch rate of transported information-defect records. The theorem-grade packet clock must derive this premise without using cluster offsets as selectors.

For a one-mode merger offset,

$$d(t) = d_0 \exp(-t/\tau_{\text{rec}}), \quad \tau_{\text{rec}} = \frac{t}{\ln(d_0/d(t))}. \quad (63)$$

With $d_0 = 200$ kpc and $t = 0.2$ Gyr:

Retained offset	Required τ_{rec}
50 kpc	0.144 Gyr
100 kpc	0.288 Gyr
134 kpc	0.500 Gyr
150 kpc	0.696 Gyr
164 kpc	1.000 Gyr

Settled galaxies impose

$$\tau_{\text{rec}} < \frac{T_{\text{set}}}{\ln(1/\epsilon)} \quad (64)$$

if the allowed tracking error after settling time T_{set} is ϵ . For $T_{\text{set}} = 5$ Gyr, the bounds are 2.17 Gyr, 1.67 Gyr, and 1.09 Gyr for 10%, 5%, and 1% tracking error.

10 CMB, BAO, Growth, And S_8

The linear cosmology contract has two distinct branches. A conserved homogeneous information-defect branch obeys

$$\rho'_A + 3\mathcal{H}\rho_A = 0, \quad \rho_A(a) = \rho_{A0}a^{-3}, \quad (65)$$

where primes denote conformal time and $\mathcal{H} = a'/a$. This branch behaves like a cold stress component if its kinetic measure is monokinetic and its homogeneous charge $Q_A = a^3\rho_A V_{\text{com}}$ is selected by OPH screen data.

No-go boundary for scalar parents. A scalar parent that emits only $\rho_A(a)$, $\rho_{A,\text{eq}}(a)$, and $B_A(k, a)$ is a source candidate, not a physical Boltzmann closure. It does not determine

$$w_A(a), \quad c_{s,A}^2(k, a), \quad \sigma_A(k, a), \quad Q_A^\mu, \quad \Gamma_{\text{rec}}(k, a),$$

and it does not by itself fix gauge-independent perturbations or the recipient stress needed when repair transfers energy-momentum. In the physical source contract B_A is built from the anomaly-frame baryon density

$$n_b^{(A)} = -u_{A\mu}J_b^\mu,$$

not from a gauge-fixed coordinate density. The physical object is a finite covariant collar-packet parent

$$\mathcal{P}_r[X, g] = (C_r, Z_r, A_r, R_r, G_r, \pi_r[X], L_r[X], Q_r, D_r),$$

where Z_r is a finite packet-state set split into anomaly and recipient channels, π_r is the convex equilibrium packet functional, L_r is the repair generator, Q_r records reaction channels, and D_r records any causal auxiliary response. The parent must emit the stress moments and source projections consumed by the Boltzmann solver; otherwise CMB/BAO/growth rows remain diagnostics. Promoting $\rho_A(a)$, $B_A(k, a)$, $\Gamma_{\text{rec}}(k, a)$, or any associated primordial scalar input also requires a source-provenance DAG and a pooled-reducer receipt. Contradictory no-data-use metadata fails closed, and nonlinear quantities such as amplitudes, ranks, condition numbers, and isocurvature leakage are computed only after additive sufficient statistics are pooled globally.

For the source-only primordial bridge, this closed branch has the stricter contract

$$\nabla_a J_A^a = 0, \quad T_A^{ab} = \rho_A u^a u^b, \quad Q_A^a = 0. \quad (66)$$

It may enter the rank-one primordial theorem only when ρ_A is a function of the same source clock as the rest of the total stress. Transport of a supplied Q_A is not an OPH derivation of the homogeneous abundance.

A repair-exchange branch instead relaxes toward a finite-collar equilibrium source:

$$\rho'_A + 3\mathcal{H}\rho_A = -a\Gamma_{\text{rec}}(\rho_A - \rho_{A,\text{eq}}). \quad (67)$$

The factor $a\Gamma_{\text{rec}}$ is the conformal-time rate associated with the physical repair rate. Define

$$q_A = \frac{\rho_{A,\text{eq}}}{\rho_A}, \quad \delta_{A,\text{eq}}(k, a) = B_A(k, a)\delta_b(k, a). \quad (68)$$

For a pressureless no-slip repair branch with energy exchange parallel to the anomaly four-velocity, the Newtonian-gauge perturbations reduce to

$$\delta'_A = -\theta_A + 3\Phi' - a\Gamma_{\text{rec}}q_A(\delta_A - \delta_{A,\text{eq}}), \quad (69)$$

$$\theta'_A = -\mathcal{H}\theta_A + k^2\Psi. \quad (70)$$

The no-slip statement here concerns the information-defect sector. The total metric slip depends on baryons, radiation, neutrinos, repair stress, and any anisotropic stress in a velocity-spread branch.

For the same primordial bridge, the repair-exchange branch is admissible only with an explicit transfer receipt:

$$\nabla_a T_A^{ab} = Q_A^b, \quad \Xi_a^A = 0, \quad h^a_b Q_A^b = 0 \quad (71)$$

or with interval bounds on the two defects. The dark-sector continuation in this paper does not derive the homogeneous anomaly amplitude Q_A , and the finite-collar kinetic stress construction remains conditional. Therefore the first source-only primordial certification run should use

$$\text{dark_continuation} = \text{OFF}.$$

A supplied dark abundance may be used only as

$$\text{dark_continuation} = \text{CONDITIONAL_SOURCE_STATE},$$

not as a derived OPH primordial prediction.

If $\Gamma_{\text{rec}} > 0$, the recipient sector is not optional. The finite parent must expose a recipient stress T_R^{ab} and reaction channel such that

$$\nabla_a (T_A^{ab} + T_R^{ab} + T_{\text{other}}^{ab}) = 0.$$

Without that explicit stress, a repair sink/source can mimic a scalar damping term but cannot be promoted to an Einstein-source prediction.

The finite scalar-load repair diagnostic gives

$$\gamma_{\text{rec}} = 0.050717471, \quad \tau_{\text{rec}}/t_{\text{commit}} = 19.717071.$$

It maps a cluster-scale relaxation window to the following commit times:

τ_{rec}	t_{commit}
0.3 Gyr	15.2152 Myr
0.5 Gyr	25.3587 Myr
1.0 Gyr	50.7175 Myr

For the same relaxation window, a simple background timing diagnostic gives:

τ_{rec}	$z = 0$	$z = 10$	$z = 1100$
0.3 Gyr	48.4092	2.36213	0.00205644
0.5 Gyr	29.0455	1.41728	0.00123387
1.0 Gyr	14.5228	0.708639	0.000616933

For a hybrid branch, repair relaxation is tiny during acoustic physics and large in low-Hubble settled environments. A cosmology claim requires a finite covariant source artifact, Boltzmann implementation, frozen likelihood hashes, and likelihood tests against CMB TT/TE/EE, CMB lensing, BAO, weak lensing, redshift-space distortions, and S_8 .

11 Abundance And Kernel Targets

The static galaxy law fixes a settled response around resolved baryonic inhomogeneity. A cosmological dark-matter branch needs OPH source outputs:

$$\rho_A(a), \quad B_A(k, a), \quad \Gamma_{\text{rec}}(k, a), \quad w_A, \quad c_{s,A}^2, \quad \sigma_A, \quad Q_A^\mu. \quad (72)$$

The first is the homogeneous anomaly abundance. The second is the linear equilibrium-source kernel in

$$\delta_{A,\text{eq}}(k, a) = B_A(k, a)\delta_b(k, a). \quad (73)$$

The remaining entries close the stress tensor and exchange law.

11.1 Static FLRW Boundary

The nonlinear galaxy law cannot be inserted directly into FLRW perturbation theory. In an exactly homogeneous background there is no preferred baryonic acceleration vector. If one tries to Taylor-expand the static RAR law at $\mathbf{g}_b = 0$, the susceptibility is singular because $\nu_{\text{OPH}}(x) \sim (\lambda_{\text{collar}}\sqrt{x})^{-1}$. The static RAR branch therefore does not select a homogeneous anomaly density or a universal CMB kernel.

This is a useful boundary. It prevents a hidden fit from being smuggled into the galaxy law. The CMB branch has to declare its background anomaly charge and its perturbation kernel from OPH state selection or finite-collar microphysics.

11.2 Flat Capacity-Saturated State Selection

One clean homogeneous branch is flat capacity saturation. The de Sitter screen capacity fixes H_{dS} , and the flat Friedmann constraint at $a = 1$ gives

$$\Omega_{A0} = 1 - \Omega_{\Lambda, \text{OPH}} - \Omega_{b0} - \Omega_{\nu 0} - \Omega_{r0}, \quad \Omega_{\Lambda, \text{OPH}} = \left(\frac{H_{\text{dS}}}{H_0} \right)^2. \quad (74)$$

The transported homogeneous dust branch is

$$\rho_A(a) = \rho_{A0} a^{-3}. \quad (75)$$

Using $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_b h^2 = 0.02237$, the OPH neutrino mass sum, and $\Omega_r = 9.17 \times 10^{-5}$, the residual is:

Quantity	Value
H_{dS}	$55.759940256 \text{ km s}^{-1} \text{ Mpc}^{-1}$
$\Omega_{\Lambda, \text{OPH}}$	0.684423332
Ω_b	0.049243191
Ω_ν	0.002127375
Ω_r	0.000091700
Ω_A	0.264114401
Ω_m	0.315484968
ρ_{A0}	$2.253649933 \times 10^{-27} \text{ kg m}^{-3}$

This matches the Planck-like cold-matter budget at the compressed-parameter level. The number is a state-selection residual. The static galaxy RAR law does not determine it. The OPH theorem that would make it standalone is a finite-screen additive-load selector for the conserved homogeneous anomaly charge.

The closed transported branch does give a theorem-grade current and charge. Let u_A^a be the branch four-velocity and let ρ_A be the coarse-grained information-defect load density. The load current is

$$J_A^a = \rho_A u_A^a. \quad (76)$$

On the closed branch,

$$\nabla_a J_A^a = 0, \quad Q_A[\Sigma] = \int_\Sigma J_A^a n_a d\Sigma \quad (77)$$

is independent of the homogeneous slice. In FLRW this gives

$$Q_A = a^3 \rho_A V_{\text{com}}, \quad \rho_A(a) = \frac{Q_A}{a^3 V_{\text{com}}}. \quad (78)$$

This proves transport of a supplied homogeneous load. It does not fix the amplitude Q_A . The missing OPH object is a nontrivial finite-screen additive load invariant

$$Q_A = F_{\text{OPH}}(N_{\text{scr}}, P, \text{screen normal-form data}). \quad (79)$$

Without that invariant, the abundance is cosmological state data or the flat capacity-saturated residual (74).

11.3 Environmental Kernel

Around a nonzero local background field $\bar{\mathbf{g}}_b$, write

$$x = \frac{|\bar{\mathbf{g}}_b|}{a_{0,\text{OPH}}}, \quad \mathbf{n} = \frac{\bar{\mathbf{g}}_b}{|\bar{\mathbf{g}}_b|}. \quad (80)$$

Linearizing the anomaly field $(\nu_{\text{OPH}} - 1)\mathbf{g}_b$ gives the susceptibility

$$A_{ij} = [\nu_{\text{OPH}}(x) - 1]\delta_{ij} + x\nu'_{\text{OPH}}(x)n_i n_j, \quad (81)$$

where

$$\nu'_{\text{OPH}}(x) = -\frac{\lambda_{\text{collar}} \exp[-\lambda_{\text{collar}}\sqrt{x}]}{2\sqrt{x}[1 - \exp(-\lambda_{\text{collar}}\sqrt{x})]^2}. \quad (82)$$

For a Fourier mode whose direction has cosine μ against $\mathbf{n}$, the density-response multiplier is

$$K_A^{(\rho)}(x, \mu) = \nu_{\text{OPH}}(x) - 1 + x\nu'_{\text{OPH}}(x)\mu^2. \quad (83)$$

An isotropic environmental average gives

$$\langle K_A^{(\rho)} \rangle = \nu_{\text{OPH}}(x) - 1 + \frac{x}{3}\nu'_{\text{OPH}}(x). \quad (84)$$

The contrast kernel used in perturbation equations is normalized by the background density ratio:

$$B_A(k, a) = \frac{\bar{\rho}_b(a)}{\bar{\rho}_A(a)} K_A^{(\rho)}(k, a). \quad (85)$$

In a realistic cosmological calculation, this local expression has to be integrated against the distribution of environmental fields,

$$K_A^{(\rho)}(k, a) = \int dx d\mu \Pi(x, \mu|k, a) [\nu_{\text{OPH}}(x) - 1 + x\nu'_{\text{OPH}}(x)\mu^2]. \quad (86)$$

This average is finite only under an explicit small-field support condition, for example

$$\int_0^\epsilon dx d\mu \Pi(x, \mu|k, a) x^{-1/2} < \infty \quad \text{for some } \epsilon > 0. \quad (87)$$

If Π has an exact atom at $x = 0$, the local nonzero-field kernel is not the correct FLRW response. The distribution Π is an OPH finite-collar output or an explicitly declared environmental closure. Exact FLRW has $x = 0$, so (83) does not provide a regular universal CMB density-response kernel by itself.

The implementation contract uses one parent finite-collar functional:

$$\rho_{A,\text{eq}}[X]c^2 = \frac{15}{8\pi^2\ell(X)^4} \int_{\mathcal{C}_X} d\mu_C I_{\omega_C}(A : D|B). \quad (88)$$

For finite samples s with weights w_s , define

$$R(X) = \frac{\sum_s w_s I_s}{\sum_s w_s}, \quad (89)$$

$$K_A^{(\rho)}(k, a) = \frac{1}{\bar{\rho}_b} \frac{\partial \rho_{A,\text{eq}}}{\partial \delta_b(k, a)}, \quad (90)$$

$$B_A(k, a) = \frac{\bar{\rho}_b(a)}{\bar{\rho}_A(a)} K_A^{(\rho)}(k, a). \quad (91)$$

The same evaluator emits the homogeneous equilibrium density, the linear density-response kernel, the settled galaxy source, and the cluster equilibrium source. A theorem-grade prediction requires the OPH collar measure $d\mu_C$, the finite-screen ensemble, and the small-field support condition. Fitting Π to CMB, weak-lensing, SPARC, or cluster data is excluded.

11.4 Compressed CMB, BAO, Growth, And S_8 Diagnostic

The local Boltzmann plumbing test passes the anomaly as a cold pressureless component with the flat-residual diagnostic parent-grid ratio $\rho_A/\rho_b = 5.363470441$. It uses CAMB 1.6.6, the OPH neutrino mass sum, and diagonal Gaussian compressed rows:

Quantity	Prediction	Target	Pull
Planck Ω_m	0.315905207	0.315 ± 0.007	+0.129
Planck σ_8	0.807787208	0.811 ± 0.006	−0.535
Planck S_8	0.828924043	0.831027707 ± 0.0111	−0.190
DESI DR1 BAO Ω_m	0.315905207	0.295 ± 0.015	+1.394
DESI DR1 BAO/BBN/ θ_* H_0	67.4	68.52 ± 0.62	−1.806
Weak-lensing S_8	0.828924043	0.790 ± 0.016	+2.433

The diagonal compressed value is $\chi^2 = 11.463$ for six rows. This is a plumbing check. It does not replace Planck, DESI, or weak-lensing likelihoods. A publication-grade calculation needs the OPH $B_A(k, a)$ kernel in a custom Boltzmann module and the full experimental covariances.

12 Likelihood Contracts

12.1 SPARC Disk-Potential Contract

The publication-grade SPARC test must solve the conservative disk equation

$$\nabla^2 \Phi = \nabla \cdot \left[\nu_\lambda \left(\frac{|\nabla \Phi_b|}{a_{0,\text{OPH}}} \right) \nabla \Phi_b \right] \quad (92)$$

on an axisymmetric baryonic disk model. The hierarchy must marginalize or profile distance, inclination, disk and bulge mass-to-light ratios, gas scale, intrinsic scatter, velocity covariance, quality cuts, and grid convergence. The unit branch and the $\mathbb{Z}_6$ /Poisson branch are fixed before this test; SPARC is used as a comparison, not as the selector for λ_{collar} .

12.2 Cluster Map Contract

The cluster forward model uses gas hydrodynamics, stellar and galaxy tracers, transported defect stress, the repair source $\rho_{A,\text{eq}}$, lensing projection, X-ray emissivity, SZ pressure, and covariance-aware map likelihoods. Merging clusters test whether the transported stress lags collisional gas. Relaxed clusters test whether the same stress law returns to the static equilibrium source. The static-only failure mode is part of the likelihood contract.

12.3 Boltzmann Contract

The Boltzmann module exposes

$$\bar{\rho}_A(a), \quad \bar{\rho}_{A,\text{eq}}(a), \quad w_A(a), \quad c_{s,A}^2(k, a), \quad \sigma_A(k, a), \quad Q_A^\mu, \quad B_A(k, a), \quad \Gamma_{\text{rec}}(k, a),$$

with the OPH neutrino mass sum in the same run. Every entry must be derived from a frozen finite covariant parent artifact with no CMB, BAO, lensing, SPARC, or cluster fit in that source step. It must recover the Λ CDM cold-component limit when the exchange and stress corrections are turned off, reproduce the compressed diagnostic rows only as a plumbing check, and then run

the full CMB TT/TE/EE, CMB lensing, BAO, supernova, weak-lensing, RSD, and S_8 likelihoods under the same nuisance model used for Λ CDM and w_0w_a . The physical likelihood run is frozen by immutable hashes for the source artifact, Boltzmann solver, and likelihood configuration before the data comparison. The same run must expose official likelihood and CDM-limit reductions at global scope. Shard-local `any()` rollups do not promote the source artifact.

13 Numerical Galaxy Comparison

The OPH capacity benchmark used here is

$$a_{0,\text{OPH}} = 1.029186271 \times 10^{-10} \text{ m s}^{-2}. \quad (93)$$

The common empirical reference is

$$a_0^{\text{emp}} = 1.2 \times 10^{-10} \text{ m s}^{-2}. \quad (94)$$

The $\mathbb{Z}_6$ /Poisson branch gives

$$\lambda_{\text{collar}} = 0.934300639, \quad a_{0,\text{eff}} = 1.179018696 \times 10^{-10} \text{ m s}^{-2}. \quad (95)$$

Quantity	Value	Comparison
$a_{0,\text{OPH}}$	$1.029186271 \times 10^{-10}$	−8.070% versus best same-function RAR
$a_{0,\text{OPH}}$	$1.029186271 \times 10^{-10}$	−14.234% versus empirical reference
$a_{0,\text{eff}}, \mathbb{Z}_6/\text{Poisson}$	$1.179018696 \times 10^{-10}$	+5.314% versus best same-function RAR
$a_{0,\text{eff}}, \mathbb{Z}_6/\text{Poisson}$	$1.179018696 \times 10^{-10}$	−1.748% versus empirical reference

13.1 SPARC RAR

Using the public SPARC RAR data [7, 8], the local diagnostic gives:

Scenario	a_0 in m s^{-2}	RMS dex	Binned RMS dex
OPH unit	$1.029186271 \times 10^{-10}$	0.134208	0.026169
$\mathbb{Z}_6$ /Poisson	$1.179018696 \times 10^{-10}$	0.132834	0.015760
Empirical reference	$1.200000000 \times 10^{-10}$	0.132913	0.016164
Best same-function	$1.119530397 \times 10^{-10}$	0.132947	0.017428

13.2 Rotation Diagnostics

The fixed stellar mass-to-light diagnostic and the first nuisance-profiled diagnostic give:

Scenario	Fixed-M/L RMS	Profiled-M/L χ^2/pt	Profiled RMS
OPH unit	23.366208	1.341948	12.420181
$\mathbb{Z}_6$ /Poisson	22.868866	1.390941	12.590203
Empirical reference	22.842145	1.400930	12.624079

Units for the RMS columns are km s^{-1} .

13.3 Systematic Likelihood Scaffold

A stronger local scaffold profiles stellar M/L, distance, inclination, and gas scale per galaxy using the SPARC sample table [6, 7]. It uses $Q \leq 2$, inclination at least 30° , and 3168 rotation-curve rows.

Scenario	χ^2/pt	χ^2/dof	RMS	Median disk M/L
OPH unit	1.016589	1.275672	11.249541	0.519102
$\mathbb{Z}_6/\text{Poisson}$	1.021093	1.281324	11.276832	0.497435
Empirical reference	1.023261	1.284045	11.285518	0.493865

The systematic scaffold mildly favors the unit branch. The $\mathbb{Z}_6/\text{Poisson}$ branch remains the better acceleration-scale and fixed-M/L RAR match. The fixed-M/L improvement alone is weak evidence for the $\mathbb{Z}_6$ bridge theorem.

13.4 Empirical Implementation Scorecard

The executable empirical pass combines four diagnostic layers: public SPARC tables, the flat capacity-saturated homogeneous anomaly state, the finite repair matrix with a cluster timing gate, and the compressed CAMB rows. The single-pass output is:

Quantity	OPH output	Measurement comparison
Ω_A	0.264114401	Planck-like matter budget
$\Omega_{\Lambda, \text{OPH}}$	0.684423332	capacity de Sitter branch
ρ_A/ρ_b	5.363470441	flat residual ratio
γ_{rec}	0.050717471	finite repair matrix gap
$a_{0, \text{OPH}}$	$1.029186271 \times 10^{-10}$	8.070% low versus best RAR
$a_{0, \text{eff}}, \mathbb{Z}_6/\text{Poisson}$	$1.179018696 \times 10^{-10}$	1.748% low versus common empirical a_0
SPARC systematic χ^2/pt , unit	1.016589	best scaffold row
SPARC systematic χ^2/pt , $\mathbb{Z}_6$	1.021093	close scaffold row
CAMB Ω_m	0.315905206	+0.129 σ versus Planck
CAMB σ_8	0.807787204	−0.535 σ versus Planck
CAMB S_8	0.828924037	+2.433 σ versus weak lensing

The compressed CMB/BAO/growth score is

$$\chi_{\text{diag}}^2 = 11.463 \quad \text{for six compressed rows.}$$

The pressure point is weak-lensing S_8 . The diagnostic value is 0.828924037. The DES/KiDS compressed row is 0.790 ± 0.016 .

For the cluster timing gate, take a 200 kpc initial anomaly separation, a 0.2 Gyr time after passage, and an example measured offset 150 ± 50 kpc. The repair matrix gives:

τ_{rec} in Gyr	t_{commit} in Myr	Retained fraction	Offset in kpc	χ^2
0.3	15.2152	0.513417	102.683	0.896
0.5	25.3587	0.670320	134.064	0.102
1.0	50.7175	0.818731	163.746	0.076

This row is a timing gate. Map likelihoods need observed lensing maps, gas maps, stellar maps, merger-age priors, and covariance.

13.5 Implementation Work Packages

The code required for paper-grade empirical tests consists of the following modules:

1. An axisymmetric conservative disk-potential solver for

$$\nabla^2 \Phi = \nabla \cdot [\nu(|\nabla \Phi_b|/a_{0,\text{OPH}})\nabla \Phi_b],$$

with grid-convergence tests and lensing-potential output.

2. A hierarchical SPARC likelihood that samples or profiles distance, inclination, disk M/L, bulge M/L, gas scale, intrinsic scatter, velocity covariance, and selection cuts under one nuisance model.
3. A finite-collar parent evaluator for $\rho_A(a)$, $K_A^{(\rho)}(k, a)$, $B_A(k, a)$, and $\rho_{A,\text{eq}}[X]$, using OPH collar samples as input. The scalar-load diagnostic grid is plumbing only.
4. A CAMB or CLASS anomaly module exposing the background density, stress variables, exchange current, relaxation kernel, environmental response kernel, and the OPH neutrino mass sum in one run.
5. A cluster forward model with gas hydrodynamics, galaxy and stellar tracers, transported anomaly stress, lensing projection, X-ray and SZ observables, merger-age priors, and map covariance.
6. A likelihood runner for Planck or ACT spectra, CMB lensing, BAO, supernovae, weak lensing, RSD, SPARC, and cluster maps, with model comparison against Λ CDM, empirical MOND functions, halo models, and $w_0 w_a$ extensions.
7. A reproducibility harness with fixed data hashes, fixed random seeds, unit checks, and independent numerical replication of the scorecard rows.

13.6 Correction Audit

The comparison mostly tests normalization because the OPH static law uses the same acceleration-family shape as the empirical RAR fit. If $a_{0,\text{OPH}}$ is held fixed, different targets imply different λ_{collar} values:

Target	Required λ_{collar}	Reserve $-\ln \lambda_{\text{collar}}$	Ratio to $P/24$
Best same-function RAR	0.958802256	0.042070424	0.619074
Binned RAR	0.936737944	0.065351711	0.961663
Common empirical a_0	0.926096769	0.076776547	1.129781
Fixed-M/L rotation	0.926654674	0.076174302	1.120919
OPH $\mathbb{Z}_6$ /Poisson	0.934300639	0.067957009	1.000000

This rules out several tempting shortcuts. In the normalization-profiled all-point objective, a generalized activation exponent near 0.503420 appears. With fixed $a_{0,\text{OPH}}$, direct RAR objectives give exponents near 0.516 all-point and 0.519 weighted-bin. These shifts are objective-dependent, and flat deep-IR rotation requires 0.5. Moving the exponent would buy little and would damage the BTFR theorem.

Finite-thickness nonuniformity can raise λ_{collar} above $\exp(-P/24)$, which moves the coefficient toward the binned RAR value. It cannot lower λ_{collar} toward the common empirical or fixed-M/L values while the mean reserve remains $P/24$ and Poisson counting is retained. An exact hit to the common empirical acceleration would require about 13% more protected inactive reserve than $P/24$, or a different microscopic selector. That extra reserve needs an OPH theorem before use.

14 Claim Boundaries

Pressure point	OPH route	Status
Unknown particle	Dark source is a carried modular/collar information-defect stress.	Conditional source target.
Galaxy RAR and BTFR	Collar/cut activation gives the square-root transition and Poisson activation law under independent-increment repair counting.	Conditional static theorem.
Acceleration scale	Capacity scale gives $a_{0,\text{OPH}}$; $\mathbb{Z}_6$ /Poisson reserve gives $a_{0,\text{eff}}$ close to the empirical scale.	Conditional coefficient theorem.
Galaxy lensing	No-slip branch gives $\Phi = \Psi$ if the information-defect stress is dust on settled galaxies.	Conditional stress theorem.
Cluster offsets	Transported anomaly can lag collisional gas; the Jacobi clock gives a no-fit scale under the tidal-mismatch premise.	Forward likelihood contract.
CMB and growth	Flat capacity state selection gives a CDM-like homogeneous residual; the closed branch has a conserved load current; environmental kernels describe nonzero-field response.	Amplitude selector and real Boltzmann likelihood work in progress.

The supported claim is:

OPH defines an imperfect-information correction to the recovered Einstein branch. Under explicit bridge premises, the static galaxy equation reproduces the RAR/BTFR functional form from collar/cut repair statistics and predicts an acceleration scale close to SPARC. The unit branch is low in normalization. The $\mathbb{Z}_6$ /Poisson branch lands within 1.748% of the common empirical acceleration reference. Profiled stellar, distance, inclination, and gas nuisances mildly favor the unit branch. The Jacobi clock gives a no-fit cluster-offset scale under its tidal-mismatch premise. Flat capacity state selection gives a Planck-like homogeneous information-defect residual, while closed-branch transport gives a conserved load current. Cluster and CMB/growth likelihoods require measured cluster covariances, a finite-screen amplitude selector, a finite covariant source parent, recipient stress for any nonzero repair exchange, and a frozen Boltzmann/likelihood implementation.

Claim outside this paper:

OPH has fully derived dark matter and solved MOND's cluster and CMB problems.

That stronger statement requires SPARC publication likelihoods, cluster likelihoods, a finite co-variant parent for the cosmological source functions, and frozen Boltzmann likelihoods.

15 Proof Ledger

Target	OPH closure	Numerical state
Galaxy source	Minimal scalar recoverability defect $R_C = I(A : D B)$.	Conditional source target.
Quotient-edge locality	Scalar repair opportunities are edge-center cut-register events on the physical quotient.	Conditional bridge.
Finite thickness	Product transverse regulator with one quotient trace makes the protected center reserve uniform through the weighted collar.	Conditional bridge.
No-slip stress	Positive kinetic collar measure; settled monokinetic branch has dust stress and vanishing information-sector anisotropic stress.	Conditional stress theorem.
Repair relaxation	Fixed repair-commit chain gives $\tau_{\text{rec}} = t_{\text{commit}}/\gamma_{\text{rec}}$.	Conditional packet theorem and clock candidate.
CMB and growth	Nonzero-environment kernel is explicit; exact FLRW static RAR kernel is blocked by a no-go theorem.	Finite-parent Boltzmann contract.
Abundance history	Flat capacity state selection gives a CDM-like residual; the closed transported branch has a conserved load current and charge.	Amplitude selector work in progress.
Linear transfer	The perturbation equation has a relaxation transfer solution between CDM-like and tracking limits.	Requires evaluated $B_A(k, a)$, recipient stress for nonzero exchange, and frozen hashes.
Parent functional	The finite collar expectation of the positive repair defect emits $\rho_{A,\text{eq}}$ and B_A if its FLRW limit is regular.	Scalar source target, not full stress closure.
Repair matrix	A reversible Metropolis kernel on packet states gives a concrete K .	Conditional on declared packet schema.
Jacobi clock	The tidal scalar $(E_{ij}E^{ij}/6)^{1/4}$ gives a no-fit dynamic repair scale under the Jacobi mismatch premise.	Conditional clock.

15.1 Positive Galaxy Source

Theorem target 0A. Let a settled galaxy collar state be judged by a scalar repair defect D_C with these properties: $D_C \geq 0$; $D_C = 0$ exactly on locally recoverable Markov states; D_C is invariant under boundary redundancy; D_C is additive over independent collar components; and D_C is monotone under local recovery maps. Then the minimal scalar branch is

$$D_C = \kappa I(A : D|B), \quad \kappa \geq 0. \quad (96)$$

The OPH dark branch uses $R_C = I(A : D|B)$.

Explicit premise. The galaxy-scale finite-stage modular nonadditivity is represented by this minimal scalar recoverability defect on settled collar states.

Conditional argument. Conditional mutual information has all listed properties on a finite collar. A different scalar with the same zero set and additivity would add an independent scalar datum to the minimal branch. The minimal scalar selector is therefore fixed up to a nonnegative multiplier. The OPH unit convention sets that multiplier to one. Since $I(A : D|B) \geq 0$, the weak-field density defined by the anomaly normalization is nonnegative. $\square$

Proof obligation. Derive the map from the finite-stage modular nonadditivity term to $I(A : D|B)$ on the galaxy branch.

Failure consequence. If the modular remainder is not this positive defect, the attractive source and normalization formula must be replaced by the derived scalar.

15.2 Quotient-Edge Scalar Opportunity

Theorem target 0B. At fixed cutoff, physical collar observables live on the quotient-local fixed-point algebra. A settled galaxy scalar repair opportunity is the local event that the quotient-local recoverability defect on a collar cut is nonzero. Then the scalar opportunity is counted on the edge-center cut register.

Explicit premise. The scalar recoverability event has no independent bulk, screen-area, or non-central collar channel.

Conditional argument. The defect is a physical scalar. Boundary-redundant representatives produce the same defect value because the quotient repair law descends through the projection onto physical collar states. The event that the defect is nonzero is therefore an observable event on the quotient-local collar algebra. For a connected cut, the fixed-cutoff edge-center decomposition supplies the central projectors that resolve physical cut sectors. A scalar event with no vector, tensor, or bulk label selects support through those central projectors. Thus the scalar opportunity count is a quotient-edge cut-register count. The protected $\mathbb{Z}_6$ center reserve is supported on the same register. $\square$

Proof obligation. Prove the fixed-cutoff uniqueness statement: every quotient-local scalar recoverability event is resolved on the edge-center cut register.

Failure consequence. If a separate scalar channel exists, the $\mathbb{Z}_6$ reserve does not fix λ_{collar} without an additional co-registration theorem.

15.3 Uniform Finite-Thickness Reserve

Theorem target 0C. Let the finite collar be a transverse thickening of one quotient-edge cut algebra with normalized transverse weight $w(y)$ and a single quotient trace on the edge algebra.

Let the protected $\mathbb{Z}_6$ reserve be a central edge projector whose mean on that trace is $\epsilon_{\mathbb{Z}_6} = P/24$. Then

$$\lambda_{\text{collar}} = \exp(-P/24). \quad (97)$$

Explicit premise. The scalar-weighted transverse regulator is a product thickening with one quotient trace and uniform protected-center density.

Conditional argument. The protected reserve belongs to the edge-center algebra. The transverse coordinate belongs to the regulator thickening. With one quotient trace on the edge algebra, the trace of the protected central projector is the same on every transverse slice that carries scalar activation weight. Hence $\epsilon_{\mathbb{Z}_6}(y) = \epsilon_{\mathbb{Z}_6}$. Poisson zero-reserve survival gives $\exp(-\epsilon_{\mathbb{Z}_6})$ at each slice. Averaging with $w(y)$ leaves the same value. Using $\epsilon_{\mathbb{Z}_6} = P/24$ gives the stated coefficient. $\square$

Proof obligation. Derive $w(y)$, the transverse regulator, and the quotient trace from base OPH screen/collar data.

Failure consequence. If the reserve density is nonuniform, the coefficient is

$$\int dy w(y) \exp[-\epsilon_{\mathbb{Z}_6}(y)],$$

not necessarily $\exp(-P/24)$.

15.4 No-Slip Settled Stress

Theorem target 0D. Let the information-defect stress be the second moment of a positive finite-collar kinetic measure $d\mathcal{E}_x(v)$ on the future unit timelike shell:

$$T_{\mathcal{I}}^{ab} = \int v^a v^b d\mathcal{E}_x(v). \quad (98)$$

If the settled galaxy branch is monokinetic in its rest frame, then

$$T_{\mathcal{I}}^{ab} = \rho_{\mathcal{I}} u_{\mathcal{I}}^a u_{\mathcal{I}}^b, \quad P_{\mathcal{I}} = 0, \quad q_{\mathcal{I}}^a = 0, \quad \pi_{\mathcal{I}}^{ab} = 0. \quad (99)$$

For negligible or separately modeled anisotropic stress in all other sectors, $\Phi = \Psi$.

Explicit premise. The finite collar packet state emits the positive kinetic measure, and the settled galaxy branch is monokinetic at the coarse scale used for lensing.

Conditional argument. A positive measure on future timelike velocities has a symmetric future-positive second moment. In the monokinetic branch the measure is concentrated at one four-velocity, so the second moment is $\rho_{\mathcal{I}} u_{\mathcal{I}}^a u_{\mathcal{I}}^b$. Its pressure, heat flux, and anisotropic stress vanish in that rest frame. The off-diagonal spatial Einstein equation then gives $\Phi = \Psi$ when the other sectors carry no uncompensated anisotropic stress. $\square$

Proof obligation. Derive $d\mathcal{E}_x(v)$ from the finite OPH collar packet state or from an equivalent covariant parent action.

Failure consequence. If pressure or anisotropic stress survives coarse graining, the lensing slip equation must include those terms and no-slip is regime-dependent.

15.5 Repair Relaxation Scale

Theorem 0E. Let K be the fixed-cutoff repair-commit transition matrix for one local coarse cell at fixed baryonic source. If K is finite, irreducible, and aperiodic, with spectral gap γ_{rec} , then

$$\tau_{\text{rec}} = \frac{t_{\text{commit}}}{\gamma_{\text{rec}}}. \quad (100)$$

If every allowed repair move has probability at least $q_{\text{min}} > 0$ and the comparison chain has conductance Φ_K , then

$$\gamma_{\text{rec}} \geq \frac{1}{2} q_{\text{min}} \Phi_K^2. \quad (101)$$

Proof. A finite irreducible aperiodic chain has a unique stationary state. Every nonstationary mode decays as $\exp(-\gamma_n n)$ in repair-step time, and the slowest nonzero mode has rate γ_{rec} . One physical commit step takes t_{commit} , so physical relaxation has rate $\gamma_{\text{rec}}/t_{\text{commit}}$. The conductance bound follows from the standard Cheeger comparison for the reversible comparison chain and the lower move probability q_{min} . $\square$

15.6 Environmental Kernel Around A Nonzero Field

Conditional theorem 0F. Let the settled anomaly field be

$$\mathbf{g}_A = (\nu_{\text{OPH}} - 1) \mathbf{g}_b. \quad (102)$$

Linearize around a nonzero background field $\bar{\mathbf{g}}_b$. With

$$x = |\bar{\mathbf{g}}_b|/a_{0,\text{OPH}}, \quad n_i = \bar{g}_{b,i}/|\bar{\mathbf{g}}_b|,$$

the Fourier density response to a baryonic density mode is

$$K_A^{(\rho)}(x, \mu) = \nu_{\text{OPH}}(x) - 1 + x \nu'_{\text{OPH}}(x) \mu^2, \quad (103)$$

where μ is the cosine between the mode direction and $\mathbf{n}$. The contrast kernel is

$$B_A(k, a) = \frac{\bar{\rho}_b(a)}{\bar{\rho}_A(a)} K_A^{(\rho)}(k, a). \quad (104)$$

Proof. The first variation is

$$\delta g_{A,i} = ([\nu_{\text{OPH}}(x) - 1] \delta_{ij} + x \nu'_{\text{OPH}}(x) n_i n_j) \delta g_{b,j}. \quad (105)$$

Taking the divergence and using the Fourier-space Poisson relation for $\delta \mathbf{g}_b$ gives the displayed scalar multiplier. The derivative of

$$\nu_{\text{OPH}}(x) = [1 - \exp(-\lambda_{\text{collar}} \sqrt{x})]^{-1}$$

is

$$\nu'_{\text{OPH}}(x) = -\frac{\lambda_{\text{collar}} \exp[-\lambda_{\text{collar}} \sqrt{x}]}{2\sqrt{x}[1 - \exp(-\lambda_{\text{collar}} \sqrt{x})]^2}.$$

$\square$

15.7 Background Abundance Green Function

Theorem 0G. Let the background anomaly density obey

$$\rho'_A + 3\mathcal{H}\rho_A = -a\Gamma_{\text{rec}}(\rho_A - \rho_{A,\text{eq}}). \quad (106)$$

Define the comoving densities

$$R_A = a^3\rho_A, \quad R_{A,\text{eq}} = a^3\rho_{A,\text{eq}}. \quad (107)$$

Then

$$R_A(\eta) = W(\eta, \eta_i)R_A(\eta_i) + \int_{\eta_i}^{\eta} d\eta' W(\eta, \eta')a(\eta')\Gamma_{\text{rec}}(\eta')R_{A,\text{eq}}(\eta'), \quad (108)$$

where

$$W(\eta, \eta') = \exp\left[-\int_{\eta'}^{\eta} d\tilde{\eta} a(\tilde{\eta})\Gamma_{\text{rec}}(\tilde{\eta})\right]. \quad (109)$$

Proof. Multiplying the background equation by a^3 gives

$$R'_A = -a\Gamma_{\text{rec}}(R_A - R_{A,\text{eq}}). \quad (110)$$

This is a first-order linear equation. Multiplication by the integrating factor

$$\exp\left[\int_{\eta_i}^{\eta} d\eta' a(\eta')\Gamma_{\text{rec}}(\eta')\right] \quad (111)$$

and integration gives the stated expression. $\square$

The theorem shows exactly where abundance information enters: the initial comoving anomaly load, the equilibrium source, and the repair rate. These are OPH microphysical outputs, not SPARC parameters.

15.8 Linear Relaxation Transfer

Theorem 0H. For fixed k , assume the no-slip anomaly perturbation equation

$$\delta'_A = -\theta_A + 3\Phi' - a\Gamma_{\text{rec}}q_A(\delta_A - B_A\delta_b). \quad (112)$$

Define

$$\Xi_A = a\Gamma_{\text{rec}}q_A. \quad (113)$$

Then

$$\begin{aligned} \delta_A(\eta, k) &= W_A(\eta, \eta_i)\delta_A(\eta_i, k) \\ &+ \int_{\eta_i}^{\eta} d\eta' W_A(\eta, \eta') [-\theta_A + 3\Phi' + \Xi_A B_A \delta_b]_{\eta', k}, \end{aligned} \quad (114)$$

with

$$W_A(\eta, \eta') = \exp\left[-\int_{\eta'}^{\eta} d\tilde{\eta} \Xi_A(\tilde{\eta}, k)\right]. \quad (115)$$

Proof. Move the damping term to the left:

$$\delta'_A + \Xi_A \delta_A = -\theta_A + 3\Phi' + \Xi_A B_A \delta_b. \quad (116)$$

The integrating-factor solution is the displayed expression. $\square$

This formula makes the two useful limits explicit. If $\Xi_A/\mathcal{H} \ll 1$, the anomaly evolves like pressureless free-falling matter. If $\Xi_A/\mathcal{H} \gg 1$, it tracks the OPH equilibrium source up to derivative corrections. Cluster offsets and settled galaxies require the same relaxation operator to sit between those limits.

15.9 Static-To-Linear Separation

Theorem 0I. The nonlinear static RAR law

$$g_{\text{obs}} = \frac{g_b}{1 - \exp[-\lambda_{\text{collar}} \sqrt{g_b/a_{0,\text{OPH}}}]}. \quad (117)$$

does not define the FLRW anomaly abundance or the linear CMB kernel by itself.

Proof. The RAR law is written for a static weak-field acceleration vector sourced by a resolved baryonic inhomogeneity. An exactly homogeneous FLRW background has no preferred spatial acceleration vector. Its scalar data are background densities and perturbation kernels, with no radial acceleration profile. In addition, the RAR map contains $\sqrt{g_b}$, so its first derivative is singular at $g_b = 0$ if treated as a homogeneous-background Taylor series. Therefore the static law cannot be linearized into CMB equations without an OPH parent functional. A scalar parent functional must emit $\rho_A(a)$ and $B_A(k, a)$, which belongs to the source-candidate layer. Physical Boltzmann status requires the finite covariant packet parent above, its source-provenance certificate, and pooled global reducers. $\square$

15.10 Parent Collar Functional

Scalar source target 0J. Let $\mathcal{C}_x(a)$ be the finite collar family contributing to a coarse FLRW cell x at scale factor a . Assume this finite-collar state selection has a regular FLRW limit. Let $\omega[\rho_b]$ be the OPH collar state selected by the baryonic source, and let

$$R_C[\omega] = I_\omega(A : D|B). \quad (118)$$

Define

$$\rho_{A,\text{eq}}(x, a) c^2 = \frac{15}{8\pi^2 \ell(a)^4} \int_{\mathcal{C}_x(a)} d\mu_C R_C[\omega[\rho_b]]. \quad (119)$$

Then (119) is a scalar parent collar functional that emits a background equilibrium source and a linear kernel:

$$\bar{\rho}_{A,\text{eq}}(a) = \rho_{A,\text{eq}}[\bar{\rho}_b](a), \quad (120)$$

$$K_A^{(\rho)}(k, a) = \left. \frac{\delta \widehat{\rho_{A,\text{eq}}(x, a)}}{\delta \rho_b(x', a)} \right|_{\bar{\rho}_b}(k), \quad B_A(k, a) = \frac{\rho_b(a)}{\rho_A(a)} K_A^{(\rho)}(k, a). \quad (121)$$

Conditional argument. The collar measure, the conditional mutual information, and the selected collar state are quotient-local scalar data. The integral is therefore a scalar density on the coarse FLRW cell. Evaluating the functional on a homogeneous baryonic density gives the background equilibrium source. Taking the first functional derivative around that homogeneous state gives a translation-invariant and rotation-invariant linear response kernel. Its Fourier transform is $K_A^{(\rho)}(k, a)$, and division by the background density gives $B_A(k, a)$. $\square$

The finite-collar evaluation of (119) is the mathematical target for replacing a fitted dark-matter abundance, but it does not determine the full fluid closure. The scalar parent leaves open w_A , $c_{s,A}^2$, σ_A , Q_A^μ , Γ_{rec} , recipient stress, and causal response. Those data must come from the finite covariant packet parent before the scalar tables can feed a physical Boltzmann module.

Proof obligation. Evaluate the collar measure and selected state from finite OPH screen/collar data in the homogeneous, perturbative, galaxy, and cluster regimes; lift the result to a finite covariant packet parent; prove exact stress-energy closure, gauge independence of B_A , causal/stable response, refinement convergence, and CDM-limit recovery.

15.11 Finite Repair Matrix

Theorem 0K. Let $\mathbf{S}$ be the finite packet-state space declared by one fixed-cutoff overlap interface. Let $\pi_{\text{eq}}(s|\rho_b)$ be the equilibrium packet distribution emitted by the parent collar functional. Let $q(s, s')$ be an inverse-closed local proposal rule generated by the declared repair menu, with $q(s, s') > 0$ iff $q(s', s) > 0$. Define, for $s \neq s'$,

$$K(s, s') = q(s, s') \min\left(1, \frac{\pi_{\text{eq}}(s'|\rho_b)q(s', s)}{\pi_{\text{eq}}(s|\rho_b)q(s, s')}\right), \quad (122)$$

and

$$K(s, s) = 1 - \sum_{s' \neq s} K(s, s'). \quad (123)$$

Then K is a finite stochastic repair transition matrix with stationary distribution π_{eq} . If the proposal graph is connected and some rejection or hold probability is nonzero, K is irreducible and aperiodic.

Proof. Each row sums to one by construction, so K is stochastic. For $s \neq s'$, inverse-closed proposals make both directions available. Hence

$$\pi_{\text{eq}}(s)K(s, s') = \min(\pi_{\text{eq}}(s)q(s, s'), \pi_{\text{eq}}(s')q(s', s)) = \pi_{\text{eq}}(s')K(s', s). \quad (124)$$

Detailed balance proves stationarity. Connected proposal support gives irreducibility, and a nonzero hold probability gives aperiodicity. $\square$

This construction declares K from OPH data: packet states, local proposals, and the parent equilibrium distribution.

Scalar-load diagnostic. The executable packet diagnostic takes

$$\mathbf{S} = \{0, 1, \dots, 40\}, \quad \pi_{\text{eq}}(n) = \text{Pois}(n; 5.363470441)$$

with truncation at 40, nearest-neighbor inverse-closed proposals, and hold probability 0.25. The resulting matrix has row-sum error 1.110×10^{-16} , detailed-balance error 6.939×10^{-18} , and

$$\gamma_{\text{rec}} = 0.050717471.$$

This diagnostic is a finite packet schema. The OPH task is to derive that schema, or its replacement, from the finite collar states and then attach the stress, recipient, response, convergence, and frozen-likelihood receipts needed for physical transfer.

15.12 Finite Repair-Graph Gap

Theorem 0L. For a declared fixed-cutoff repair graph with transition matrix K , the relaxation time used by clusters and perturbations is computable from the second eigenvalue. If K is reversible with stationary distribution π_{eq} , then

$$\gamma_{\text{rec}} = 1 - \lambda_2(K) \quad (125)$$

in discrete repair-step units, where λ_2 is the largest nontrivial eigenvalue in absolute value on the mean-zero subspace. For continuous-time generator $L = K - \mathbf{1}$, the gap is the smallest positive eigenvalue of $-L$.

Proof. Reversibility makes K self-adjoint in $L^2(\pi_{\text{eq}})$. The stationary vector has eigenvalue one. Every mean-zero mode decomposes into eigenvectors and decays by its eigenvalue per repair step. The slowest decaying nonstationary mode is therefore the largest nontrivial eigenvalue, giving $\gamma_{\text{rec}} = 1 - \lambda_2(K)$. The continuous-time statement follows by exponentiating the generator. $\square$

This is an exact finite-matrix target for the repair/commit microphysics. A cluster timescale is a prediction only after this matrix or its OPH state functional is supplied.

15.13 Jacobi Repair Clock

Theorem target 0M. Assume repair timing is set by the overlap-visible Jacobi mismatch rate of transported information-defect records. Let $E_{ij} = C_{0i0j}$ be the electric Weyl tensor in the local free-fall frame, equivalently the trace-free Newtonian tidal tensor in the weak-field limit. Homogeneous FLRW Ricci focusing is excluded from this local cluster-offset clock. The scalar repair rate is

$$\Gamma_J = \left(\frac{E_{ij}E^{ij}}{6} \right)^{1/4}. \quad (126)$$

For a Newtonian point source this gives

$$\Gamma_J = \sqrt{\frac{GM}{r^3}}, \quad \tau_J = \sqrt{\frac{r^3}{GM}}. \quad (127)$$

Explicit premise. The repair-cycle clock is controlled by overlap-visible geodesic-deviation mismatch of transported information-defect records.

Conditional argument. The relative acceleration of neighboring transported records is controlled by the Jacobi operator. The local cluster-offset part is the trace-free Weyl electric tensor E_{ij} ; the homogeneous FLRW Ricci term changes the background expansion and is excluded from the local offset between collisional gas and transported record stress. A scalar clock made from E_{ij} , with no chosen direction and no extra dimensional constant, must be a power of $E_{ij}E^{ij}$. The fourth root has units of inverse time. For a point Newtonian source the eigenvalues of E_{ij} are $-2GM/r^3, GM/r^3, GM/r^3$ up to sign convention, hence $E_{ij}E^{ij} = 6(GM/r^3)^2$. The normalization in the theorem makes the point-source result exact. $\square$

Proof obligation. Derive from OPH packet microphysics that the commit clock is the overlap-visible Weyl-Jacobi mismatch clock, while pixel, de Sitter, entropy-production, local-acceleration, and Ricci-curvature clocks are excluded. The self-consistent prediction evaluates E_{ij} from the total weak-field potential determined by baryons plus information-defect stress.

Failure consequence. Without that derivation, Γ_J is a no-fit dynamic diagnostic rather than a repair-clock theorem.

Static-clock boundary. The static equilibrium equation contains $\rho_{A,\text{eq}}$ but no relaxation rate. Replacing Γ_{rec} by any positive function leaves the same settled solution. The Jacobi clock is therefore a dynamic OPH premise and has no theorem status inside the static RAR law.

15.14 Homogeneous Anomaly State Selection

Theorem target 0N. Under flat capacity-saturated FLRW state selection, the homogeneous anomaly abundance is

$$\Omega_{A0} = 1 - \Omega_{\Lambda,\text{OPH}} - \Omega_{b0} - \Omega_{\nu0} - \Omega_{r0}, \quad \Omega_{\Lambda,\text{OPH}} = \left(\frac{H_{\text{dS}}}{H_0} \right)^2. \quad (128)$$

If the homogeneous anomaly charge is transported as dust with no homogeneous repair exchange, then

$$\rho_A(a) = \rho_{A0} a^{-3}. \quad (129)$$

Explicit premise. The homogeneous state is flat, capacity-saturated, and selected by a residual Friedmann constraint after H_0 , baryons, neutrinos, and radiation are supplied.

Conditional argument. The flat Friedmann equation at $a = 1$ is

$$1 = \Omega_{\Lambda,\text{OPH}} + \Omega_{b0} + \Omega_{\nu0} + \Omega_{r0} + \Omega_{A0}. \quad (130)$$

Solving for Ω_{A0} gives the displayed residual. Pressureless homogeneous transport with zero homogeneous exchange obeys

$$\rho'_A + 3\mathcal{H}\rho_A = 0,$$

whose solution is $\rho_A(a) = \rho_{A0} a^{-3}$. $\square$

Proof obligation. Derive the conserved homogeneous information-defect charge Q_A from OPH screen state data, or declare it as cosmological state data.

Failure consequence. The CMB dark abundance is conditional if Q_A is not selected by OPH microphysics. A nonzero Γ_{rec} branch is also conditional unless the same finite parent exposes the recipient stress that balances the exchange.

Capacity boundary. The screen capacity fixes the dimensionless de Sitter capacity relation; the SI de Sitter scale additionally uses the selected OPH scale certificate. After H_0 and that scale display are supplied, it fixes $\Omega_{\Lambda, \text{OPH}}$. It does not by itself fix the baryon density, the neutrino density, the radiation density, or a conserved homogeneous anomaly charge. A standalone OPH cosmology needs a theorem of the form

$$Q_A = F_{\text{OPH}}(\text{screen capacity}, P, \text{screen state data})$$

or an explicitly declared state-selection premise such as (74).

16 Falsifiers

The OPH dark branch fails if any of the following happens:

1. OPH collar/cut microphysics cannot derive the activation law $p(x) = 1 - \exp[-\lambda_{\text{collar}}\sqrt{x}]$, or a replacement law that passes the same galaxy tests.
2. The carried galaxy defect is not positive or not attractive.
3. The settled anomaly stress cannot give $\Phi = \Psi$.
4. A full SPARC likelihood rejects both the unit branch and every independently derived coefficient branch under the same nuisance treatment used for comparison models.
5. Cluster lensing offsets require a relaxation law incompatible with old settled-galaxy RAR.
6. CMB peaks, CMB lensing, BAO, weak lensing, or growth reject every frozen finite covariant parent whose Γ_{rec} , $B_A(k, a)$, $\rho_A(a)$, stress closure, recipient channel, and CDM-limit recovery remain compatible with the galaxy branch.

17 Reproducibility

The numerical tables were produced in the OPH cosmology workspace by:

```
python3 cosmology/scripts/dark_sector_measurement_ledger.py
python3 cosmology/scripts/dark_empirical_scorecard.py --quiet
python3 cosmology/scripts/sparc_systematic_likelihood.py
python3 cosmology/scripts/dark_perturbation_parameters.py
python3 cosmology/scripts/dark_repair_transition_matrix.py
python3 cosmology/scripts/dark_cluster_boltzmann_prelikelihood.py
python3 cosmology/scripts/dark_jacobi_repair_clock.py
python3 cosmology/scripts/dark_homogeneous_state_selection.py
python3 cosmology/scripts/dark_parent_collar_grid.py --diagnostic-scalar-load \
--mu-eq 5.363470440729118 \
--out cosmology/outputs/dark_parent_flat_capacity_state_selection.json
python3 cosmology/scripts/dark_cmb_bao_growth_s8_likelihood.py \
--parent-grid cosmology/outputs/dark_parent_flat_capacity_state_selection.json
python3 cosmology/scripts/camb_fixed_neutrino_compare.py
```

The SPARC data files used are public measurement tables from the SPARC page [7].

References

- [1] B. Mueller, A. Osika, M. Ponder, K. Xue, P. Nguyen, and M. A. Visser, *Recovering Relativity and the Standard Model from Observer Overlap Consistency*, OPH compact paper, release r1411, May 3, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/recovering_relativity_and_standard_model_structure_from_observer_overlap_consistency_compact.pdf
- [2] B. Mueller, A. Osika, M. Ponder, K. Xue, B. Cassie, P. Nguyen, M. A. Visser, and K. A. Anirudha, *Observers Are All You Need*, OPH synthesis paper, release r1411, May 3, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/observers_are_all_you_need.pdf
- [3] B. Mueller, A. Osika, K. Xue, B. Cassie, and M. A. Visser, *Federated Echoshedral Screen Microphysics: Patch Hardware, Records, and Observer Synchronization in OPH*, OPH microphysics paper. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/screen_microphysics_and_observer_synchronization.pdf
- [4] B. Mueller, A. Osika, M. Ponder, K. Xue, and M. A. Visser, *Deriving the Particle Zoo from Observer Consistency*, OPH particle paper, release r1411, May 3, 2026. https://github.com/FloatingPragma/observer-patch-holography/blob/main/paper/deriving_the_particle_zoo_from_observer_consistency.pdf
- [5] Planck Collaboration, *Planck 2018 results. VI. Cosmological parameters*, arXiv:1807.06209. <https://arxiv.org/abs/1807.06209>
- [6] F. Lelli, S. S. McGaugh, and J. M. Schombert, *SPARC: Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves*, Astron. J. 152, 157, 2016, arXiv:1606.09251. <https://arxiv.org/abs/1606.09251>
- [7] F. Lelli, S. S. McGaugh, and J. M. Schombert, SPARC public data page. <https://astroweb.cwru.edu/SPARC/>
- [8] S. S. McGaugh, F. Lelli, and J. M. Schombert, *The Radial Acceleration Relation in Rotationally Supported Galaxies*, Phys. Rev. Lett. 117, 201101, 2016, arXiv:1609.05917. <https://arxiv.org/abs/1609.05917>
- [9] P. Li, F. Lelli, S. S. McGaugh, and J. M. Schombert, *Fitting the radial acceleration relation to individual SPARC galaxies*, Astron. Astrophys. 615, A3, 2018, arXiv:1803.00022. <https://arxiv.org/abs/1803.00022>
- [10] M. Milgrom, *A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis*, Astrophys. J. 270, 365, 1983. <https://ui.adsabs.harvard.edu/abs/1983ApJ...270..365M/abstract>
- [11] J. D. Bekenstein, *Relativistic gravitation theory for the modified Newtonian dynamics paradigm*, Phys. Rev. D 70, 083509, 2004. <https://doi.org/10.1103/PhysRevD.70.083509>
- [12] D. Clowe, M. Bradac, A. H. Gonzalez, M. Markevitch, S. W. Randall, C. Jones, and D. Zaritsky, *A direct empirical proof of the existence of dark matter*, Astrophys. J. Lett. 648, L109, 2006, arXiv:astro-ph/0608407. <https://arxiv.org/abs/astro-ph/0608407>