

Pellicer Creek Aquatic Preserve

SEACAR Water Quality Analysis

Last compiled on 30 September, 2025

Contents

Indicators	2
Nutrients	2
Total Nitrogen - Discrete	2
Total Phosphorus - Discrete	4
Water Quality	6
Dissolved Oxygen - Discrete	6
Dissolved Oxygen - Continuous	8
Dissolved Oxygen Saturation - Discrete	10
Dissolved Oxygen Saturation - Continuous	12
Salinity - Discrete	14
Salinity - Continuous	16
Water Temperature - Discrete	18
Water Temperature - Continuous	20
pH - Discrete	22
pH - Continuous	24
Water Clarity	26
Turbidity - Discrete	26
Turbidity - Continuous	28
Total Suspended Solids - Discrete	30
Chlorophyll a, Uncorrected for Pheophytin - Discrete	32
Chlorophyll a, Corrected for Pheophytin - Discrete	34
Secchi Depth - Discrete	36
Colored Dissolved Organic Matter - Discrete	38

Indicators

Nutrients

Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1971	22	2002 - 2025	0.638	-0.00864	0.6359	-0.00046	0.8722

Total nitrogen showed no detectable trend between 2002 and 2025.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	3679	24	2002 - 2025	0.081	-0.17588	0.08735	-0.00066	1e-04

Monthly average total phosphorus decreased by less than 0.01 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Quality

Dissolved Oxygen - Discrete




Figure 5: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 3: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	2199	24	2002 - 2025	5.63	-0.01079	5.14493	-0.00194	0.8433

Dissolved oxygen showed no detectable trend between 2002 and 2025.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen - Continuous




Figure 7: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 4: Seasonal Kendall-Tau Results - Dissolved Oxygen

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
gtmpcwq	No significant trend	721785	25	2001 - 2025	5.5	0.05	5.24	0	0.251

No detectable change in monthly average dissolved oxygen was observed at one location.




Figure 8: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Discrete




Figure 9: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 5: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	201	11	2004 - 2023	55.5	0.08238	60.80703	0.14922	0.6126

Dissolved oxygen saturation showed no detectable trend between 2004 and 2023.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 11: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 6: Seasonal Kendall-Tau Results - Dissolved Oxygen Saturation

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
gtmpcwq	No significant trend	723278	25	2001 - 2025	71.4	0.01	67.88	0.02	0.6851

No detectable change in monthly average dissolved oxygen saturation was observed at one location.




Figure 12: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Discrete




Figure 13: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 7: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	No significant trend	3012	24	2002 - 2025	16.2	-0.03621	15.92762	-0.06274	0.4357

Salinity showed no detectable trend between 2002 and 2025.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 15: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 8: Seasonal Kendall-Tau Results - Salinity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
gtmpcwq	Significantly decreasing trend	731565	25	2001 - 2025	16.6	-0.14	17.1	-0.21	0.0013

At one program location, monthly average salinity decreased by 0.21 ppt per year.




Figure 16: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Discrete




Figure 17: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 9: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	2353	24	2002 - 2025	23.8	0.07972	24.04755	0.04074	0.0648

Water temperature showed no detectable trend between 2002 and 2025.




Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 19: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 10: Seasonal Kendall-Tau Results - Water Temperature

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
gtmpcwq	Significantly increasing trend	740131	25	2001 - 2025	24.4	0.17	23.61	0.05	0

At one program location, monthly average water temperature increased by 0.05°C per year.




Figure 20: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Discrete




Figure 21: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 11: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	2261	24	2002 - 2025	7.29	-0.0957	7.4608	-0.01043	0.0306

Monthly average pH decreased by 0.01 pH units per year.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 23: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 12: Seasonal Kendall-Tau Results - pH

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
gtmpcwq	Significantly decreasing trend	716012	25	2001 - 2025	7.4	-0.1	7.44	0	0.0191

At one program location, monthly average pH decreased by less than 0.01 pH units per year.




Figure 24: Map showing location of ph continuous water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Clarity

Turbidity - Discrete




Figure 25: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 13: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Insufficient data to calculate trend	237	9	2004 - 2021	7	-	-	-	-

There was insufficient data to fit a model for turbidity.




Figure 26: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Continuous




Figure 27: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 14: Seasonal Kendall-Tau Results - Turbidity

Program Location	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
gtmpcwq	Significantly decreasing trend	707002	25	2001 - 2025	9	-0.17	9.57	-0.11	1e-04

At one program location, monthly average turbidity decreased by 0.11 NTU per year.




Figure 28: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Total Suspended Solids - Discrete




Figure 29: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 15: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1868	16	2005 - 2025	14	-0.00865	15.5404	-0.02778	0.8867

Total suspended solids showed no detectable trend between 2005 and 2025.




Figure 30: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 31: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 16: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	3634	24	2002 - 2025	6.9	-0.03708	7.3778	-0.02645	0.3883

Chlorophyll a, uncorrected for pheophytin, showed no detectable trend between 2002 and 2025.




Figure 32: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 33: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 17: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	4554	24	2002 - 2025	5.2	0.02955	5.36244	0.02275	0.4906

Chlorophyll a, corrected for pheophytin, showed no detectable trend between 2002 and 2025.




Figure 34: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Secchi Depth - Discrete




Figure 35: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 18: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	359	17	2002 - 2022	-0.6	0.12513	-0.6937	0.00556	0.0435

Monthly average Secchi depth became shallower by 0.01 m per year, indicating a decrease in water clarity.




Figure 36: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Colored Dissolved Organic Matter - Discrete




Figure 37: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 19: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	959	11	2007 - 2025	54.6	-0.06685	63.35181	-0.44077	0.3172

Colored dissolved organic matter showed no detectable trend between 2007 and 2025.




Figure 38: Map showing location of discrete water quality sampling locations within the boundaries of *Pellicer Creek Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.