

Coupon Bight Aquatic Preserve

SEACAR Habitat Analyses

Last compiled on 22 May, 2025

Contents

Funding & Acknowledgements	2
Threshold Filtering	2
Value Qualifiers	3
Water Column	5
Seasonal Kendall-Tau Analysis	5
Water Quality - Discrete	5
Chlorophyll a, Uncorrected for Pheophytin - Discrete	6
Dissolved Oxygen - Discrete	7
Dissolved Oxygen Saturation - Discrete	9
Salinity - Discrete	11
Total Nitrogen - Discrete	13
Total Phosphorus - Discrete	16
Turbidity - Discrete	18
Water Temperature - Discrete	20
Water Quality - Continuous	23
Dissolved Oxygen - Continuous	25
Dissolved Oxygen Saturation - Continuous	27
pH - Continuous	29
Salinity - Continuous	31
Turbidity - Continuous	33
Water Temperature - Continuous	35
Coral Reef	37
References	39

Funding & Acknowledgements

The data used in this analysis is from the Export Standardized Tables in the SEACAR Data Discovery Interface (DDI). Documents and information available through the SEACAR DDI are owned by the data provider(s) and users are expected to provide appropriate credit following accepted citation formats. Users are encouraged to access data to maximize utilization of gained knowledge, reducing redundant research and facilitating partnerships and scientific innovation.

With respect to documents and information available from SEACAR DDI, neither the State of Florida nor the Florida Department of Environmental Protection makes any warranty, expressed or implied, including the warranties of merchantability and fitness for a particular purpose arising out of the use or inability to use the data, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

This report was funded in part, through a grant agreement from the Florida Department of Environmental Protection, Florida Coastal Management Program, by a grant provided by the Office for Coastal Management under the Coastal Zone Management Act of 1972, as amended, National Oceanic and Atmospheric Administration. The views, statements, findings, conclusions and recommendations expressed herein are those of the author(s) and do not necessarily reflect the views of the State of Florida, NOAA or any of their sub agencies.

Published: 2025-05-22



Threshold Filtering

Threshold filters, following the guidance of Florida Department of Environmental Protection's (*FDEP*) Division of Environmental Assessment and Restoration (*DEAR*) are used to exclude specific results values from the SEACAR Analysis. Based on the threshold filters, Quality Assurance / Quality Control (*QAQC*) Flags are inserted into the *SEACAR_QAQCFlagCode* and *SEACAR_QAQC_Description* columns of the export data. The *Include* column indicates whether the *QAQC* Flag will also indicate that data are excluded from analysis. No data are excluded from the data export, but the analysis scripts can use the *Include* column to exclude data (1 to include, 0 to exclude).

Table 1: Continuous Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Dissolved Oxygen	mg/L	-0.000001	50
Dissolved Oxygen Saturation	%	-0.000001	500
Salinity	ppt	-0.000001	70
Turbidity	NTU	-0.000001	4000
Water Temperature	Degrees C	-5.000000	45
pH	None	2.000000	14

Table 2: Discrete Water Quality threshold values

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Ammonia, Un-ionized (NH3)	mg/L	-	-
Ammonium, Filtered (NH4)	mg/L	-	-
Chlorophyll a, Corrected for Pheophytin	ug/L	-	-
Chlorophyll a, Uncorrected for Pheophytin	ug/L	-	-
Colored Dissolved Organic Matter	PCU	-	-

<i>Parameter Name</i>	<i>Units</i>	<i>Low Threshold</i>	<i>High Threshold</i>
Dissolved Oxygen	mg/L	-0.000001	25
Dissolved Oxygen Saturation	%	-0.000001	310
Fluorescent Dissolved Organic Matter	QSE	-	-
Light Extinction Coefficient	m ⁻¹	-	-
NO ₂ +3, Filtered	mg/L	-	-
Nitrate (NO ₃)	mg/L	-	-
Nitrite (NO ₂)	mg/L	-	-
Nitrogen, organic	mg/L	-	-
Phosphate, Filtered (PO ₄)	mg/L	-	-
Salinity	ppt	-0.000001	70
Secchi Depth	m	0.000001	50
Specific Conductivity	mS/cm	0.005000	100
Total Kjeldahl Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Nitrogen	mg/L	-	-
Total Phosphorus	mg/L	-	-
Total Suspended Solids	mg/L	-	-
Turbidity	NTU	-	-
Water Temperature	Degrees C	3.000000	40
pH	None	2.000000	13

Table 3: Quality Assurance Flags inserted based on threshold checks listed in Table 1 and 2

<i>SEACAR QAQC Description</i>	<i>Include</i>	<i>SEACAR QAQCFlagCode</i>
Exceeds maximum threshold	0	2Q
Below minimum threshold	0	4Q
Within threshold tolerance	1	6Q
No defined thresholds for this parameter	1	7Q

Value Qualifiers

Value qualifier codes included within the data are used to exclude certain results from the analysis. The data are retained in the data export files, but the analysis uses the *Include* column to filter the results.

STORET and WIN value qualifier codes

Value qualifier codes from *STORET* and *WIN* data are examined with the database and used to populate the *Include* column in data exports.

Table 4: Value Qualifier codes excluded from analysis

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>MDL</i>	<i>Description</i>
STORET-WIN	H	0	0	Value based on field kit determination; results may not be accurate
STORET-WIN	J	0	0	Estimated value
STORET-WIN	V	0	0	Analyte was detected at or above method detection limit
STORET-WIN	Y	0	0	Lab analysis from an improperly preserved sample; data may be inaccurate

Discrete Water Quality Value Qualifiers

The following value qualifiers are highlighted in the Discrete Water Quality section of this report. An exception is made for **Program 476 - Charlotte Harbor Estuaries Volunteer Water Quality Monitoring Network** and data flagged with Value Qualifier **H** are included for this program only.

H - Value based on field kit determination; results may not be accurate. This code shall be used if a field screening test (e.g., field gas chromatograph data, immunoassay, or vendor-supplied field kit) was used to generate the value and the field kit or method has not been recognized by the Department as equivalent to laboratory methods.

I - The reported value is greater than or equal to the laboratory method detection limit but less than the laboratory practical quantitation limit.

Q - Sample held beyond the accepted holding time. This code shall be used if the value is derived from a sample that was prepared or analyzed after the approved holding time restrictions for sample preparation or analysis.

S - Secchi disk visible to bottom of waterbody. The value reported is the depth of the waterbody at the location of the Secchi disk measurement.

U - Indicates that the compound was analyzed for but not detected. This symbol shall be used to indicate that the specified component was not detected. The value associated with the qualifier shall be the laboratory method detection limit. Unless requested by the client, less than the method detection limit values shall not be reported

Systemwide Monitoring Program (SWMP) value qualifier codes

Value qualifier codes from the *SWMP* continuous program are examined with the database and used to populate the *Include* column in data exports. *SWMP* Qualifier Codes are indicated by *QualifierSource=SWMP*.

Table 5: SWMP Value Qualifier codes

<i>Qualifier Source</i>	<i>Value Qualifier</i>	<i>Include</i>	<i>Description</i>
SWMP	-1	1	Optional parameter not collected
SWMP	-2	0	Missing data
SWMP	-3	0	Data rejected due to QA/QC
SWMP	-4	0	Outside low sensor range
SWMP	-5	0	Outside high sensor range
SWMP	0	1	Passed initial QA/QC checks
SWMP	1	0	Suspect data
SWMP	2	1	Reserved for future use
SWMP	3	1	Calculated data: non-vented depth/level sensor correction for changes in barometric pressure
SWMP	4	1	Historical: Pre-auto QA/QC
SWMP	5	1	Corrected data

Water Column

The water column habitat extends from the water's surface to the bottom sediments, and it's where fish, dolphins, crabs and people swim! So much life makes its home in the water column that the health of marine and coastal ecosystems, as well as human economies, depend on the condition of this vulnerable habitat. Local patterns of rainfall, temperature, winds and currents can rapidly change the condition of the water column, while global influences such as [El Niño/La Niña](#), large-scale fluctuation in sea temperatures and climate change can have long-term effects. Inputs from the prosperity of our day-to-day lives including farming, mining and forestry, and emissions from power generation, automobiles and water treatment can also alter the health of the water column. Acting alone or together, each input can have complex and lasting effects on habitats and ecosystems.

SEACAR evaluates water column health with several essential parameters. These include nutrient surveys of nitrogen and phosphorus, and water quality assessments of salinity, dissolved oxygen, pH, and water temperature. Water clarity is evaluated with Secchi depth, turbidity, levels of chlorophyll a, total suspended solids, and colored dissolved organic matter. Additionally, the richness of nekton is indicated by the abundance of free-swimming fishes and macroinvertebrates like crabs and shrimps.

Seasonal Kendall-Tau Analysis

Indicators must have a minimum of five to ten years, depending on the habitat, of data within the geographic range of the analysis to be included in the analysis. Ten years of data are required for discrete parameters, and five years of data are required for continuous parameters. If there are insufficient years of data, the number of years of data available will be noted and labeled as “insufficient data to conduct analysis”. Further, for the preferred Seasonal Kendall-Tau test, there must be data from at least two months in common across at least two consecutive years within the RCP managed area being analyzed. Values that pass both of these tests will be included in the analysis and be labeled as *Use_In_Analysis* = **TRUE**. Any that fail either test will be excluded from the analyses and labeled as *Use_In_Analysis* = **FALSE**. The points for all Water Column plots displayed in this section are monthly averages. Trend significance will be denoted as “Significant Trend” (when $p < 0.05$), or “Non-significant Trend” (when $p \geq 0.05$). Any parameters with insufficient data to perform Seasonal Kendall-Tau test will have their monthly averages plotted without a corresponding trend line.

Water Quality - Discrete

The following files were used in the discrete analysis:

- *Combined_WQ_WC_NUT_Chlorophyll_a_corrected_for_pheophytin-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Chlorophyll_a_uncorrected_for_pheophytin-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Colored_dissolved_organic_matter_CDOM-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Dissolved_Oxygen_Saturation-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_pH-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Salinity-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Secchi_Depth-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Total_Nitrogen-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Total_Phosphorus-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Total_Suspended_Solids_TSS-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Turbidity-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_Water_Temperature-2025-Mar-06.txt*

Chlorophyll a, Uncorrected for Pheophytin - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 1: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 6: Seasonal Kendall-Tau Trend Analysis for Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	Significantly increasing trend	184	23	1995 - 2024	0.6	0.435	0.0325	0.0189	0

Monthly average chlorophyll a, uncorrected for pheophytin, increased by 0.02 $\mu\text{g/L}$ per year, indicating a decrease in water clarity.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 7: Programs contributing data for Chlorophyll a, Uncorrected for Pheophytin

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	93	2019	2024
297	61	1995	2011
103	32	2020	2021

Program names:

103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX¹

297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²

5002 - Florida STORET / WIN³

Dissolved Oxygen - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 3: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 8: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	Significantly decreasing trend	159	20	1995 - 2024	5.9905	-0.2504	6.5239	-0.039	0.022

Monthly average dissolved oxygen decreased by 0.04 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 9: Programs contributing data for Dissolved Oxygen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
297	111	1995	2011
5002	33	2020	2024
118	7	2020	2020
69	4	2000	2000
103	4	2021	2021

Program names:

69 - Fisheries-Independent Monitoring (FIM) Program⁴
 103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX¹
 118 - National Aquatic Resource Surveys, National Coastal Condition Assessment⁵
 297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²
 5002 - Florida STORET / WIN³

Dissolved Oxygen Saturation - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 5: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 10: Seasonal Kendall-Tau Trend Analysis for Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	No significant trend	192	22	1995 - 2024	93.6199	-0.018	94.3848	-0.0947	0.7714

Dissolved oxygen saturation showed no detectable trend between 1995 and 2024.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 11: Programs contributing data for Dissolved Oxygen Saturation

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
297	112	1995	2011
5002	85	2020	2024

Program names:

297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²

5002 - Florida STORET / WIN³

Salinity - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 7: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 12: Seasonal Kendall-Tau Trend Analysis for Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
All	Significantly increasing trend	202	22	1995 - 2024	36.7	0.3294	36.2011	0.0415	0

Monthly average salinity increased by 0.04 ppt per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 13: Programs contributing data for Salinity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
297	111	1995	2011
5002	86	2020	2024
118	6	2020	2020
69	4	2000	2000

Program names:

69 - Fisheries-Independent Monitoring (FIM) Program⁴

118 - National Aquatic Resource Surveys, National Coastal Condition Assessment⁵

297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²

5002 - Florida STORET / WIN³

Total Nitrogen - Discrete

Total Nitrogen Calculation:

The logic for calculated Total Nitrogen was provided by Kevin O'Donnell and colleagues at FDEP (with the help of Jay Silvanima, Watershed Monitoring Section). The following logic is used, in this order, based on the availability of specific nitrogen components.

- 1) $TN = TKN + NO_3O_2$;
- 2) $TN = TKN + NO_3 + NO_2$;
- 3) $TN = ORGN + NH_4 + NO_3O_2$;
- 4) $TN = ORGN + NH_4 + NO_2 + NO_3$;
- 5) $TN = TKN + NO_3$;
- 6) $TN = ORGN + NH_4 + NO_3$;

Additional Information:

- Rules for use of sample fraction:
 - Florida Department of Environmental Protection (FDEP) report that if both “Total” and “Dissolved” components are reported, only “Total” is used. If the total is not reported, then the dissolved components are used as a best available replacement.
 - Total nitrogen calculations are done using nitrogen components with the same sample fraction, nitrogen components with mixed total/dissolved sample fractions are not used. In other words, total nitrogen can be calculated when TKN and NO₃O₂ are both total sample fractions, or when both are dissolved sample fractions. *Future calculations of total nitrogen values may be based on components with mixed sample fractions.*
- Values inserted into data:
 - ParameterName = “Total Nitrogen”
 - SEACAR_QAQCFlagCode = “1Q”
 - SEACAR_QAQC_Description = “SEACAR Calculated”

Seasonal Kendall-Tau Trend Analysis




Figure 9: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 14: Seasonal Kendall-Tau Trend Analysis for Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	Significantly increasing trend	163	23	1995 - 2024	0.2887	0.3505	0.1608	0.0068	0

Monthly average total nitrogen increased by 0.01 mg/L per year.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 15: Programs contributing data for Total Nitrogen

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	99	2019	2024
297	64	1995	2011

Program names:

297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²

5002 - Florida STORET / WIN³

Total Phosphorus - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 11: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 16: Seasonal Kendall-Tau Trend Analysis for Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	Significantly increasing trend	180	23	1995 - 2024	0.007	0.2005	0.0062	0.0001	0.0081

Monthly average total phosphorus increased by less than 0.01 mg/L per year.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 17: Programs contributing data for Total Phosphorus

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	102	2019	2024
297	64	1995	2011
103	16	2020	2021

Program names:

103 - EPA STOrage and RETrieval Data Warehouse (STORET)/WQX¹

297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²

5002 - Florida STORET / WIN³

Turbidity - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 13: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 18: Seasonal Kendall-Tau Trend Analysis for Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Lab	Insufficient data to calculate trend	97	6	2019 - 2024	0.7	-	-	-	-

There was insufficient data to fit a model for turbidity.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 19: Programs contributing data for Turbidity

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
5002	99	2019	2024
297	74	1995	2011
103	16	2020	2021

Program names:

103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX¹

297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²

5002 - Florida STORET / WIN³

Water Temperature - Discrete

Seasonal Kendall-Tau Trend Analysis




Figure 15: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 20: Seasonal Kendall-Tau Trend Analysis for Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
Field	No significant trend	242	27	1995 - 2024	28.5	0.0096	27.7916	0	1

Water temperature showed no detectable trend between 1995 and 2024.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Table 21: Programs contributing data for Water Temperature

<i>ProgramID</i>	<i>N_Data</i>	<i>YearMin</i>	<i>YearMax</i>
297	110	1995	2011
5002	98	2020	2024
982	23	2015	2023
103	13	2020	2021
69	4	2000	2000
95	1	2010	2010

Program names:

- 69 - Fisheries-Independent Monitoring (FIM) Program⁴
- 95 - Harmful Algal Bloom Marine Observation Network⁶
- 103 - EPA STORage and RETrieval Data Warehouse (STORET)/WQX¹
- 297 - Florida Keys National Marine Sanctuary Water Quality Monitoring Project²
- 982 - Florida Keys Bleach Watch⁷
- 5002 - Florida STORET / WIN³

Water Quality - Continuous

The following files were used in the continuous analysis:

- *Combined_WQ_WC_NUT_cont_Dissolved_Oxygen_SE-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Dissolved_Oxygen_Saturation_SE-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_pH_SE-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Salinity_SE-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Turbidity_SE-2025-Mar-06.txt*
- *Combined_WQ_WC_NUT_cont_Water_Temperature_SE-2025-Mar-06.txt*

Continuous monitoring locations in Coupon Bight Aquatic Preserve

Table 22: Station overview for Continuous parameters by Program

<i>ProgramID</i>	<i>ProgramLocationID</i>	<i>Years of Data</i>	<i>Use in Analysis</i>	<i>Parameters</i>
10004	FKCB	1	FALSE	DO , DOS , pH , Sal , Turb , TempW

Program names:

10004 - Florida Keys Aquatic Preserves Continuous Water Quality Monitoring⁸




Figure 17: Map showing continuous water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. Sites marked as *Use In Analysis* (green) are featured in this report.

Dissolved Oxygen - Continuous




Figure 18: Scatter plot of monthly average dissolved oxygen over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 23: Seasonal Kendall-Tau Results for Dissolved Oxygen - All Stations

Station	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
FKCB	Insufficient data to calculate trend	16262	1	2024 - 2024	6.8	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 19: Map showing location of dissolved oxygen continuous water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Dissolved Oxygen Saturation - Continuous




Figure 20: Scatter plot of monthly average dissolved oxygen saturation over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 24: Seasonal Kendall-Tau Results for Dissolved Oxygen Saturation - All Stations

Station	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
FKCB	Insufficient data to calculate trend	16263	1	2024 - 2024	103.3	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 21: Map showing location of dissolved oxygen saturation continuous water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

pH - Continuous




Figure 22: Scatter plot of monthly average pH over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 25: Seasonal Kendall-Tau Results for pH - All Stations

Station	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
FKCB	Insufficient data to calculate trend	16263	1	2024 - 2024	8.2	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 23: Map showing location of pH continuous water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Salinity - Continuous




Figure 24: Scatter plot of monthly average salinity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 26: Seasonal Kendall-Tau Results for Salinity - All Stations

Station	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
FKCB	Insufficient data to calculate trend	16258	1	2024 - 2024	36.5	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 25: Map showing location of salinity continuous water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Turbidity - Continuous




Figure 26: Scatter plot of monthly average turbidity over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 27: Seasonal Kendall-Tau Results for Turbidity - All Stations

Station	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
FKCB	Insufficient data to calculate trend	16240	1	2024 - 2024	4	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 27: Map showing location of turbidity continuous water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Water Temperature - Continuous




Figure 28: Scatter plot of monthly average water temperature over time at continuously monitored program locations. Each location is analyzed separately, with significant (blue) or non-significant (magenta) trend lines shown for time series that included five or more years of observations.

Table 28: Seasonal Kendall-Tau Results for Water Temperature - All Stations

Station	Statistical Trend	Sample Count	Years with Data	Period of Record	Median	tau	Sen Intercept	Sen Slope	p
FKCB	Insufficient data to calculate trend	16263	1	2024 - 2024	26.8	-	-	-	-

There was insufficient data to fit a model for one location.




Figure 29: Map showing location of water temperature continuous water quality sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

Coral Reef

The data file used is: **All_CORAL_Parameters-2025-Mar-06.txt**
Species Richness




Figure 30: Line graph of annual average species richness of grazers and reef-dependent species over time. If the time series included more than one year of observations, a line connects the data points for visualization.

Table 29: Coral Species Richness

<i>Sample Count</i>	<i>Number of Years</i>	<i>Period of Record</i>	<i>Median N of Taxa</i>	<i>Mean N of Taxa</i>
72	16	1999 - 2017	281	182.125

The median annual number of taxa was 281 based on 72 observations collected between 1999 and 2017.




Figure 31: Map showing location of coral species richness sampling locations within the boundaries of *Coupon Bight Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

References

1. U.S. Environmental Protection Agency (EPA). [EPA STORage and RETrieval Data Warehouse \(STORET\)/WQX](#). (2023).
2. Florida International University (FIU). [Florida Keys National Marine Sanctuary Water Quality Monitoring Project](#). (2023).
3. Florida Department of Environmental Protection (DEP). [Florida STORET / WIN](#). (2024).
4. Florida Fish and Wildlife Conservation Commission (FWC). [Fisheries-Independent Monitoring \(FIM\) Program](#). (2022).
5. U.S. Environmental Protection Agency (EPA); Office of Water; National Oceanic and Atmospheric Administration (NOAA); U.S. Geological Survey (USGS); U.S. Fish and Wildlife Service (USFWS); National Estuary Program (NEP); coastal states. [National Aquatic Resource Surveys, National Coastal Condition Assessment](#). (2021).
6. Florida Fish and Wildlife Conservation Commission (FWC); Florida Fish and Wildlife Research Institute (FWRI). [Harmful Algal Bloom Marine Observation Network](#). (2018).
7. Mote Marine Laboratory. [Florida Keys Bleach Watch](#). (2023).
8. Florida Department of Environmental Protection (DEP); Office of Resilience and Coastal Protection (RCP); Florida Keys Aquatic Preserves. [Florida Keys Aquatic Preserves Continuous Water Quality Monitoring](#). (2024).