

# St. Andrews Aquatic Preserve

## SEACAR Water Quality Analysis

Last compiled on 10 July, 2025

### Contents

<b>Indicators</b>	<b>2</b>
Nutrients . . . . .	2
Total Nitrogen - Discrete . . . . .	2
Total Phosphorus - Discrete . . . . .	4
Water Quality . . . . .	6
Dissolved Oxygen - Discrete . . . . .	6
Dissolved Oxygen Saturation - Discrete . . . . .	8
Salinity - Discrete . . . . .	10
Water Temperature - Discrete . . . . .	12
pH - Discrete . . . . .	14
Water Clarity . . . . .	16
Turbidity - Discrete . . . . .	16
Total Suspended Solids - Discrete . . . . .	18
Chlorophyll a, Uncorrected for Pheophytin - Discrete . . . . .	20
Chlorophyll a, Corrected for Pheophytin - Discrete . . . . .	22
Secchi Depth - Discrete . . . . .	24
Colored Dissolved Organic Matter - Discrete . . . . .	26

# Indicators

## Nutrients

### Total Nitrogen - Discrete




Figure 1: Scatter plot of monthly average total nitrogen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only nitrogen values obtained from laboratory analyses (triangles) are included in the plot.

Table 1: Seasonal Kendall-Tau Results for - Total Nitrogen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	1142	33	1990 - 2024	0.18	0.34247	0.0362	0.00438	0

Monthly average total nitrogen increased by less than 0.01 mg/L per year.




Figure 2: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Total Phosphorus - Discrete




Figure 3: Scatter plot of monthly average total phosphorus over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only phosphorus values obtained from laboratory analyses (triangles) are included in the plot.

Table 2: Seasonal Kendall-Tau Results for - Total Phosphorus

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly increasing trend	985	33	1990 - 2024	0.008	0.46597	-0.00209	0.00033	0

Monthly average total phosphorus increased by less than 0.01 mg/L per year.




Figure 4: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Water Quality

### Dissolved Oxygen - Discrete




Figure 5: Scatter plot of monthly average dissolved oxygen over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen values measured in the field (circles) are included in the plot.

Table 3: Seasonal Kendall-Tau Results for - Dissolved Oxygen

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	2835	28	1996 - 2024	7.16	0.02036	7.34659	0.00274	0.6469

Dissolved oxygen showed no detectable trend between 1996 and 2024.




Figure 6: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Dissolved Oxygen Saturation - Discrete




Figure 7: Scatter plot of monthly average dissolved oxygen saturation over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only dissolved oxygen saturation values measured in the field (circles) are included in the plot.

Table 4: Seasonal Kendall-Tau Results for - Dissolved Oxygen Saturation

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly decreasing trend	901	18	2001 - 2024	101.9	-0.39775	115.5735	-0.895	0

Monthly average dissolved oxygen saturation decreased by 0.9% per year.




Figure 8: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Salinity - Discrete




Figure 9: Scatter plot of monthly average salinity over time. If the time series included ten or more years of discrete observations, significant (blue) or non-significant (magenta) trend lines are also shown. Discrete salinity values derived from grab samples analyzed in the field (circles) or the laboratory (triangles) are both included in the plot.

Table 5: Seasonal Kendall-Tau Results for - Salinity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
All	Significantly decreasing trend	2557	30	1974 - 2024	31.8	-0.09099	33.61385	-0.043	0.0378

Monthly average salinity decreased by 0.04 ppt per year.




Figure 10: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Water Temperature - Discrete




Figure 11: Scatter plot of monthly average water temperature over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only water temperature measurements taken in the field (circles) are included in the plot.

Table 6: Seasonal Kendall-Tau Results for - Water Temperature

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	2913	29	1974 - 2024	25.38	-0.0508	21.48245	-0.01606	0.2582

Water temperature showed no detectable trend between 1974 and 2024.




Figure 12: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## pH - Discrete




Figure 13: Scatter plot of monthly average pH over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only pH values measured in the field (circles) are included in the plot.

Table 7: Seasonal Kendall-Tau Results for - pH

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	No significant trend	2708	26	1998 - 2024	8.1	0.00459	8.10826	0	1

pH showed no detectable trend between 1998 and 2024.




Figure 14: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Water Clarity

### Turbidity - Discrete




Figure 15: Scatter plot of monthly average turbidity over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only turbidity values measured in the laboratory (triangles) are included in the plot.

Table 8: Seasonal Kendall-Tau Results for - Turbidity

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	622	19	2003 - 2024	0.775	-0.43163	1.09285	-0.03202	0

Monthly average turbidity decreased by 0.03 NTU per year, indicating an increase in water clarity.




Figure 16: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Total Suspended Solids - Discrete




Figure 17: Scatter plot of monthly average total suspended solids (TSS) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only TSS values obtained from laboratory analyses (triangles) are included in the plot.

Table 9: Seasonal Kendall-Tau Results for - Total Suspended Solids

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	Significantly decreasing trend	263	13	2003 - 2015	9	-0.23727	10.55284	-0.45833	0.004

Monthly average total suspended solids decreased by 0.46 mg/L per year, indicating an increase in water clarity.



Figure 18: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Chlorophyll a, Uncorrected for Pheophytin - Discrete




Figure 19: Scatter plot of monthly average levels of chlorophyll a, uncorrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 10: Seasonal Kendall-Tau Results for - Chlorophyll a, Uncorrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	1251	33	1990 - 2024		1.3	-0.07317	1.72091	-0.0055 0.1142

Chlorophyll a, uncorrected for pheophytin, showed no detectable trend between 1990 and 2024.




Figure 20: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Chlorophyll a, Corrected for Pheophytin - Discrete




Figure 21: Scatter plot of monthly average levels of chlorophyll a, corrected for pheophytin, over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed chlorophyll a (triangles) is included in the plot.

Table 11: Seasonal Kendall-Tau Results for - Chlorophyll a, Corrected for Pheophytin

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	852	19	2003 - 2024	1.2	0.0093	1.27184	0.00038	0.9261

Chlorophyll a, corrected for pheophytin, showed no detectable trend between 2003 and 2024.




Figure 22: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Secchi Depth - Discrete




Figure 23: Scatter plot of monthly average Secchi depth over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Secchi depth is only measured in the field (circles).

Table 12: Seasonal Kendall-Tau Results for - Secchi Depth

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Field	Significantly increasing trend	2186	32	1991 - 2024	-2.6	0.17014	-4.50922	0.04803	0

Monthly average Secchi depth became shallower by 0.05 m per year, indicating a decrease in water clarity.




Figure 24: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.

## Colored Dissolved Organic Matter - Discrete




Figure 25: Scatter plot of monthly average colored dissolved organic matter (CDOM) over time. If the time series included ten or more years of discrete observations, a significant (blue) or non-significant (magenta) trend line is also shown. Only laboratory-analyzed CDOM (triangles) is included in the plot.

Table 13: Seasonal Kendall-Tau Results for - Colored Dissolved Organic Matter

Activity Type	Statistical Trend	Sample Count	Years with Data	Period of Record	Median Result Value	Tau	Sen Intercept	Sen Slope	P
Lab	No significant trend	161	23	2001 - 2024	6	0.17801	3	0	0.8787

Colored dissolved organic matter showed no detectable trend between 2001 and 2024.




Figure 26: Map showing location of discrete water quality sampling locations within the boundaries of *St. Andrews Aquatic Preserve*. The bubble size on the maps above reflect the amount of data available at each sampling site.